

Original Research Article

Enhancing Zn and Fe content in Blackgram seeds by adapting drip fertigation for nutritional security

ABSTRACT

Aims: To study the influence of Zn and Fe content in blackgram seeds by adapting drip fertigation.

Study design: Randomized Block design.

Place and Duration of Study: Research block, Department of Agronomy, Agricultural College and Research Institute, Tamil Nadu Agricultural University, Madurai; March – May 2023.

Methodology: Field experiment was conducted with blackgram var.VBN 8 variety under drip irrigation on sandy clay textured soil. The treatments comprising of control measures, seed treatment, soil application, foliar spraying of Zn and Fe, a combination of seed and foliar spraying and fertigation with ZnSO₄ and FeSO₄. Fertigation with 100 per cent recommended dose of NPK was given as water soluble fertilizer for all treatments. A field-based study was executed to explore the impact of zinc sulphate and ferrous sulphate on growth, yield and grain quality of micronutrient uptake were recorded. The economics and Water Use Efficiency of different treatments were also calculated.

Results: Among the different treatments, the application of ZnSO₄ @ 25 kg ha⁻¹ and FeSO₄ @ 10 kg ha⁻¹ through drip fertigation at 15, 30 & 45 DAS recorded the highest plant growth and yield of blackgram and it also enhanced the zinc content 17.72 per cent and iron content 23.82 per cent in blackgram seeds compared to the control seeds.

Conclusion: Drip fertigation of zinc sulphate and ferrous sulphate improved the grain quality and it also used to alleviate malnutrition in humans.

Keywords: Zinc; iron; drip fertigation; blackgram.

1. INTRODUCTION

India is a major producer of pulses, contributing significantly to global production. Pulses are the cheapest sources of proteins, vitamins and micronutrients and can be supplied to people through daily diet. Despite being a prominent protein source in human diets, pulses cultivated in nutrient-depleted soils generally exhibit diminished micronutrient content within their grains, potentially leading to micronutrient deficiencies among human populations. Micronutrient malnutrition is impacting rural communities in developing nations who have limited access to a variety of foods because of their constrained purchasing ability [12]. Iron (Fe) and zinc (Zn) stand out as the most critical micronutrients for human health, and their deficiencies have widespread consequences across the globe. Zinc serves as a catalyst for essential enzymes, including ribonucleic acid (RNA) polymerase, superoxide dismutase, and

Comment [B1]: A

Comment [B2]: remove

Comment [B3]: the

Comment [B4]: Add proper space

Comment [B5]: remove

Comment [B6]: their

cellular signaling proteins. On the other hand, iron plays a vital role in proteins like hemoglobin and facilitating oxygen transport throughout the human body and activating numerous enzymes crucial for various biological functions. Among the various agricultural strategies available to combat malnutrition, agronomic bio-fortification stands out as the leading approach for enhancing the iron (Fe) and zinc (Zn) levels in grains. Applying fertilizers either directly to the soil or through foliar spraying is a convenient, efficient and low cost method to enhance the levels of Zn and Fe in the edible portions of plants. This approach is commonly referred to as agronomic bio-fortification [4].

Comment [B7]: please check

In India, blackgram holds the third position among the pulse crops, following pigeonpea and chickpea. In Tamil Nadu, blackgram cultivated in an area of 4.07 lakh hectares, 2.69 lakh tonnes production with a productivity of 660 kg per hectare [7]. Blackgram is highly susceptible to both water stress and water logging throughout its growing period. Effective management of irrigation water is an important issue in crop production, since irrigation is a precondition for crop growth, development and production per mm of water and productivity per unit area. Drip irrigation can be regarded as an efficient irrigation method because it precisely moistens the soil, ensuring that the root zone maintains an optimal moisture level. Drip fertigation is a precise and efficient agricultural practice that combines drip irrigation with the application of fertilizers and nutrients directly to the root zone of plants. In this contrast, blackgram is being considered a viable choice for augmenting micronutrient content and providing nourishing food options through drip fertigation.

Comment [B8]: s

Comment [B9]: proper space

2. MATERIAL AND METHODS

2.1. Study Area

A field experiment was conducted during summer, 2023 at field number C-46 of the research block, Agricultural College and Research Institute, Madurai, which is geographically located at 9°54' N latitude and 78°54' E longitude at an elevation of 147 m above the Mean Sea Level. The soil of the experiment site was sandy clay in texture. The main objective of this experiment was to improve the Zn and Fe content and yield of blackgram under drip fertigation.

Comment [B10]: at

2.2. Experimental Details

The experiment was laid out statistically in a Randomized Block Design with eight treatments and three replications allocated randomly to reduce the experimental error. The treatment comprising of T₁- Control (No Zn and Fe application), T₂- Soil application of RDF of ZnSO₄ @ 25kg ha⁻¹ + FeSO₄ @ 10 kg ha⁻¹ at basal, T₃- Seed treatment with ZnSO₄ @ 5 g ha⁻¹ + FeSO₄ @ 5 g ha⁻¹, T₄- Foliar spray of ZnSO₄ @ 0.5% + FeSO₄ @ 0.5% at 30 & 45DAS, T₅- Seed treatment with ZnSO₄ @ 5g ha⁻¹ + FeSO₄ @ 5g ha⁻¹ + foliar spray of ZnSO₄ @ 0.5% + FeSO₄ @ 0.5 % at 30 & 45DAS, T₆- Fertigation of ZnSO₄ @ 12.5 kg ha⁻¹ + FeSO₄ @ 5 kg ha⁻¹ at 15, 30 & 45 DAS, T₇- Fertigation of ZnSO₄ @ 18.75kg ha⁻¹ + Fe SO₄ @ 7.5 kg ha⁻¹ at 15, 30 & 45 DAS and T₈- Fertigation of ZnSO₄ @ 25 kg ha⁻¹ + FeSO₄ @ 10 kg ha⁻¹ at 15, 30 & 45 DAS. Blackgram var.VBN 8 was taken for the experiment. Raised beds were formed with the plot size of 30 x 1.2 m². Line sowing of blackgram seeds was done by adapting four rows per bed with a spacing of 30 x 10 cm. Macronutrients were supplied as water soluble fertilizer through drip fertigation for all the treatments. Irrigation was done once in 3 days based on 100 per cent potential evapo-transpiration (PET).

Comment [B11]: a

Comment [B12]: remove

2.3. Biometric observation

Five plants were chosen randomly in each plot and tagged. From the tagged plants growth and yield parameters were recorded.

2.3.1. Plant height (cm)

Plant height is the indication of crop growth. With a linear meter scale, the plant height was measured from the ground level to the tip of the last opened leaf at harvest and their average was calculated and expressed in cm.

2.3.2. Leaf Area Index (LAI)

The maximum leaf length and breadth of the third leaf from the top of the plant was measured using measuring scale and total number of green leaves was counted from the tagged plants in each treatment. The leaf area index was worked out as suggested by Puttaswamy *et al.*[14] using the formula as given below.

$$LAI = \frac{L \times B \times K \times \text{No. of leaves plant}^{-1}}{\text{Unit land area (cm}^2\text{)}}$$

Where,

- L – Maximum length of the third leaf from top (cm)
- B – Maximum breadth of the third leaf from top (cm)
- K – Constant factor (0.7)

2.3.3. Yield and yield attributes

The numbers of pods per plant was counted from five tagged plants and mean number of pods per plant was recorded. For determining grain yield, all the pods were harvested separately and threshed manually, cleaned and were dried to 12 per cent moisture level and the grain yield from net plot was calculated and expressed in kg ha⁻¹.

2.4. Analysis of Zn and Fe content in blackgram seeds

Seed samples were subjected to drying in an oven at a temperature of 60°C. Subsequently, the oven-dried seed samples were finely ground and then digested using a tri-acid solution. The micronutrient content of Zn and Fe in the seeds was quantified using an Atomic Absorption Spectrophotometer (AAS) at 213 nm and 248.33 nm respectively.

2.5. Water Use Efficiency

Water use efficiency (WUE) is the yield that can be produced from a given quantity of water Viets, [19]. It was worked out by using the following formula and expressed as kg mm⁻¹ ha⁻¹.

$$WUE = \frac{\text{Economic yield of crop (kg ha}^{-1}\text{)}}{\text{Total water used (mm)}}$$

2.6. Economics

To assess the economic returns of the each treatments, the following parameters were computed, they are Cost of cultivation: The expenditure incurred from field preparation to harvest was worked out and expressed as Rs ha⁻¹. Gross return: The crop yield was computed per hectare and the total income was worked out based on the minimum market rate which was prevalent during the time of this study. Net return: Net returns were obtained by subtracting the cost of cultivation from gross return for each treatment. Net return = Gross return (Rs. ha⁻¹) - Cost of cultivation (Rs. ha⁻¹) and Benefit Cost Ratio was calculated using crop yield and cost of cultivation for each treatment. Finally, the collected experimental data were statistically analyzed by ANOVA method suggested by Gomez and Gomez [5] with R software packages (R v.4.2.1).

Comment [B13]: assess

Comment [B14]: remove

Comment [B15]: remove

Comment [B16]: remove

3. RESULTS AND DISCUSSION

3.1. Plant height (cm)

Among various treatments, maximum plant height of 65.38 cm was significantly observed by fertigation of ZnSO₄@ 25 kg ha⁻¹ and FeSO₄ @10 kg ha⁻¹ at 15, 30 and 45 DAS (T₈) as compared to control (T₁) plot. (Table1.). This might be due to split application of Zn and Fe throughout the cropping period through fertigation. Greater availability of Zn and Fe promotes the meristematic growth of plants by stimulation of auxin production. These results were in accordance with the findings of Kumar *et al.* [11]; Vinoth *et al.* [21] and Rathor *et al.* [15].

3.2. Leaf Area Index (LAI)

LAI was significantly influenced, when application of Zn and fertigation of ZnSO₄ (25 kg ha⁻¹) and FeSO₄ (10 kg ha⁻¹) at 15, 30 and 45 DAS (T₈) showed higher LAI of 5.55 than the control (T₁) (3.37) as indicated in Table 1. The reason might be due to continues supply of water and Zn and Fe produces quality of leaves with optimum size, more number of leaves and greater photosynthetic activity. The findings were confirmed with the result of Anwar *et al.* [2] and Gupta *et al.* [6] who reported greater photosynthetic activity and chlorophyll synthesis due to micronutrient fertilization resulting into higher leaf area index.

Comment [B17]: Capital letter

3.3. Yield and yield attributes

The maximum number of 64 pods per plant and seed yield of 1267 kg ha⁻¹ were recorded in fertigation of ZnSO₄ (25 kg ha⁻¹) and FeSO₄ (10 kg ha⁻¹) at 15, 30 and 45 DAS (Table1& Fig.1) which was at par with (T₇). The minimum number of pods per plant (36) and seed yield (910 kg ha⁻¹) was recorded (fig 1). Increased in zinc and iron led to improved capabilities for the effective translocation of photosynthates from source to sink and enhance the enzymatic activity which helps in catalyzing reaction for growth, finally leading to development of more yield attributing characters like number of pods per plant, pod weight and seed yield. Similar findings were related to yield confirmed by Singh *et al.* [18]; Kakade *et al.* [10] and Saakshi *et al.* [16].

3.4. Zn and Fe content in blackgram grains

Among various treatments as indicated in Table 1, drip fertigation of ZnSO₄ (25 kg ha⁻¹) and FeSO₄ (10 kg ha⁻¹) at 15, 30 and 45 DAS (T₈) released significantly increased Zn content of 76 kg ha⁻¹ and Fe content of 184 kg ha⁻¹ in seeds of blackgram compared to control (T₁). This could be attributed to the balanced nutrient absorption achieved through fertigation, which

Comment [B18]: Capital letter

has an impact on pollen production, fertility, and seed quality production. The finding was confirmed with the result of Barla *et al.* [3]; Singh *et al.* [17] and Hanumanthappa *et al.* [7].

Table: 1. Influence of zinc sulphate and ferrous sulphate on growth, yield and seed quality of blackgram under drip irrigation.

Treatments	Plant height (cm)	LAI	No. of pods per plant	Seed yield (kg ha ⁻¹)	Zn content in seed (kg ha ⁻¹)	Fe content in seed (kg ha ⁻¹)	WUE (kg ha ⁻¹ mm ⁻¹)
T ₁ -Control (No Zn and Fe application)	45.14	3.37	36	910	54.50	125.03	1.83
T ₂ - Soil application of RDF of ZnSO ₄ @ 25kg ha ⁻¹ + FeSO ₄ @ 10 kg ha ⁻¹ at basal	52.64	4.39	47	1040	63.13	140.28	2.09
T ₃ - Seed treatment with ZnSO ₄ @ 5 g ha ⁻¹ + FeSO ₄ @ 5 g ha ⁻¹	48.87	3.86	41	975	59.51	132.13	1.96
T ₄ - Foliar spray of ZnSO ₄ @ 0.5% + FeSO ₄ @ 0.5% at 30 & 45DAS	56.34	4.76	52	1107	66.01	148.83	2.23
T ₅ -Seed treatment with ZnSO ₄ @ 5g ha ⁻¹ + FeSO ₄ @ 5g ha ⁻¹ + foliar spray of ZnSO ₄ @ 0.5% + FeSO ₄ @ 0.5 % at 30 & 45DAS	57.02	4.81	54	1113	66.77	152.49	2.24
T ₆ - - Fertigation of ZnSO ₄ @ 12.5 kg ha ⁻¹ + FeSO ₄ @ 5 kg ha ⁻¹ at 15 ,30 & 45 DAS	60.82	5.12	58	1178	70.13	167.48	2.37
T ₇ - Fertigation of ZnSO ₄ @ 18.75kg ha ⁻¹ + Fe SO ₄ @ 7.5 kg ha ⁻¹ at 15, 30 & 45 DAS	64.52	5.48	62	1253	75.15	182.91	2.52
T ₈ - Fertigation of ZnSO ₄ @ 25 kg ha ⁻¹ + FeSO ₄ @ 10 kg ha ⁻¹ at 15, 30 & 45 DAS	65.38	5.55	64	1267	76.73	184.15	2.55
SEd	1.20	0.14	1.26	30.16	1.20	3.16	1.83
CD (P=0.05)	2.57	0.31	2.69	64.68	2.56	6.78	2.09

Comment [B19]: SEd (±)

(RDF- recommended dose of fertilizer, LAI – leaf Area Index, WUE- Water Use Efficiency, Zn – Zinc, Fe – Iron)

3.5. Water Use Efficiency

The data shown in Table 1, indicated that maximum water use efficiency of $2.55 \text{ kg ha}^{-1} \text{ mm}^{-1}$ was noticed in drip fertigation of ZnSO_4 (25 kg ha^{-1}) and FeSO_4 (10 kg ha^{-1}) at 15, 30 and 45 DAS (T_8) compared to control (T_1). This might be due to application of Zn and Fe through drip fertigation, provides essential nutrients directly to the root zone, limiting nutrient loss and leading to higher uptake and improved productivity. These findings were in accordance with Vanishree *et al.* [19] and Fanish and Selvam, [1].

Comment [B20]: Capital letter

Comment [B21]: space

Table: 2. Influence of zinc sulphate and ferrous sulphate on economics of blackgram under drip irrigation.

Treatment	COC (Rs. ha ⁻¹)	GMR (Rs. ha ⁻¹)	NMR (Rs. ha ⁻¹)	BCR
T ₁ – control	41913	68001	26088	1.62
T ₂ - Soil application of micronutrients	43802	77652	33850	1.77
T ₃ - Seed treated with micronutrients	41963	72825	30862	1.74
T ₄ - Foliar application of micronutrients	42104	82641	40537	1.96
T ₅ - Combination of seed and foliar treatment with micronutrients	42153	83184	41031	1.97
T ₆ - Fertigation at 50 % RDF of micronutrients	42833	88005	45172	2.05
T ₇ - Fertigation at 75 % RDF of micronutrients	43273	93486	50213	2.16
T ₈ - Fertigation at 100 % RDF of micronutrients	43943	94551	50608	2.15

(COC – Cost of Cultivation, GMR – Gross Monetary Return, NMR – Net Monetary Return, BCR – Benefit cost ratio)

3.6. Economics

Application of ZnSO_4 (25 kg ha^{-1}) and FeSO_4 (10 kg ha^{-1}) by drip fertigation method at 15, 30 and 45 DAS (T_8) registered higher gross return of Rs.94551 ha⁻¹, net return of Rs.50608 ha⁻¹ and benefit cost ratio of 2.15 over control treatment (T_1) (Table 2). Drip fertigation of micronutrients enhanced the blackgram productivity led to increase in economics of blackgram cultivation. Similar results were supported by Jha *et al.* [9] and Muniyappa *et al.* [13].




Fig 1. Effect of zinc sulphate and ferrous sulphate on seed yield (kg ha^{-1}) of blackgram under drip irrigation.

4. Conclusion

It is concluded that application of ZnSO_4 @ 25 kg ha^{-1} and FeSO_4 @ 10 kg ha^{-1} at 15, 30 and 45 DAS through drip fertigation increased 28.17 percent of blackgram grain yield and enhanced Zn and Fe content in grain of blackgram was 17.72 and 23.82 per cent respectively. Hence, the combined application of Zn and Fe had positive interaction under drip fertigation and the quality of Zn and Fe content was improved through agronomic approaches.

Comment [B22]: a

REFERENCES

1. Anitta Fanish S. and S. Paneer Selvam. Enhancing Productivity of Blackgram through drip Fertigation. Biological Forum – An International Journal. 2023;15(1): 196-200.
2. Anwar Z, Basharat Z, Bilal Hafeez M, Zahra N, Rafique Z, Maqsood M. Biofortification of maize with zinc and iron not only enhances crop growth but also improves grain quality. Asian Journal of Agriculture and Biology. 2021.
3. Barla S, Sahoo HK, Patra BP, Biswasi S, Phonglosa A. Foliar Nutrition of Zinc and Iron on Quality and Productivity of Summer Mungbean (*Vigna radiata* L.).
4. Cakmak I. Enrichment of fertilizers with zinc: An excellent investment for humanity and crop production in India. Journal of trace elements in medicine and biology. 2009 Nov 1; 23(4):281-9.
5. Gomez KA, Gomez AA. Statistical procedures for agricultural research. John Wiley & sons; 1984 Feb 17.
6. Gupta N, Ram H, Kumar B. Mechanism of Zinc absorption in plants: uptake, transport, translocation and accumulation. Reviews in Environmental Science and Bio/Technology. 2016 Mar; 15:89-109.
7. Hanumanthappa D, Vasudevan SN, Shakuntala JB, Muniswamy NM, Macha SI. Enrichment of iron and zinc content in pigeonpea genotypes through agronomic biofortification to mitigate malnutrition. International Journal of Current Microbiology and Applied Sciences. 2018; 7: 4334-42.
8. Indiatat. (2023). India stat Data Base. <https://www.indiastat.com/>
9. Jha DP, Sharma SK, Amarawat T. Effect of organic and inorganic sources of nutrients on yield and economics of blackgram (*Vigna mungo* L.) grown during kharif. Agricultural Science Digest-A Research Journal. 2015; 35(3):224-8.

Comment [B23]: Anitta FS and Selvam SP.

Comment [B24]: Please check

10. Kakade SU, Deshmukh JP, Parlawar ND, Goud VV. Response of split application of nutrients through fertigation in pigeonpea. *Legume Research-An International Journal*. 2023; 46(6):713-20.
11. Kumar R, Baba AY, Kumar M, Bhusan A, Singh K. Growth, nodulation and yield of black gram (*Vigna radiata* L.) as influenced by sulphur and iron under sandy loam soil. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(3):614-6.
12. Kumar S, Palve A, Joshi C, Srivastava RK. Crop biofortification for iron (Fe), zinc (Zn) and vitamin A with transgenic approaches. *Heliyon*. 2019 Jun 1; 5(6).
13. Muniyappa M, Ramachandrappa BK, Nagaraju SA. Studies on the influence of depth and interval of drip irrigation on yield, water use efficiency and economics of chickpea (*Cicer arietinum* L.). *Int J Pure App Biosci*. 2017; 5(1):771-6.
14. Puttaswamy S, Timmagowda S, Krishnamurthy K. Determination of leaf area in pulses. *Curr. Res*. 1976; 5(3):47.
15. Rathor KM, Sharma MK, Meena H, Yadav VK, Yadav RK, Ghasil BP, Meena RK, Yadav SL. Conjunctive Use of Fe, Zn and Bio-Fertilizers on Growth, Nodulation and Chlorophyll Content of Urdbean (*Vigna mungo* L.). *International Journal of Environment and Climate Change*. 2023 Apr 5; 13(5):392-9.
16. Saakshi AR, Rathod PS, Rachappa V, Dodamani BM, Ananda N. Growth, yield and economics of pigeonpea as influenced by biofortification of zinc and iron. *International Journal of Current Microbiology and Applied Sciences*. 2020;9: 3088-97.
17. Singh D, Prasanna R. Potential of microbes in the biofortification of Zn and Fe in dietary food grains. A review. *Agronomy for Sustainable Development*. 2020 Apr; 40:1-21.
18. Singh SP. Effect of micronutrients on nodulation, growth, yield and nutrient uptake in black gram (*Vigna mungo* L). *Annals of Plant and Soil Research*. 2017; 19(1):66-70.
19. Vanishree H, Anand N, Chittapur BM, Umesh MR, Bhat SN. Standardization of fertigation technology for transplanted pigeonpea [*Cajanus cajan* (L.) Millsp.]. *Legume Research-An International Journal*. 2019;42(2):243-6.
20. Viets FG. Fertilizers and the efficient use of water. *Advances in agronomy*. 1962 Jan 1; 14: 223-64.
21. Vinoth D, Vaiyapuri K, Thavaprakash N, Selvakumar S. Influence of different irrigation methods on growth and yield of summer greengram (*Vigna radiata* L.). 2022.