

# **Novel records of testate amoebae (Amoebozoa and Cercozoa) from ShenthurineyWLS, Kerala**

## **ABSTRACT**

This report summarizes the results hitherto achieved in the study of moss inhabitant testate amoebae in Shenthuriney WLS, part of Western Ghats, Kerala. The study reveals the novel records of 28 species of testate amoebae span over 6 families and 9 genera from Shenthuriney WLS, Kerala.

Key words :Shenthuriney WLS, Kerala, Moss, Protozoa, Testate amoebae

## **1. INTRODUCTION**

Shendurney Wildlife Sanctuary, part of Western Ghats of Kerala owes its name to a tree locally called Chenkurinji, an endemic tree which is confined to this tract. Shendurney was declared as a Wildlife sanctuary in 1984 with a total extent of 172 sq. kms. The sanctuary lies on the western part of Agastyamala –Ashambu hill range with an altitudinal span of 100 to 1550m above sea level. This tract is drained by a good number of rivers namely Shendurney, Kazhuthuruthy, Kulathupuzha and their tributaries such as Parappan, Uruli, Pasmakandamthodu, Aruvi and Umayar. The sanctuary falls between 8°44' & 9°14' N latitude and 76°59' & 77°16' E longitude and is located in the Pathanapuram taluk of Kollam district of Kerala state. The reserve forests of Thenmala, Thiruvananthapuram, Punalur forest divisions and Thirunelveli forest division of Tamil Nadu surrounds the sanctuary from all four sides. The terrain is undulating with elevation ranging from 100-1550m ASL. The sanctuary is a continuous stretch of vast and valuable forest area bounded on the east by the Sahyadri hills which act as a great barrier separating the two states, Kerala and Tamil Nadu. The sanctuary lies on the either side of the Shendurney river and is located on the north of Kulathupuzha valley separated by Churuttumala ridge. The whole area is hilly in character with a gentle slope towards the west. The upper slopes are rugged, steep and inaccessible in many places. The highest peak is Alwarkurichi peak. Even though this sanctuary is amidst in Western Ghats with rich biodiversity no attempts were so far made to explore the diversity of testate amoebae. Testate amoebae (TA) are a diverse and abundant group of protists found in

a wide range of habitats around the world and are particularly abundant in wetlands [1]. Testate amoebae are routinely used as indicators of past changes in peatland hydrology [2,1]. These single-celled organisms respond quickly to environmental change, produce decay-resistant and taxonomically distinctive shells, and are generally well preserved and abundant in Holocene peat deposits and present in a variety of habitats like terrestrial, freshwater, estuarine and marine from the tropics to polar areas [3,4]. Testate amoebae form a very sensitive group of organisms [5]. Their short generation times make them useful indicators of environmental changes [6,7,8]. Their well-defined ecological preferences in relation to important ecological variables in different type of ecosystems have made them useful in biomonitoring [9,10].

Testate amoebae research has increased significantly over the past two decades due to their increasing use in different applied aspects as bioindicators for palaeoecological studies, in environmental monitoring, studies on their role in the cycling of elements in the terrestrial ecosystems and biogeographical and evolutionary studies [11]. It is very important to understand the diversity of free-living protists because it plays a very significant role in the ecological health and make up a large part of earth's biodiversity [11,12]. Even though its wide range of applications, not much serious studies have been done and herewith reporting 29 species as new records to this sanctuary.

## 2. MATERIALS AND METHODS

Moss samples (100-200grams) were collected by quadrant sampling (1m<sup>2</sup>) by scrapping from rock and tree bark from the study area during the faunistic survey to Western Himalaya in October 2019. The samples were processed with non-flooded petri dish method as described by Foissner [13] and from each sample permanent mounts were prepared and studied under Nikon 50 i compound microscope for species level identification.

## 3. RESULTS AND DISCUSSION

The present study results the reports of 28 species of testate amoebae from Shenthuriney WLS belong to 9 genera under 6 families (Images 1-29 as Annexure-1). Of these *Cyclopyxisaplanata* Deflandre, 1929, *Cyclopyxisleidyi* (Coûteaux et Chardez, 1981), *Cyclopyxisputeus* Thomas, 1960 and *Cyclopyxistronconica* Godeanu, 1972 were earlier reported only from Corbett NP [14] in India. Even though Western Ghats are rich in biodiversity only 26 species have been reported from Peppara WLS, part of Western Ghats in Kerala [15]. Family hyalosphenidae represented the maximum number of

species(7) followed by the families Netzeiliidae and Centropyxidae (6 each). This is a part of study conducted by Zoological survey of India and this is only a baseline information from the rich biodiversity area and intensive studies should be done to reveal the actual testate fauna from this biodiversity hotspot.

### **Systematic list of testate amoebae recorded as per the classification of Adlet *et al.*, 2019**

Phylum Tubulinea Smirnov *et al.*, 2005

Class Elardia Kang *et al.*, 2017

Order Arcellinida Kent, 1880

#### **Family Arcellidae Ehrenberg, 1843**

1. *Arcella conica* (Playfair, 1918)
2. *Galeripora arenaria*(Greeff, 1866) González-Miguéns *et al.*, 2021
3. *Galeriporadiscoides*(Ehrenberg, 1871) González-Miguéns *et al.*, 2021

#### **Family Netzeiliidae Kosakyan *et al.*, 2016**

4. *Cyclopyxis aplanata* Deflandre, 1929
5. *Cyclopyxis eurystoma* Deflandre, 1929
6. *Cyclopyxis kahli* Deflandre, 1929
7. *Cyclopyxis leidyi* (Coûteaux et Chardez, 1981)
8. *Cyclopyxis puteus* Thomas, 1960
9. *Cyclopyxis tronconica* Godeanu, 1972

#### **Family Diffugiidae Wallich, 1864**

10. *Diffugia oblonga* Ehrenberg, 1838
11. *Diffugialithophila* Penard, 1902
12. *Diffugioglobulosa* Penard, 1902
13. *Diffugialebes* Penard, 1899
14. *Diffugia lucida* Penard, 1890

#### **Family Centropyxidae Jung, 1942**

15. *Centropyxis aculeata* Ehrenberg, 1838
16. *Centropyxis constricta* Penard, 1902
17. *Centropyxis elongata*(Penard, 1890) Thomas, 1959

18. *Centropyxis oblonga* Deflandre, 1929
19. *Centropyxis plagiostoma* Bonnet and Thomas, 1955
20. *Centropyxis platystoma* Deflandre, 1929

**Family Plagiopyxidae** Bonnet and Thomas, 1960

21. *Plagiopyxis declivis* Bonnet, 1955

**Family Hyalospheniidae** Schultze, 1977, emend. Kosakyan and Lara, 2012

22. *Nebelabohemica* Taranek, 1882
23. *Padaungiella lageniformis* (Penard, 1890) Lara and Todorov, 2012
24. *Padaungiella tubulata* (Brown, 1910) Lara and Todorov, 2012
25. *Padaungiella wailesi* (Deflandre, 1936) Lara and Todorov, 2012
26. *Quadrullella variabilis* Kosakyan et al., 2016
27. *Quadrullella symmetrica* (Wallich, 1864) Schulze, 1875
28. *Quadrullella tropica* Wailes, 1912

4. ACKNOWLEDGEMENTS

The author is grateful to the Director, Zoological Survey of India for approving the project and for providing the facilities to do the work. Thanks are also due to the survey team for collecting the samples for the study.

5. COMPETING INTERESTS

Author has declared that no competing interest exists.



6. REFERENCES

1. Mitchell, E.A.D., Charman, D.J. & Warner, B.G. Testate amoebae analysis in ecological and paleoecological studies of wetlands: past, present and future. *Biodiversity and Conservation*, 2008;17, 2115–2137.
2. Charman, D.J. Biostratigraphic and palaeoenvironmental applications of testate amoebae. 2001; *Quaternary Science Reviews*, 20, 1753–1764.
3. Foissner, W. Soil protozoa: fundamental problems, ecological significance, adaptations in ciliates and testaceans, bioindicators, and guide to the literature. *Progressive Protistology*, 1987; 2: 69-212

4. Foissner, W. Description of two new, mycophagous soil ciliates (Ciliophora, Colpodea): *Fungiphryastrobli* n. g., n. sp. and *Grossglockneria ovata* n. sp. *Journal of Eukaryotic Microbiology*, 1999; 46: 34-42.
5. Qin, Y., Mitchell, E.A.D., Lamentowicz, M., Payne, R.J., Lara, E., Gu, Y., Huang, X. and Wang, H. Ecology of testate amoebae in peatlands of central China and development of a transfer function for palaeohydrological reconstruction. *Journal of Paleolimnology*, 2013;50(3): 319-330.
6. Vincke S, Gremmen N, Beyens L, van de Vijver B. The moss dwelling testacean fauna of Île de Possession. *Polar Biol.* 2004a; 27:753–766. doi:10.1007/s00300-004-0655-8
7. Vincke S, Ledeganck P, Beyens L, van de Vijver B. Soil testate amoebae from sub-Antarctic Îles Crozet. *Antarct Sci.* 2004b; 16:165–174.
8. Mattheeussen R, Ledeganck P, Vincke S, van de Vijver B, Nijs I, Beyens L. Habitat selection of aquatic testate amoebae communities on Qeqertarsuq (Disko Island), west Greenland. *Acta Protozool.* 2005; 44:253–263
9. Charman DJ. Modelling hydrological relationships of testate amoebae (Protozoa: Rhizopoda) on New Zealand peatlands. *J R Soc.* 1997; N Z 27:465–483. doi:10.1080/03014223.1997/9517549
10. Mitchell, E.A. D., Buttler, A., Grosvernier., Hydin, H., Albinsson, C., Greenup, A. L., Heijmans, M. P. D., Hoosbeek, M. R., Saarinen, T. Relationships among testate amoebae (Protozoa), vegetation and water chemistry in five sphagnum-dominated peatlands in Europe. *Res New Phytol.*, 2000; 145: 95- 106.
11. Nguyen, H., Gilbert, D., Bernard, N., Mitchell, E.A.D. and Badot, P.M. Relationship between atmospheric pollution characterized by NO<sub>2</sub> concentrations and testate amoebae and diversity. *Acta Protozoologica*, 2004; 43: 233-329.
12. Payne, R.J. Seven reasons why protists make useful bioindicators. *Acta Protozoologica*, 2013; 52: 95-105.
13. Foissner, W. Estimating the species richness of soil protozoa using the non-flooded petridish method. In *Protocols in Protozoology*. 1992; Lee, J.J and Soldo, A.T (eds.), Allen Press, Lawrence, Kansas B-10.1-B10.2.
14. Bindu. L and Sathish Kumar V.M. Novel reports of moss inhabitant testate amoebae (Protozoa: Amoebozoa) from Jim Corbett National park, Uttarakhand, *Int. J. Glob. Sci. Res.*, 2022; 9 (2): 1884-1891.



15. Bindu L. Novel records of testate amoebae (Protozoa : Tubulinea and Cercozoa) from Peppara Wildlife Sanctuary, Agastyamala Biosphere Reserve, Kerala. *Biological Forum-An International Journal*, 2022;14(1) : 1284-1286.

**Annexure-1**




*Arcella conica* (Playfair, 1918) *Galeripora arenaria* (Greeff, 1866)

González-Miguéns *et al.*, 2021




*Galeriporadiscoides* (Ehrenberg, 1871) *Cyclopyxis aplanata* Deflandre, 1929


González-Miguéns *et al.*, 2021





*Cyclopyxis eurystoma* Deflandre, 1929 *Cyclopyxis kahli* Deflandre, 1929



*Cyclopyxis leidyi* (Coûteaux et Chardez, 1981) *Cyclopyxis puteus* Thomas, 1960



*Cyclopyxis tronconica* Godeanu, 1972 *Diffugia oblonga* Ehrenberg, 1838




*Difflugialithophila* Penard, 1902 *Difflugiaglobulosa* Penard, 1902



*Diffflugialebes* Penard, 1899 *Diffflugia lucida* Penard, 1890



*Centropyxis aculeata* Ehrenberg, 1838 *Centropyxis constricta* Penard, 1902




*Centropyxis elongata* (Penard, 1890) Thomas, 1959 *Centropyxis oblonga* Deflandre, 1929




*Centropyxis plagiostoma* *Centropyxis platystoma* Deflandre, 1929

Bonnet and Thomas, 1955




*Plagiopyxisdeclivis* Bonnet, 1955    *Nebelabohemica* Taranek, 1882




*Padaungiellalageniformis* (Penard, 1890)    *Padaungiellatubulata* (Brown, 1910)

Lara and Todorov, 2012    Lara and Todorov, 2012



*Padaungiellawailesi* (Deflandre, 1936)    *Quadrullella variabilis* Kosakyan et al., 2016

Lara and Todorov, 2012



*Quadrullella tropica* Wailes, 1912    *Quadrullella asymmetrica*

(Wallich, 1864)Schulze, 1875

UNDER PEER REVIEW

UNDER PEER REVIEW