

# Review Article

# Applications of Protoplast Fusion in Plant Biotechnology

## ABSTRACT

Plant biotechnology is the deliberate application of biotechnology tools such as protoplast fusion, DNA extraction, plant bioinformatics, PCR and cloning in creating plants with new, improved and desired traits for human benefit. Protoplasts are often referred to as plant cells which the cell wall has been removed and it has many applications in plant biotechnology. The applications of protoplast fusion ranges from the following in plant biotechnology production of useful metabolites, gene editing (crispr mutagenesis), improvement of food nutrition content, creation nitrogen-fixing symbioses, introduction and establishment of disease resistance, production of herbicide resistant plants, e.t.c. The knowledge of protoplast fusion can be use by plant biotechnologist to improve plants trait for human benefits. The application of plant biotechnology is also very important to any nation that will achieve food security.

*Keywords: [Disease Resistance, Food Security, Herbicide Resistant, Nitrogen-Fixing Symbioses, Gene Editing and Nutrition Content]*

## 1. INTRODUCTION

Protoplasts are often referred to as plant cells which the cell wall has been mechanically or enzymatically removed. In theory it is assumed that protoplasts are totipotent, which means they have the ability to dedifferentiate and regenerate into various organs [1] Protoplast fusion has become a very important technique in producing crops with useful agronomic traits and which can also be sold on commercial scale. In the past few decades there have an increased in research using protoplast fusion due to public antagonism with genetically modified crops [2,3]. Protoplast technology is promising technique that can be exploited by breeders to increase germplasm accessibility and can also bring about improvement in different crop varieties [4].

Plant protoplast can be used as versatile cell-based experimental system which can analyze gene expression during a transient time period [5,6,7] and macromolecules like proteins, RNA and DNA can be delivered into protoplasts using various techniques such as microinjection and PEG-calcium fusion electroporation [8,9].

Biotechnology techniques which use DNA analysis have shown significant advancement for the past year and it has been used continuously in characterizing somatic hybrids. SSR has been used extensively for somatic hybrids characterization analysis as a biotechnology technique [10,11,]. [12] Studied the use of proteomics as better toll in regulation and inheritance rules in somatic hybridization. [7] Suggest the use of next generation sequencing which is cheaper and faster as a tool in somatic hybrid genome screening and stability studies for future scientist. High resolution melting analysis which is also another tool that can be use for screening technique based upon insertions, SNPs induce alteration or deletion of double stranded DNA dissociation behavior [13,14]. PCR-RFLP and CAPS analysis has also been found as efficient and reliable tools in cytoplasmic genome characterization [7].

This review, aim at reviewing the biotechnology application of protoplast fusion such as production of useful metabolites, gene editing (crispr mutagenesis), improvement of nutrition

production, introduction and establishment of disease resistance plants, creating nitrogen-fixing symbioses and production of herbicide resistant plants.

## **2.0 PLANT BIOTECHNOLOGY APPLICATIONS**

### **2.1 Production of Useful Metabolites**

Useful metabolite such as anticancer agents, functional proteins, enzymes and antiviral proteins are found in the cell walls of plants, between cell membrane and the cell walls [15,16,17,18,19].

The major challenge is that the accumulations of these metabolites are usually very low. The use of protoplast fusion allows large amount of the metabolite to be released into the culture [20]. To avoid the regeneration of the cell wall, immobilization matrix together with an inhibitor makes the production of the metabolite to be more efficient [20,21,22].

*Catharanthus roseus* protoplasts isolated from callus culture was entrapped in alginate gel to study the extracellular production of enzymes (peroxidase and alpha-galactosidase). However, free protoplasts extracellular production of these enzymes was higher than the protoplasts immobilized in 0.7–2.5% alginate gel beads. Callus culture protoplasts of *C. roseus* immobilized in alginate gels rich in guluronic acid to study the production of secondary metabolite of indole alkaloids (ajmalicine, catharanthine, and tryptamine), which are synthesized through many enzyme reaction steps. Protoplasts immobilized in alginate produced extracellular ajmalicine much higher than protoplasts immobilized in agarose. Addition of 30 mM CaCl<sub>2</sub> to the broth, maintained the active protoplast for 15 days with neither cell wall regeneration nor inhibition of indole alkaloid production [23].

### **2.2 Gene Editing (Crispr Mutagenesis)**

The direct alteration of specific DNA sequence is a vital element of genome editing which is called gene editing. Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) which is associated with Protein 9 (CAS9) method is suitable genome editing technique which requires just two reagent; single guide RNA (sgRNA) and Cas9 protein [24,25,26,27,28,] Other endonucleases like Cpf1 can cause mutations apart from cas9 [29,30,31,32,]. The genome editing reagents can be synthesized and assemble in vitro which can form active ribonucleoprotein (RNP) complexes where the complexes can be delivered into protoplast which can mutagenize the target gene [29,33,34].

In biotechnology, protoplast is usually used to determine target site mutagenesis efficiency and which can also be generated into plants [33]. The regeneration of protoplast is a major challenge when using CRISPR mutagenesis most especially in monocot plants therefore there will be need for high regeneration efficiency protocols which are unavailable [34,35]. CRISPER/Cas9 technology is a very useful tool in research and plant breeding which still a new technology at the infant stage [35]. Using the CRISPER-mediated mutagenesis it is possible to remove the integrated transgenes encoding gene editing reagents from the genome through genetic segregation thereby reducing the fear of the public about genetically modified foods [36].

### **2.3 Improvement of Food Nutrition Content**

The aquatic food chain of fauna aquaculture, microalgae denotes the major natural nutrition. *Chlorella vulgaris* is one example of the sources of aquatic nutrition [37]. The *C. vulgaris* species have been used extensively as nutritional supplements or aquaculture feeds [38]. Microalgae are regarded as one of the organism producing distinct range of bioactive chemical compounds, primarily vitamins, pigments, proteins, minerals, lipids and polysaccharides. The main reasons for their consideration as important source of nutrition for diverse purposes are the high nutritional content of the microalgal species [39,40,41]. [43] Reported the documented the production of genetically improved strain of algae by somatic fusion and hybridization.

Algae-algae protoplast fusion has been reported as a valuable technique to improve their nutrition. Protoplast fusion technique is an in vitro genetic manipulation process which are considered more effectual compared to the conventional techniques used for strain improvement such as selection and mutation [44,45]. Somatic hybridization put up by this procedure has demonstrated strong efficacy in increasing nutrition and valuable metabolites production [46].

According to [47], the use of protoplast fusion technique for microalgae *Chlorella* had been carried out to improve their carotenoid for animal aquatic supplement. During the application of protoplast fusion on interspecific microalgae of *C. vulgaris*, the nutrition content of fusant was subjected for analysis by GCMS methods on *C. vulgaris* powder from 100 L liquid cultivation of the fusant. The study resulted in fusant with high mass production level. 17 amino acid with high concentration of firstly, glutamic acid (14495.52 ppm) secondly, leucine (10856.97 ppm) and thirdly, Aspartic acid (10378 ppm) was showed on the nutrition analysis of fusant. Palmitic acid (1.59%) was showed highest concentration in its lipid acid profile [46].

#### **2.4 Creating Nitrogen-Fixing Symbioses**

Nitrogen which is a compound of many bio-molecules is very important for growth and development of plants. Most nitrogen exist in the atmosphere and the ability fix atmospheric nitrogen through the nitrogenase enzyme complex is restricted to some bacteria and where the bacterian abd plant live in symbiotic relationship provide the reiches sources of nitrogen to plants [48,49]. For the last few decades there have been scientific research on bacteria having nitrogen-fixing symbioses with legumes, which are mostly made of the following genera *Sinorhizobium*, *Mesorhizobium*, *Bradyrhizobium*, and *Rhizobium*, *Azorhizobium* [50,51]. Leguminous plants characterization for nitrogen fixation is mainly dependent on their ability in developing nitrogen-fixing nodules through interaction with the symbiotic bacteria [51]. Actinomycete *Frankia* forms nitrogen-fixing root nodules on non legumes plants thua the genetic basis of the symbiotic interactions that occurs between *Frankia* strains and host plants is poorly understood [53].

Early works on the genetics of nitrogen fixation was studied on free-living nitrogen-fixing bacteria *Klebsiella pneumoniae*. The presence of 17 nil (nitrogen fixation) genes that encode nitrogen fixation in *K. pneumoniae* is responsible for nitrogen fixation [54]. Intergeneric transfer of *K. pneumoniae* nil clusters to other non-nitrogen-fixing bacteria and yeast though nitrogen fixation has only been observed in closely related species [55,56,57].

[58] reported the successful use of Protoplast fusion to create novel actinomycete capable of fixing atmospheric nitrogen. Protoplasts of *Streptomyces griseofuscus*, which is a fast-growing actinomycete and *Frankia* which is a a slow-growing actinomycete were both allowed to fuse and regenerate on media which does not have supply of nitrogen. The regenerated colonies were able to acquire the fast growing characteristics of the streptomycetes and tha ability on growon a media lacking nitrogen from *Frankia*.

[59] Louis and Ensign (1987) also reported the use of four *Frankia* strains ACN1, EAN1pec, Cpl1 and Eullc for the formation and regeneration of protoplasts of the actinorhizal nitrogen-fixing actinomycete *Frankia* where the protoplast was sandwiched between a layer of a nutritionally rich osmotically stabilizing medium and a layer of low-melting-point agarose. It was observed that the regeneration of the four strains varied widely and the maximum regeneration efficiency was only observed on two strains.

[60] Also reported an effective role by protoplast fusion in enhancing nodulation of Rhizobial species. The bacteria abilities to produce nodulation were observed on two weak strains (Rtl1 and Rtl2) and one efficient strain (RtA1) which were selected for protoplast fusion and the numbers of nodules formed by the intra-specific protoplast fusion strains were observed. The Protoplast fusion of the indigenous *R. leguminosarum* biovar *trifolii* strains resulted in nodulation increases by 1.93- to 5.67-fold when compared to their parent strains. This is an

excellent result for agricultural practices for the formation of nitrogen-fixing root nodules on leguminous crops.

The prospect of Green Nitrogen Revolution will be a great achievement in producing stable food crops that has substitute for mineral nitrogen fertilizer which can be achieved by using nitrogen-fixing fertilizers [61].

## **2.5 Production of Herbicide Resistant Plants**

The production of herbicide resistant plant can be achieved through protoplast fusion. Attempts have been made to make plants tolerance herbicides such as bromoxynil, atrazine, sulphonylureas, glyphosate and phosphinothricin [62]. Many herbicide operate by inactivating some plant enzymes (target proteins) which are very important for functions such as the photosynthetic or other biosynthetic pathways which are unique to plants [63].

[64] Reported the use of bar gene in combination with the herbicide Basta to select transformed rice (*Oryza sativa* L. cv. Radon) protoplasts for the production of herbicide-resistant rice plants. The presence of Phosphinothricin acetyltransferase (PAT) assays was used to confirm the expression of the bar gene in plants obtained from phosphinothricin resistant calli. Both the bar and gusA genes were transmitted to progeny as confirmed by Southern analysis, where the progeny having the bar gene was found to be resistant to Basta. Thus Herbicide or Basta can be used as a post-emergence on rice plants transformed with the bar gene.

[65] also reported the use of Terbutryn-resistant plastids of the *Nicotiana plumbaginifolia* TBR2 mutant which was introduced into *N. tabacum* plants by protoplast fusion following X-irradiation of TBR2 protoplasts where the Cybrid plants were founded resistant to high levels of atrazine (10 kg/ha). The level of atrazine resistance (to 10 kg/ha) is likely to be sufficiently enough to protect the crop under field conditions because atrazine is mostly applied at a rate of 2-4.5 kg/ha.

[66] Confirmed that an important Indica rice cultivar *Oryza sativa* cv. IR72 was transformed with the application of direct gene transfer to protoplasts. The transformed rice showed resistance to high dosage level of phosphinothricin.

## **2.6 Introduction and Establishment of Disease Resistance**

[67] Reported the use of protoplast fusion to overcome sexual incompatibility between cultivated potatoes and diploid *Solanum*. They developed a systematic protocol for the isolation of huge number of high quality protoplasts from variety of Mexican wild species that has high levels insect [Colorado potato beetle (CPB)] resistance and disease (late blight). Using the protocol, new somatic hybrids of one Argentina wild species, two Mexican and cultivated potato clones and the successful somatic hybrids were from the cell fusion of *S. tuberosum* and the diploid species *S. pinnatisectum*, *S. cardiophyllum*, and *S. chacoense* which shows higher level of resistance to both late blight and CPB than was found in *S. tuberosum*. [68] reported *Camelina sativa* and *Brassica napus* hybrids which has increased linolenic acid content when compared to the *B. napus* partner.

Cybridizations and Somatic hybridizations in citrus produced rootstocks that is resistant to abiotic and biotech constraints which increased the yield and quality of the fruit [69] also in brown spot resistant scions [70]. [71] also reported the production of the first resistant raphano-brassica asymmetric and symmetric hybrids. This new development showed new resistance types along with multiple resistances which include turnip mosaic virus.

## **3. DISCUSSION**

Protoplast fusion is a major breeding tool that is used to produce new genetic combination which is different from other scientific tools and it also transfers mono and polygenic traits [72,73].

Genomic variation is of important interest in most plants for yield and quality improvement. Quality improvement, disease resistance, cytoplasmic male sterility (CMS) transfer, Salt

tolerance, rootstock improvement and seedless triploids are the most frequent goal of protoplast fusion [74].

Other researchers have also report important use of protoplast fusion in plant biotechnology. [75] reported the protoplast fusion for intergeneric hybrid cells. [76] Studied the microbead encapsulation of living plant protoplasts which is seen as a new tool for the handling of single plant cells. [77] reported the interspecific T-DNA transfer through plant protoplast fusion. [78] Reported the Protoplast fusion for production of triploids and tetraploids which is used for rootstock and scion breeding in citrus. [79] Describe plant adenine base editor based on an evolved tRNA adenosine deaminase fused to the nickase CRISPR/Cas9 which enable A•T to G•C conversion in protoplasts and regeneration in rice and wheat plants.

Many countries especially some of the developing countries have many fears about the application of plant biotechnology but have forgotten also about the two major challenges of plant biotechnology which are the continuous increase in population at geometric rate and the current climatic changes which are posing a very serious threat to the human population and the growth of our plants (crops). If the challenges of the plant biotechnology will be solved then the application of plant biotechnology has become very necessary. Since protoplast fusion allows the introduction of new genes into plants without genetically modified plants, which is the fear of common then protoplast fusion, offers an option.

Plant biotechnology tool such protoplast fusion has therefore become essential for plant (crops) improvement for the future. The use of protoplast fusion will go a long way to remove the fear of genetically modified food in the mind of the common man.

#### **4. CONCLUSION**

The fear of increase in population and climatic change has left us with no other option in feeding ourselves than the use of plant biotechnology for improved food production to meet our ever increasing population. Plant biotechnology tool such protoplast fusion has therefore become essential for plant improvement for the future. The use of protoplast fusion will go a long way to remove the fear of genetically modified food in the mind of the common man

#### **CONSENT**

Not applicable

#### **ETHICAL APPROVAL**

Not applicable

#### **REFERENCES**

- [1] Johnson A, Veilleux R. Somatic hybridization and application in plant breeding. In: Janick J (ed) Plant Breeding Rev 20. John Wiley & Sons, New York, (2001) 167–225
- [2] Davey M, Anthony P, Patel D, Power J. Plant Protoplasts: Isolation, Culture and Plant Regeneration. In: Davey M, Anthony P (eds) Plant cell culture: essential methods. John Wiley & Sons, New York, 2010;153–173
- [3] Grosser J, Calovic M, Louzada E. Protoplast fusion technology— somatic hybridization and cybridization. In: Davey M, Anthony P (eds) Plant cell culture: essential methods. John Wiley & Sons, New York, 2010;175–198
- [4] Hawkes JG. The potato: Evolution, biodiversity and genetic resources. Am. J. Pot Res. 1990;67, 733–735

- [5] Wu FH, Shen SC, Lee, LY, Lee SH, Chan MT, Lin CS; Tape-Arabidopsis Sandwich - a simpler Arabidopsis protoplast isolation method. *Plant Methods*. 2009;5:16.
- [6] Lung SC, Yanagisawa M, Chuong SDX. Protoplast isolation and transient gene expression in the single-cell C4 species, *Bienertia sinuspersici*. *Plant Cell Rep*, 2011;30(4):473–484
- [7] Eeckhaut T, Lakshmanan SP, Deryckere D, Bockstaele VE, and Huylenbroeck VJ. Progress in plant protoplast research. *Planta*. 2013;238:991–1003.
- [8] Hassanein A, Hamama L, Loridon K, Dorion N. Direct gene transfer study and transgenic plant regeneration after electroporation into mesophyll protoplasts of *Pelargonium x hortorum*, 'Panache Sud'. *Plant Cell Rep*. (2009);28:1521–1530
- [9] Masani MYA, Noll GA, Parveez GKA, Sambanthamurthi R, Prüfer D. Efficient transformation of oil palm protoplasts by PEG-mediated transfection and DNA microinjection. *PLoS One*. 2014;9(5)
- [10] Toppino L, Menella G, Rizza F, D'Alessandro A, Sihachakr D, Rotino GL. ISSR and isozyme characterization of androgenetic dihaploids reveals tetrasomic inheritance in tetraploid somatic hybrids between *Solanum melongena* and *Solanum aethiopicum*. Group Gilo. *J Hered*. 2008;99:304–315
- [11] Sarkar D, Tiwari J, Sharma S, Sharma PS, Gopal J, Singh B, Luthra S, Pandey S, Pattanayak D. Production and characterization of somatic hybrids between *Solanum tuberosum* L. and *S. pinnatisectum* Dun. *Plant Cell Tiss Org* . 2011;107:427–440
- [12] Gancle A, Grimplet J, Sauvage F, Ollitrault P, Brillouet J. Predominant expression of diploid mandarin leaf proteome in two citrus mandarin-derived somatic allotetraploid hybrids. *J Agr. Food Chem*;2006;54:6212–6218
- [13] Wu S, Wirthensohn M, Hunt P, Gibson J, Sedgley M. High resolution melting analysis of almond SNPs derived from ESTs. *Theor Appl Genet* 118:1–14
- [14] Deryckere D, De Keyser E, Eeckhaut T, Van Huylenbroeck J, Van Bockstaele E. High-resolution melting analysis as a rapid and highly sensitive method for *Cichorium* plasmotype characterization. *Plant Mol Biol Rep*. 2013;31:731–740
- [15] Millar, D.J., Whitelegge, J.P., Bindschedler, L.V., Rayon, C., Boudet, A.M., Rossignol, M., Borderies, G. and Bolwell, G.P.,. The cell wall and secretory proteome of a tobacco cell line synthesising secondary wall. *Proteomics*, (2009)9(9), 2355-2372.
- [16] Cassab GI, Varner JE. Cell wall proteins. *Annual Review of Plant Physiology and Plant Molecular Biology*. , 1988;39(1):321-353.
- [17] Bonness MS, Ready MP, Irvin JD, Mabry TJ, Pokeweed antiviral protein inactivates pokeweed ribosomes; implications for the antiviral mechanism. *The Plant Journal*. 1994; 5(2):173-183.
- [18] Fett-Neto AG, Pennington JJ, Di Cosmo F. Effect of white light on taxol and baccatin III accumulation in cell cultures of *Taxus cuspidata* and *Zucc*. *J Plant Physiol*. 1995;146:584–90

- [19] Emmanuel DB, Gali AI, Peingurta AF, Afolabin SA. Callus Culture for the Production of Therapeutic Compounds. *American Journal of Plant Biology* 2019;4(4):76-84
- [20] Aoyagi H. Application of plant protoplasts for the production of useful metabolites. *Biochemical Engineering Journal*. 2011.56(1-2):1-8.
- [21] Mera N., Aoyagi H, DiCosmo F, Tanaka H. Production of cell wall accumulative enzymes using immobilized protoplasts of *Catharanthus roseus* in agarose gel. *Biotechnology letters*. 2003; 25(20);1687-1693.
- [22] Bodeutsch T, James E, Lee J. The effect of immobilization on recombinant protein production in plant cell culture. *Plant Cell Rep*. 2001;20;562–566
- [23] Aoyagi H, Sakamoto Y, Asada M, Tanaka H. Indole alkaloids production by *Catharanthus roseus* protoplasts with artificial cell walls containing of guluronic acid rich alginate gel. *Journal of fermentation and bioengineering*. 1998;85(3);306-311
- [24] Nekrasov V, Staskawicz B, Weigel D, Jones JDG, Kamoun S. Targeted mutagenesis in the model plant *Nicotiana benthamiana* using Cas9 RNA-guided endonuclease. *Nat. Biotechnol*. 2013;31:691-693.
- [25] Li JF, Norville JE, Aach J, McCormack M, Zhang DD, Bush J, *et al*. Multiplex and homologous recombination-mediated genome editing in *Arabidopsis* and *Nicotiana benthamiana* using guide RNA and Cas9. *Nat. Biotechnol*. 2013;31:688-691.
- [26] Gaj T, Gersbach CA, Barbas CF. ZFN, TALEN, and CRISPR/Cas-based methods for genome engineering. *Trends Biotechnol*. 2013;31:397-405.
- [27] Feng ZY, Zhang BT, Ding WN, Liu XD, Yang DL, Wei PL, *et. a*. Efficient genome editing in plants using a CRISPR/Cas system. *Cell Res*. 2013;23:1229-1232
- [28] Shan QW, Wang YP, Li J, Zhang Y, Chen KL, Liang Z, Zhang K, *et al*. Targeted genome modification of crop plants using a CRISPR-Cas system. *Nat. Biotechnol*. 2013;31: 686-688
- [29] Kim H, Kim ST, Ryu J, Kang BC, Kim JS, Kim SG. CRISPR/Cpf1-mediated DNA-free plant genome editing. *Nat. Commun*. 2017;8:14406
- [30] Endo A, Masafumi M, Kaya H, Toki S. Efficient targeted mutagenesis of rice and tobacco genomes using Cpf1 from *Francisella novicida*. *Sci. Rep*. 2016;6:38169
- [31] Xu R, Qin R, Li H, Li D, Li L, Wei P. Generation of targeted mutant rice using a CRISPR-Cpf1 system. *Plant Biotechnol J*. 2017;15:713-717
- [32] Mahfouz MM. Genome editing: The efficient tool CRISPR-Cpf1. *Nat. Plants*. 2017;3:17028.
- [33] Woo JW, Kim J, Kwon SI, Corvalan C, Cho SW, Kim H, *et al*. DNA-free genome editing in plants with preassembled CRISPR-Cas9 ribonucleoproteins. *Nat. Biotechnol*. . 2015;33:1162-1164.
- [34] Liang Z, Chen K, Li T, Zhang Y, Wang Y, Zhao Q, Liu J. Efficient DNA-free genome editing of bread wheat using CRISPR/Cas9 ribonucleoprotein complexes. *Nat. Commun*. 2017;8:14261.

- [35] Zhang Y, Liang Z, Zong Y, Wang Y, Liu J, Chen K. Efficient and transgene-free genome editing in wheat through transient expression of CRISPR/Cas9 DNA or RNA. *Nat. Commun.* 2016;7:12617.
- [36] Huang S, Weigel D, Beachy RN, Li J. A proposed regulatory framework for genome-edited crops. *Nat. Genet.* 2016;48,109-11.
- [37] Kusumaningrum HP, Zainuri M. Detection of bacteria and fungi associated with *Penaeus monodon* postlarvae mortality. *Procedia Environmental Sciences.* 2015;23:329-337.
- [38] Pulz O, and Gross W. Valuable products from biotechnology of microalgae. *Applied microbiology and biotechnology*, 2004;65(6):635-648.
- [39] Mathias D, Hammantola S, Ishaku G. Isolation and Characterization of Biofloculant-Producing Bacteria from Wastewater at Jimeta, Adamawa State, *J. Adv. Biol. Biotechnol.*, 2017;15( 1):1–7.
- [40] Adebisi A, Basseyy EE, Ayo R, Bello I, Habila J, Ishaku GA. Anti-mycobacterial, Antimicrobial and phytochemical evaluation of *Pulicaria crispa* and *Scoparia dulcis* plant extracts. *Journal of Advances in Medical and Pharmaceutical Sciences.* 2016;7(4):1-11.
- [41] Njobdi S, Gambo M, Gali AI. Antibacterial Activity of *Zingiber officinale* on *Escherichia coli* and *Staphylococcus aureus*. *Journal of Advances in Biology and Biotechnology.* 2018;19(1):1-8.
- [42] Sukorno IF, Islam S, Kabir LA, de la Cruz VC, Zaman S, Gali AI. Phytochemicals are natural resources of food supplement for happier people. *Horticult Int J.* 2019;3(6):300–305
- [43] Lee YK, Tan HM. Interphylum protoplast fusion and genetic recombination of the algae *Porphyridium cruentum* and *Dunaliella* spp. *Microbiology.* 1988;134(3):635-641.
- [44] Wang L, Li Y, Chen P, Min M, Chen Y, Zhu J. Anaerobic digested dairy manure as a nutrient supplement for cultivation of oil-rich green microalgae *Chlorella* sp. *Bioresource technology*2010;101(8):2623-2628.
- [45] Lu Y, Kong R, Hu L. Preparation of protoplasts from *Chlorella protothecoides*. *World Journal of Microbiology and Biotechnology.* 2012;28(4):1827-1830.
- [46] Kusumaningrum HP, Zainuri M. Aplikasi Pakan Alami Kaya Karotenoid untuk Post Larvae *Penaeus monodon* Fab.(Applications Rich Natural Carotenoids Feed for Post Larvae of *Penaeus monodon* Fab.). *ILMU KELAUTAN: Indonesian Journal of Marine Sciences.* 2013;18(3):143-149.
- [47] Kusumaningrum HP, Zainuri M, Improvement of Nutrition Production by Protoplast Fusion Techniques in *Chlorella vulgaris*. *J Food Process Technol.* 2018;9: 711
- [48] Kneip C, Lockhart P, VoB C and Maier U..Nitrogen fixation in eukaryotes – New models for symbiosis *BMC Evolutionary Biology.* 2007;7:55
- [49] Johansson C. Bergman B. Early events during the establishment of the *Gunnera/Nostoc* symbiosis. *Planta.* 1992;188:403-413

- [50] MacLean MA, Finan MT, Sadowsky JM. Genomes of the Symbiotic Nitrogen-Fixing Bacteria of Legumes. *Plant Physiology*. 2007;144:615–622
- [51] Hakoyama T, Niimi K, Yamamoto T, Isobe S, Sato S, Nakamura Y. *et al.* The Integral Membrane Protein SEN1 is Required for Symbiotic Nitrogen Fixation in *Lotus japonicus* Nodules. *Plant Cell Physiol*. 2012;53(1): 225–236.
- [52] Laporte P, Satiat-Jeunemaître B, Velasco I, Csorba T, Van de Velde W, Campalans A *et al.* A novel RNA-binding peptide regulates the establishment of the *Medicago truncatula*–*Sinorhizobium meliloti* nitrogen-fixing symbiosis. *The Plant Journal*. 2010;62:24–38
- [53] Normand P, Lapierre P, Tisa LS, Gogarten PJ, Alloisio N, Bagnarol E. *et al.*, Genome characteristics of facultatively symbiotic *Frankia* sp. strains reflect host range and host plant biogeography. *Genome Research*. 2007;17:7–15
- [54] MacNeil T, MacNeil D, Roberts GP, Supaino MA, Brill N J: Fine-structure mapping and complementation analysis of *nif* (nitrogen fixation) genes in *Klebsiella pneumoniae*. *J Bacteriol*. 1978;136:253-266
- [55] Dixon RA, Postgate JR: Genetic transfer of nitrogen fixation from *Klebsiella pneumoniae* to *Escherichia coli*. *Nature*. 1972; 237:102-103.
- [56] Krishnapillai V, Postgate JR. Expression of *Klebsiella* *his* and *nif* genes in *Serratia marcescens*, *Erwinia herbicola* and *Proteus mirabilis*. *Arch Microbiol*. 1980;127: 115-118.
- [57] Zamir A, Mainn CV, Fink GR, Szalay AA: Stable chromosomal integration of the entire nitrogen fixation gene cluster from *Klebsiella pneumoniae* in yeast. *Proc Natl Acad Sci USA* 1981;78:3496-3500.
- [58] Prakash RK, Cummings B, 1988. Creation of novel nitrogen-fixing actinomycetes by protoplast fusion of *Frankia* with streptomycetes. *Plant molecular biology*. 10(3):281-289
- [59] Louis ST, Ensign JC, 1987. Formation and Regeneration of Protoplasts of the Actinorhizal Nitrogen-Fixing Actinomycete *Frankia*. *Applied And Environmental Microbiology*. 53(1)53-56
- [60] Sabir MS J, El-Bestawy E. Enhancement of nodulation by some arid climate strains of *Rhizobium leguminosarum* biovar *trifolii* using protoplast fusion. *World J Microbiol Biotechnol*. 2009;25:545–552
- [61] Dent D, Cocking E. Establishing symbiotic nitrogen fixation in cereals and other non legume crops: The Greener Nitrogen Revolution. *Agric & Food Secur*. 2017;6:7
- [62] Oxtoby E, Monica A. Hughes Breeding for herbicide resistance using molecular and cellular techniques. *Euphytica*. 1989;40:173-180
- [63] Botterman J, Leemans J. Engineering of Herbicide Resistance in Plants, *Biotechnology and Genetic Engineering Reviews*. 1988;6(1): 321-340
- [64] Rathore SK, Chowdhury KV, Hodges KT. Use of *bar* as a selectable marker gene and for the production of herbicide-resistant rice plants from protoplasts. *Plant Molecular Biology*. 1993;21: 871-884.

- [65] Menczel L, Polsby SL, Steinbaek EK, Maliga P. Fusion-mediated transfer of triazine-resistant chloroplasts: Characterization of *Nicotiana tabacum* cybrid plants. *Mol Gen Genet*. 1986;205:201-205
- [66] Datta K.S. Karabi Datta K., Soltanifar N., Donn G. and Potrykus I. Herbicide-resistant *Indica* rice plants from IRR1 breeding line IR72 after PEG-mediated transformation of protoplasts. *Plant Molecular Biology*. 1992;20: 619-629.
- [67] Chen Q, Li H Y, Shi Y Z, Beasley D, Bizimungu B, Goettel MS. Development of an effective protoplast fusion system for production of new potatoes with disease and insect resistance using Mexican wild potato species as gene pools. *Can. J. Plant Sci*. 2008;88: 611-619
- [68] Jiang J, Zhao X, Tian W, Li T, Wang Y. Intertribal somatic hybrids between *Brassica napus* and *Camelina sativa* with high linolenic acid content. *Plant Cell Tiss Org*. 2009 99:91–95
- [69] Dambier D, Benyahia H, Pensabene-Bellavia G, Kacar Y, Froelicher Y, Belfalah Z. Somatic hybridization for Citrus rootstock breeding: an effective tool to solve some important issues of the Mediterranean Citrus industry. *Plant Cell Rep*. 2011;30:883–900
- [70] Soriano L, Assis F, Camargo L, Cristofani-Yaly M, Rocha R, Andrade C. Regeneration and characterization of somatic hybrids combining sweet orange and mandarin/mandarin hybrid cultivars for citrus scion improvement. *Plant Cell Tiss Org*. 2012;111:385–392
- [71] Xiao Z, Wan L, Han B. An interspecific somatic hybrid between *Actinidia chinensis* and *Actinidia kolomikta* and its chilling tolerance. *Plant Cell Tiss Org*. 2004;79:299–306
- [72] Grosser J, Gmitter F. Applications of somatic hybridization and cybridization in crop improvement, with Citrus as a model. *In Vitro Cell Dev Plant*. 2005;41:220–225
- [73] Thieme R, Darsow U, Rakosy-Tican L, Kang Z, Gavrilenko T, Antonova O, Heimbach U, Thieme T. Use of somatic hybridization to transfer resistance to late blight and potato virus Y (PVY) into cultivated potato. *Plant Breed Seed Sci*. 2004;50:113–118
- [74] Wang J, Jiang J, Wang Y. Protoplast fusion for crop improvement and breeding in China. *Plant Cell Tiss Org*. 2013;112:131–142
- [75] Kao KN, Constabel F, Michayluk MR, Gamburg OL. *Plant Protoplast Fusion and Growth of Intergeneric Hybrid Cells*. *Planta (Berl.)*. 1974;120:215--227
- [76] Matthew SG, Philip M. Lintilhac Microbead encapsulation of living plant protoplasts: A new tool for the handling of single plant cells. *Applications in Plant Sciences* 2016;4(5): 1500140
- [77] Miiller-Gensert E, Schieder O. Interspecific T-DNA transfer through plant protoplast fusion. *Mol Gen Genet* 1987;208:235-241.
- [78] Grosser JW, Gmitter GF, Jr Protoplast fusion for production of tetraploids and triploids: applications for scion and rootstock breeding in citrus. *Plant Cell Tiss Organ Cult*. 2011;104:343–357

[79] Li C, Zong Y, Wang Y, Jin S, Zhang D, Song Q. Expanded base editing in rice and wheat using a Cas9-adenosine deaminase fusion. *Genome Biology*. 2018;19:59

UNDER PEER REVIEW