

Original Research Article

Allelopathic Effects of Selected Weed Species on Seed Germination and Seedling Growth of Mungbean (*Vigna radiata* L.)

ABSTRACT

Four weed species viz., Durba (*Cynodondactylon*), White cock's comb (*Celociaargentia*), Shyama (*Echinochloa crus-galli*), Shak notae (*Amaranthus viridis*) were used on Mungbean (BARI Mug 6). Boiled and unboiled weed extracts reduced the germination and primary growth of mungbean. The ranking of weed species in respect of inhibitory effect on seed germination of mungbean was *Celociaargentia* > *Cynodondactylon* > *Amaranthus viridis* > *Echinochloa crus-galli*. Durba (*Cynodondactylon*) and White cock's comb (*Celociaargentia*) had strong detrimental effect on primary growth of mungbean. The extracts of all the parts of weeds inhibited germination and primary growth of this crop. The relative efficiency of inhibition of germination and primary growth by different weed extracts was in the order of stem > whole plant > leaf > root. The reduction of germination and primary growth increased with increasing extract concentrations. So, it could be suggested that *Cynodondactylon* and *Celociaargentia* had strong detrimental effect on mungbean. Therefore, the cited weeds must be taken into well care and it should be avoided in seed bed for growing mungbean seedlings.

Keywords: Allelopathic effect, weed species, mungbean, seed germination, seedling growth

INTRODUCTION

Allelopathy is considered as a defense mechanism available to plants, which release chemicals and influence the growth and development of neighboring plants in both natural and agricultural ecosystems. This process happens through root exudation, leaching, volatilization, residue decomposition and other means by creating direct and indirect effects on the adjacent microenvironment (Khanh *et al.*, 2007) due to chemical substances released by plants (Einhelling, 2008).

Allelochemicals results in detrimental effects and is categorized into two which are true allelopathy and functional allelopathy (Duke, 2015). The true allelopathy is the release of substances which are toxic in nature from its origin as they are produced in plants (Duke, 2015). Functional allelopathy is the release of toxic things that are resultants of transformation by microorganism (Inderjit *et al.*, 2002; Jabran and Farooq, 2013). These chemicals accrue and

persevere for a substantial time in the plant, result significant interference on growth and development of neighboring plants (Einhelling, 2008) that can be either a crop or a weed. With considerable work conducted in past decades, the presence of restrictive compounds in a wide array of plant types and its parts at varying concentrations were investigated and with remarkable attention being paid to allelopathy (Inderjit *et al.*, 2002) and its related effects.

As the understanding of the mechanisms of these allelochemicals in plants grew, a great significant attention was given to the potential use of allelopathy as a weed management technique in achieving sustainable agriculture particularly in organic farming (Dayan *et al.*, 2009). Integration of allelopathy into natural and agricultural management systems might be helped to reduce the usage of synthetic agrochemicals. Ultimately, it will reduce the negative effects on the environment, especially environment pollution. Exploring crop allelopathy against weeds might be a useful strategy to eliminate problems created by synthetic agro-chemicals (Anjum and Bajawa, 2005). Recently, several crops and weeds showed clear weeds suppressive ability and with strong use in agricultural fields, by exuding allelochemical compounds either from living plants or from decomposing residues (Xuan *et al.*, 2006). Further, this positive effect on weed control will be helpful in introducing potential allelopathic crops into crop rotation programmes, either as a cover or mulch to smother crops, or as a green manure in sustainable agricultural practices. This is a very positive and important trend in modern agriculture, which could apply for the benefit of human beings by reducing excess use of chemical usage. Further, it helps to control severe health problems that could be caused due to unsafe handling of agrochemicals (Belz, 2007). Moreover, the idea of allelopathy can be utilized effectively to produce eco-friendly natural herbicides (Sodaeizadehet *et al.*, 2009).

During past decades, the suppressive ability of allelochemicals in weeds has drawn significant attention. However, the specific weed or plant species and its unique responses were not clearly studied yet. Therefore, the focus of this study was to assessment the allelopathic effect of some naturally grown weed species that commonly occur in agricultural fields. The objective of the study was to observe the allelopathic effect of four selected common broadleaf weed species on mungbean seeds and seedlings and to identify the weed plant parts which have more allelopathic response on crops.

MATERIALS AND METHODS

Plant materials:

Mungbean is almost important pulse crops containing high nutritive value. It not only plays important role in human diet but also in improving the soil fertility by fixing the atmospheric nitrogen. Its seed is more palatable, nutritive, digestible and non-flatulent than the other pulses. BARI Mug 6 was used in this experiment.

Allelopathic plant species based on a literature review performed as active allelopathic plant species will be chosen. Aqueous extracts of Durba (*Cynodon dactylon*), White cock's comb (*Celocia argentea*), Shyama (*Echinochloa crus-galli*), Shak notae (*Amaranthus viridis*) considered as allelopathic materials.

Experimental treatment

The boiled and unboiled selected weed extracts were used. Stem, leaf, root and whole plant extracts of different weed species were considered.

Collection of weeds and preparation of weed extracts

Weed species collected from the research field, Bangladesh Agricultural Research Institute, Gazipur. After collecting they were washed in distilled water for removing dust and soil particles, then chopped and macerated. In a set fresh weed mass (250g) of each weed species was boiled in 1 litre water and kept for 3 days with intermittent stirring. The extracts were filtered through filter paper. The filtrates were used as boiled extracts of weeds. In another set, 250g fresh weed mass were decomposed in 1 litre water for 7 days at normal room temperature ($28 \pm 1^\circ\text{C}$). The extracts were filtered and used as unboiled extract. Thus boiled and unboiled extracts of each weed species were prepared and used in this experiment following the technique as stated by Prasad and Srivastava (1991).

Experimental Procedure

The germination of mungbean seeds was studied by petridish method. There are two sets of trials, each with unboiled and boiled extracts for each crop, were performed. Twenty five seeds of each test crops were placed in each petridish with double layer of filter paper and treated with weed extracts. A control set with distilled water was set simultaneously every time. The filter papers were kept moist with distilled water constantly. The experiment was arranged in a completely randomized design (CRD) with 3 replications at room temperature ($28 \pm 1^\circ\text{C}$).

Parameters

- a) Germination (%)
- b) Days to complete germination
- c) Seedling growth and weight
 - i) Root and shoot lengths
 - ii) Root and shoot weights
- d) Chlorophyll content

In this experiment, 10 plants were selected to each parameter and the average results were calculated from these collected plants. The number of germinated seeds was counted from the beginning of the seed germination up to completion of the seed germination. The root and shoot lengths, fresh & dry weights of respective root and shoot of mungbean seedlings were measured after 7 and 14 days respectively of the germination setting.

Determination of Chlorophylls:

Chlorophyll was estimated by Arnon's method (1949). 0.5gm of sample grinds into 5.0ml of 80% acetone in clean mortar and pestle. It was centrifuged at 5000 rpm for 10 minutes and supernatant will be used for read the absorbance at 645 and 663 nm against the solvent (80% Acetone) blank.

RESULTS & DISCUSSION

Effect of boiled and unboiled weed extracts on mungbean

Effect on seed germination

The experimental data on the effect of boiled and unboiled weed extracts on seed germination have been shown in Table 1. It was found that the seed germination varied significantly due to the effect of boiled and unboiled weed extracts on seed germination of mungbean seed. The allelopathic effect of different species of weeds induced significant reduction in mungbean seed (BARI mug 6) germination. Maximum germination rate (98.80 %) was observed in control treatment while the minimum (33.40%) was obtained in *Celociaargentina* extract treated seeds. The ranking of the weed species in respect of inhibitory effect on seed germination of BARI mug 6 was *Celociaargentina*>*Cynodondactylon*>*Amaranthus viridis*>*Echinochloa crus-galli*. Again the number of days required to complete the germination of seeds was also affected significantly by the weed extracts. The number of days required to complete the germination was recorded minimum (3.0 days) in control treatment whereas it was maximum (5.50 days) due to the effect of *Cynodondactylon* extract which was statistically similar with that of *Celociaargentina*. Similar inhibitory effect of weed species on seed germination was noted by Rahman *et al.*, (1996). Binita *et al.*, (2001) found the inhibitory effect on germination and seedling growth of vegetables by different weed species.

On the other hand, the effect of boiled and unboiled extracts of different weed species induced significant variation in germination rate and number of days required to complete the germination of mungbean seeds which have been shown in Table 2. Highest germination rate

(99.20%) was recorded in control treatment (boiled water) which was statistically similar with that of control (unboiled water). On the other hand, the lowest germination rate was found (30.0%) due to the effect of unboiled extract of *Celociaargentina*. The number of days required to complete germination was also significant in boiled and unboiled extracts of different weed species. Maximum days (5.7 days) required to complete the germination was noted due to the effect of unboiled extract of *Cynodondactylon* which was statistically similar with unboiled extract of *Celociaargentina* and minimum days (3.0 days) in control treatment (both boiled and unboiled water).

Table 1. Effect of extract of different weed species on germination and primary growth of Mungbean

Weed extracts	Germination (%)	Days to complete germination	Root length (cm)	Shoot length (cm)	Fresh wt. of root (mg)	Dry wt. of root (mg)	Fresh wt. of shoot (mg)	Dry wt. of shoot (mg)	Total Chloro phyll content
Control (water)	98.8	3.0	6.87	16.72	44.90	4.35	284.70	31.60	14.63
<i>Cynodondactylon</i>	45.4	5.5	1.28	4.29	16.95	1.53	110.25	15.25	11.57
<i>Celociaargentina</i>	33.4	5.3	0.21	1.15	14.8	1.12	99.3	13.9	9.48
<i>Echinochloacrusgalli</i>	88.3	4.1	3.77	7.91	36.2	3.49	270.5	27.05	13.76
<i>Amaranthus viridis</i>	86.3	4.3	4.34	8.72	39.15	3.86	265.9	26.65	14.50
Level of significance	**	**	**	**	**	**	**	**	**
CV (%)	0.69	3.42	10.30	3.78	0.29	3.26	0.30	2.52	0.62

**Significant at 1% level of probability.

Effect on root and shoot lengths

Root and shoot lengths of Mungbean seedlings reduced significantly due to the allelopathic effect of boiled and unboiled weed extracts as presented in Table 1 & 2. The highest root length (6.87cm) was observed in control treatment while it was lowest (0.21 cm) in the extract of *Celociaargentina*. Again, the highest shoot length (16.72 cm) was found in control and the lowest (1.15 cm) was found in that of *Celociaargentina* (Table 1).

The effect of boiled and unboiled extracts of weeds on BARI mung 6 have been presented in Table 2 indicating significant variation in root and shoot lengths of seedlings. Maximum root length (7.63 cm) was recorded in control while the minimum (0.16 cm) was in boiled extract of *Celociaargentina*. Minimum root length was statistically similar with those of unboiled and boiled extract of *Celociaargentina*. Maximum shoot length (18.12 cm) was also observed in control and the minimum (1.09 cm) was due to the effect of boiled extract of *Celociaargentina*. Minimum shoot length was statistically similar with that of unboiled extract of *Celociaargentina*.

Table 2: Effect of boiled and unboiled weed extracts on germination and primary growth of Mungbean

Weed extracts	Germination (%)	Days to complete germination	Root length (cm)	Shoot length (cm)	Fresh wt. of root (mg)	Dry wt. of root (mg)	Fresh wt. of shoot (mg)	Dry wt. of shoot (mg)
Control (water)								
BW	98.5	3.0	7.63	18.12	44.2	4.1	284.3	31.0
UBW	99.2	3.0	6.1	15.32	45.6	4.6	285.1	32.2
<i>Cynodondactylon</i>								
BE	46.3	5.3	1.17	4.37	16.3	1.44	109	14.2
UBE	44.5	5.7	1.38	4.21	17.6	1.62	111.5	16.3
<i>Celociaargentia</i>								
BE	36.8	5.0	0.16	1.09	14.40	1.06	98.4	13.8
UBE	30.0	5.6	0.25	1.21	15.20	1.18	100.2	14.0
<i>Echinochloacrusgalli</i>								
BE	84.6	4.0	2.65	6.68	35.80	3.43	266.6	26.7
UBE	88.0	4.2	4.89	9.14	36.60	3.56	274.4	27.4
<i>Amaranthus viridis</i>								
BE	90.3	4.0	3.05	7.55	38.00	3.78	256.8	25.4
UBE	82.3	4.6	5.62	9.88	40.30	3.94	275.0	27.9
Level of significance	**	**	**	**	**	**	**	**
CV (%)	0.69	3.42	10.30	3.78	0.29	3.26	0.30	2.52

Note: BE = Boiled extract; UBE= Unboiled extract; BW= Boiled water; UBW= unboiled water;

**Significant at 1% level of probability.

Effect on fresh and dry weights of root

Fresh and dry weights of roots of mungbean reduced significantly due to the allelopathic effect of boiled and unboiled weed extracts as have been presented in table 1 & 2. Maximum fresh (44.90 mg) and dry (4.35 mg) weights were observed in control while the minimum fresh (14.80 mg) and dry (1.12 mg) weights of root were found in aqueous extract of *Celociaargentia* (table 1). The effect of boiled and unboiled extracts of weeds on Mungbean (table 2) reflecting significant reduction in fresh and dry weights of roots of seedlings. Maximum fresh (45.60 mg) and dry (4.60 mg) weights of root were recorded in boiled water control. On the other hand, minimum fresh (14.40 mg) and dry (1.06 mg) weights were due to the effect of boiled extract of *Celociaargentia*.

Effect on fresh and dry weights of shoot

Fresh and dry weights of shoot of mungbean seedling were reduced significantly due to the allelopathic effect of different weed species. Maximum fresh (284.70 mg) and dry (31.60 mg)

weights were observed in control. Again, the minimum fresh (99.30 mg) and dry (13.90 mg) weights of shoot was found in extract of *Celociaargentina* (Table 1). The effects of boiled and unboiled extracts of weeds on mungbean have been presented in table 2 indicating significant inhibition on the fresh and dry weights of shoot of seedlings. Maximum fresh (285.10 mg) and dry (32.20 mg) weights of shoot were recorded in boiled water control treatment. On the other hand, the minimum fresh (98.4 mg) and dry (13.80 mg) weights were recorded due to the effect of boiled extract of *Celociaargentina*. Minimum dry weight of shoot was statistically similar with those of boiled and unboiled extract of *Cynodondactylon* and unboiled extract of *Celociaargentina*.

Chlorophyll content:

Total Chlorophyll content of wheat seedlings was reduced significantly by the allelopathic effect of different weed species. Maximum Chlorophyll content (14.63) was observed in control and minimum rate (9.48) was found in *Celociaargentina* extract treated seeds (table 1). It was observed that the chlorophyll content was comparatively higher due to less allelopathic effect of *Echinochloacrusgalli* (13.76), *Amaranthus viridis* (14.50) extract treated seeds. The ranking of weed species in respect of inhibitory effect on seed germination of wheat was *Cynodondactylon*>*Celociaargentina*>*Amaranthus viridis*>*Echinochloacrusgalli*.

Allelopathic effect of different weed plants parts on mungbean

Celociaargentina and *Cynodondactylon* showed more inhibitory effect on BARI mug 6 compared to other weed species, these two weed species were selected further study.

Effect on germination

The results revealed that the germination of mungbean was suppressed significantly due to the extract of different weed plant parts as presented in Table 3 & 4. Highest germination rate (98.85%) was observed in control and the lowest (30.05%) was in stem extract of different weeds (Table 3). The variation in germination of mungbean seed (BARI mug 6) was significant at 1% due to the interaction between different plant parts of two weeds species. Highest germination rate (98.85%) was found in control while the lowest (24.30%) was in stem extract of *Celociaargentina* (Table 4). Roy (1995) observed that the mass of stem of the weed species was more detrimental than leaf and root. The germination of mungbean seed was delayed by the allelopathic effect of extract of different weed plant parts. Maximum days required (6.1 days) to complete the germination was recorded in aqueous extract of stem and the minimum days (3.0 days) were found in control as presented in Table 3.

Effect on root and shoot lengths

Root and shoot lengths of mungbean seedlings were reduced significantly by the allelopathic effect of different plant parts. Highest root length (6.87 cm) was observed in control and the lowest length (0.84 cm) was found in stem extract of weeds. Lowest length of root was statistically similar with leaf, root and whole plant extracts of weeds. Again the highest shoot length (16.72 cm) was observed in control and the lowest length (2.26 cm) was found in stem extract of weeds. Lowest shoot length was also similar with that of leaf extract of weeds (Table 3). While considering the effect of plant parts of two species of weeds it was observed that the variation in root and shoot lengths of mungbean seedling was significant at 1%. Highest root (6.87cm) and shoot (16.72 cm) lengths were recorded in control and the lowest root (0.55 cm) and shoot (1.07 cm) lengths were found in stem extract of *Celociaargenia*. Minimum length of root and shoot was statistically similar with that of leaf, root and whole plant extracts of *Celociaargenia* (Table 4).

Table 3: Effects of extract of different weed plant parts on germination and primary growth of Mungbean

Weed extracts	Germination (%)	Days to complete germination	Root length (cm)	Shoot length (cm)	Fresh wt. of root (mg)	Dry wt. of root (mg)	Fresh wt. of shoot (mg)	Dry wt. of shoot (mg)
Control (water)	98.85	3.00	6.87	16.72	44.90	4.28	274.10	30.80
Stem Extract	30.05	6.1	0.84	2.26	16.40	1.24	110.60	14.50
Leaf Extract	33.70	5.9	0.95	2.42	17.70	1.45	116.35	19.54
Root Extract	42.80	5.6	1.52	2.61	19.30	1.62	105.24	12.62
Whole weed Extract	32.90	5.4	1.0	2.49	18.44	1.51	115.20	15.90
Level of significance	**	**	**	**	**	**	**	**
CV (%)	0.62	3.42	2.02	1.76	0.28	3.30	0.31	2.58

**Significant at 1% level of probability

Table 4: Interaction effect of weeds species and weed plants parts on germination and primary growth of Mungbean

Weed extracts	Germination (%)	Days to complete germination	Root length (cm)	Shoot length (cm)	Fresh wt. of root (mg)	Dry wt. of root (mg)	Fresh wt. of shoot (mg)	Dry wt. of shoot (mg)
Control (water)	98.85	3.00	6.87	16.72	44.90	4.28	274.10	30.80
Extract of <i>Cynodondactylon</i>								
Stem	40.20	6.0	1.11	3.34	16.70	1.34	112.60	16.50
Leaf	44.10	5.8	1.19	3.47	17.65	1.55	115.35	19.34
Root	48.50	5.3	1.13	4.95	15.30	1.19	103.21	13.60
Whole plant	42.00	5.5	1.21	3.98	17.10	1.48	117.20	21.10
Extract of <i>Celociaargenia</i>								
Stem	24.30	6.1	0.55	1.07	16.50	1.28	116.20	15.32
Leaf	26.70	5.7	0.63	1.12	17.10	1.44	122.85	21.17
Root	36.40	5.6	0.71	1.17	16.80	1.05	116.00	15.32
Whole plant	28.00	5.5	0.69	1.21	17.30	1.33	115.30	14.79
Level of significance	**	**	**	**	**	**	**	**
CV (%)	0.62	3.42	2.02	1.76	0.28	3.30	0.31	2.58

**Significant at 1% level of probability

Effect on fresh and dry weights of root

Fresh and dry weights of root of mungbean seedling were significantly inhibited due to the allelopathic effect of weed plant parts as presented in Table 3. Maximum fresh (44.90 mg) and dry (4.28 mg) weights of root were recorded in control and minimum fresh (16.40 mg) and dry (1.24 mg) weights were recorded due to the root extract of two different weeds. The inhibition of fresh and dry weight of root of BARI mug 6 seedling as reduced by the extract of different plant parts of two weed species have been shown in Table 4. Maximum fresh (44.90 mg) and dry (4.28 mg) weights were found in control. Minimum fresh (15.30 mg) weight found in root extracts of *Cynodondactylon* and dry (1.05 mg) weights was found in root extract of *Celociaargenia*.

Effect on fresh and dry weights of shoot

Fresh and dry weights of shoot of seedling of BARI mug 6 were significantly reduced due to the allelopathic effect of plant parts of two weed species. Maximum fresh (274.10 mg) and dry (30.80 mg) weights of shoot was observed in control while the minimum fresh (105.24 mg) and dry (12.62 mg) weights was recorded due to the root extract of different weeds (Table 3). The variation in fresh and dry weights of shoot of mungbean seedlings was significant at 1% due to

the effect of extract of different plant parts of two weed species. Maximum fresh (274.10 mg) and dry (30.80 mg) weights were found in control treatment. Minimum fresh (103.21 mg) and dry (13.60 mg) weights of root were recorded in root extract of *Cynodondactylon*. Respectively, Minimum dry weight of shoot was statistically similar with that of whole plant extract of *Celociaargentina* (Table 4).

CONCLUSIONS

The allelopathic influences of Durba (*Cynodondactylon*), White cock's comb (*Celociaargentina*), Shyama (*Echinochloacrusgalli*), Shak notae (*Amaranthus viridis*) were tested in bioassays on Mungbean. The aqueous extracts of above weeds significantly suppressed the seed and seedlings of above crops. All the selected weeds had more or less inhibitory effect on seed germination, number of days to complete germination, root and shoot lengths, fresh and dry weights of root and shoot of mungbean. Among the plant parts, stem extract was most inhibitory followed by leaves. From the above findings of the present experiment it could be suggested that *Cynodondactylon* and *Celociaargentina* had strong detrimental effect on mungbean cultivation. Therefore the cited weeds must be taken into well care and it should be avoided in seed bed for growing mungbean seedlings. The results of this experiment proved that it was important to exclude these allelopathic weeds specially *Cynodondactylon* and *Celociaargentina*.

REFERENCES

- Anjum, T., Bajwa, R., (2005). A bioactive annuionone from sunflower leaves. *Phytochemistry* 66(16):1919-1921.
- Arnon, D. I., (1949). Copper enzyme in isolated chloroplast polyphenol oxidase in *Beta vulgaris*, *plant physiol* [give full name of journal](#) 24:1-15
- Belz, R. G., (2007). Allelopathy in crop/weed interactions-an update. *Pest Mmanagement Science* 63(4):308-326.
- Binita, H., Sannigrahi. A. K., Hazarika, B., (2001). Allelopathic research in vegetable production. *Enriron Ecol.* [give full name of journal](#) 19(4): 799-806.
- Dayan, F. E., Charles L., Cantrell Stephen O., Duke., (2009). Natural products in crop protection. *Bioorganic & Medicinal Chemistry*. Volume 17, Issue 12, 15 June 2009, Pages 4022-4034
- Duke, S. O., (2015). Proving allelopathy in crop–weed interactions. *Weed Science* 63(1):121-132.

Formatted: Underline

- Einhelling, F. A., (2008). Mechanism of Action of Allelochemical in Allelopathy. In: Allelopathy Organism Processes and Application. American Chemical Society, Washington, USA pp. 96-116.
- Inderjit, S. J. C., Olofsdotter M., (2002). Joint action of phenolic acid mixtures and its significance in allelopathy research. *Physiologia Plantarum* 114(3):422-428.
- Jabran, K., Farooq M., (2013). Implications of potential allelopathic crops in agricultural systems. In *Allelopathy* pp. 349-385. Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-30595-5_15
- Khanh, T. D., Xuan T. D., Chung I. M., (2007). Rice allelopathy and the possibility for weed management. *Annals of Applied* [??/ 151\(3\):325-339](#).
- Prasad, K., Shrivastava V. C., (1991). Teletoxic effect of some weeds on germination and initial growth of rice (*Oryza saliva* L.). [Indian J. Agric. Sci.give full name of journal](#) 61 (8): 501-502.
- Rahman, M. M., Rahman M. M., Roy B., Ahmed S., (1996). Allelopathic effects of weeds on germination and growth of rice and jute seeds. [Progress. Agric.give full name of journal](#)7(1):69-72.
- Roy, K. C., 1995. Allelopathic effect of five selected weed species on corn (cv. Barnali) seed germination. plant height and biomass weight. M. Sc. (Ag.) [Thesis. Dept.of Agronomy. Bangladesh Agricultural link ersitN Mymensingh.???](#)
- Sodaeizadeh, H., Rafieiolhossaini M., Havlík J., Van Damme P., (2009). Allelopathic activity of different plant parts of *Peganum harmala* L. and identification of their growth inhibitors substances. *Plant Growth Regulation* 59(3):227.
- Xuan, T. D., Elzaawely A. A., Deba F., Fukuta M., Tawata S., (2006). Mimosine in *Leucaena* as a potent bio-herbicide. *Agronomy for Sustainable Development* 26(2):89-97.

Formatted: Underline

Formatted: Underline

Formatted: Underline

Formatted: Underline