

Original Research Article

Chemical composition, Antibacterial and Antioxidant activities of extracts from dry leaves and ash-dry leaves of *Luffa cylindrica* (L.) Roem cultivated in Vietnam

Abstract

Luffa cylindrica (L.) Roem was traditionally used to treat stomachaches, as an antihyperlipidemic and antioxidant, particularly for atherosclerosis therapy, as a suppository to cure constipation spleenopathy, as an anthelmintic, carminative, emmenagogue, galactagogue, and as an antiseptic. Therefore, the aim of this study was to study the chemical composition, antibacterial and antioxidant, of an extract from dry leaves (LuL) compared to ash-dry leaves (LuA) of *Luffa cylindrica* (L.) Roem.

The result showed the physical-chemical effect, phytochemical effect, antioxidant activity, and antibacterial activity, including metal ion content. However, in comparison between the extract from dry leaves (LuL) and ash-dry leaves (LuA) of *Luffa cylindrica* (L.) Roem, there was the most quantitative phytochemical determination, such as the cardiac glycosides, alkaloids, phenolics, flavonoids, and triterpenoids effects presented in the LuL extract because the LuA sample was burned out incompletely into ash. Thus, this study implied that the activity of extracts from dry leaves (LuL) and ash-dry leaves (LuA) contained both bacteriostatic and bactericidal effects, antioxidation, the flavonoids w metal ions which may contribute to the wound healing effect and should be further study.

Keywords: *Luffa cylindrica* (L), inorganic herbal metal ions , antibacterial, antioxidant.

1. Introduction

Luffa cylindrica (L.) Roem belongs to the family *Cucurbitaceae* [1]. The origin of *Luffa cylindrica* is believed to be in South America [2]. *Luffa cylindrica* is commonly grown in Guinea, Ivory Coast, the Philippines, India, and China [3]. The flowers, buds, and young leaves can be used as food [4]. When the fruit is old and dry, cleared of its epidermis and seeds, it gives an excellent sponge called "vegetable sponge," which can be used as a body scrub, pot, or appliance. It is also used as a heavy metal absorber for dehydration [5], [6]. The seed oil is edible. In America, oil is used as an ingredient in soapmaking [7], [8]. The traditional use has been reported in Africa, China, Vietnam, Cambodia, Thailand, Laos, and the Philippines [9]–[11]. The fruit is used as a galactagogue, the roots as a hydragogue and purgative [12], and the root and the whole plant as a suppository to cure constipation [13]. Seed acts as an anthelmintic drug, an inducing vomiting drug, and a laxative [9], [14], [15].

The leaves are prescribed for skin diseases, to treat wounds, to reduce swelling, and to treat stomachaches, antihyperlipidemic and antioxidant particularly for atherosclerosis therapy [16] . Freshly crushed leaves act as emmenagogues, blood detoxifiers, and are used to treat papules and swelling skin [17]. A decoction of leaves is used as a diuretic [18]. Past research has found that leaf extract contains saponin, flavonoids, alkaloids, and cardiac glycosides, and the extract can inhibit *Bacillus subtilis*, *Escherichia coli*, *Staphylococcus aureus*, and *Salmonella typhi* [19]. Aqueous extracts also have an oxytocic activity [13], [20].

The present knowledge of the wound healing process comprises coagulation, inflammation, proliferation, formation and accumulation of fibrous tissues, collagen deposition, epithelialization, contraction of the wound with the formation of granulation tissues, remodeling, and maturation [17], [21].

The constituents of the plant extracts modulate one or more of the above stages. It was the endeavor to identify the active constituents responsible for antimicrobial activity, free radical scavenging properties, stimulators of enhanced collagen production, and/or angiogenesis promoters through the identification of lead scaffold chemical structures [20], [21].

Some studies have shown that *Luffa Cylindrica* is able to affect wound healing, which is a wisdom of folk medicine in many countries [22], but in Vietnam it is used in a different way by using only the leaves [23]. Used to treat wounds to make the wound heal faster. Minerals in organics are known to have an effect on wound healing, such as zinc and chromium shots, which speed wound healing [24]. In addition, diabetic patients are characterized by scarring and chronic wounds, which are rare [22]. Studies on the trend of using *Luffa* leaves for wound healing suggest that this may be a product that helps with diabetes [22], [25].

In our previous research, the trace of traditional use of *Luffa* leaves was done in Hai Duong province, which is located in the center of the Red River Delta with a total area of 1,668.28 km² and a population of more than 1.9 million people. The province has good conditions for agriculture, transportation, and industrial production and plays an important role in the social and economic development of the country. A total of six traditional medicine practitioners were interviewed for this survey. Informed consent was obtained from all, and the survey was explained to them in detail, including the information that the survey results may be published internationally. Findings showed that *luffa* leaf was used long ago by both traditional medicine doctors and the old generation themselves to treat open wounds that were affected long-term by bacteria or fungi. The conservative burned ash from *luffa leaf* was pound into dried powder and then applied to the acne, boils, pressure ulcers, and fungal infection areas. For many cases of pressure ulcers and fungal infections in the area between the toes during flood season, this treatment was very effective. The wounds were quickly healed and recovered. The aim of this study was to study the chemical composition, antibacterial, antioxidant of extract from dry leaves (LuL) compared to ash-dry leaves (LuA) of *Luffa cylindrica* (L.) Roem. as a preliminary study that sample extracts obtained from Vietnam have properties that contribute to the cause of wound healing.

Materials and Methods

Plant material, extraction, and chemicals

Luffa leaves were collected from a Vietnamese farm in Hui Dong Province, Vietnam. Voucher specimen No. 0023302 was identified and kept at the Herbarium of the Faculty of Pharmacy, Chiang Mai University, Thailand. The chemical ingredients and solvent used for extraction of the leaves and ash are pharmaceutical grade and were purchased from Union Sciences Co. Ltd., Thailand. The leaf was dried in a hot air oven at 60 °C and ground to powder (LuL). *Luffa* ash (LuA) was prepared by burning the *Luffa* dried leaves at a normal temperature in open air until the blackish-grey ash was obtained in an uncompleted burning condition. This process was done by the local people.

The Pharmacogenetic Evaluation of the Raw Material of Crude Dried Leaf

A microscopic examination of powdered LuL was studied. The TLC, moisture content, and extractive value were done according to Thai Herbal Pharmacopoeia V.I. in order to prove the scientific database for further uses. Two systems of developing solvents for TLC plates were used: hexane: ethyl acetate (6:4) and dichloromethane: ethyl acetate (9:1). TLC patterns were

determined under UV light at 254 and 366 nm detectors. The plate was sprayed with a freshly prepared anisaldehyde-sulfuric acid reagent (AS).

Sample extraction

Samples of LuL and LuA were ground to 60–80 mesh size with an electric grinder. Each sample was extracted with 95% ethanol in a ratio of 1:10. Sonication was done under an ultrasonic device for 1 hour, separated the clear parts, repeated three times, and then evaporated under pressure.

Quantitative Phytochemical Determination

1) Determination of Total Phenolic Content The extract solution (1 mg/ml in methanol, 1 ml) was mixed with 10% Folin-Ciocalteu reagent (Sigma-Aldrich, Germany, 1 ml), then mixed for 5 minutes, added saturated sodium carbonate (60 g/l, 1 ml), and allowed to stand for 90 minutes in the dark at ambient temperature. The absorbance of the reaction mixture was measured by a UV/Vis spectrophotometer (Shimadzu, Japan) at a wavelength of 725 nm using gallic acid as a standard.

2) Determination of Total Flavonoids Content The extract solution (1 mg/mL in ethanol: water 1:1, 1 ml) was mixed with a 2% AlCl₃ solution (1 ml) and kept in a dark place at ambient temperature for 25 minutes. The absorbance was determined at 415 nm compared with rutin.

3) Determination of Total Alkaloids Content The extract solution (0.1 µg /ml in purified water, 1 ml) was mixed with phosphate buffer solution (pH 4.7, 2 ml). The bromocresol green solution (2 ml) was added to the mixture and then extracted with 1, 2, and 2 ml of chloroform. The absorbance was determined at 415 nm by using berberine chloride as a standard.

4) Determination of Total Triterpenoids Content The extract solution (1 mg/ml in glacial acetic acid, 200 µl) was mixed with a 5% vanillin-acetic acid solution (1 ml) and sulfuric acid (1.8 ml). The sample solutions were allowed to stand at 70°C for 30 minutes and then cooled down to room temperature before adding glacial acetic acid (2 ml). The absorbance of sample solutions was measured at 573 nm by using ursolic acid (Tokyo Chemical, Japan) as a standard.

5) Determination of Total Cardiac Glycoside Content The extract (1 mg/ml in 50% aqueous ethanol, 1 ml) was mixed with 1 ml of freshly prepared Baljet's reagent (95 mL of 1% picric acid and 5 ml of 10% sodium hydroxide solution). The reaction mixture was incubated for 1 h, then diluted with 2 ml of purified water. The absorbance was quantitatively determined at 495 nm by using digoxin as a standard.

Biological activities of LuL and LuA extract

Determination of Total Phenolic Content (TPC)

The total phenolic content of the sample was examined by the Folin-Ciocalteu colorimetric method modification [26]. The sample solutions (1 mL) were mixed with 5 mL of the Folin-Ciocalteu reagent (diluted with distilled water in a ratio of 1:10). After 8 min, a sodium carbonate solution (4 mL, 7.5% w/v) was added and incubated in the dark at room temperature for 2 hrs. Finally, the absorbance of the test samples was measured at 765 nm by a Milton Roy Spectronic 21D spectrophotometer. The gallic acid equivalent values (GAE mg/100g) were calculated and compared with the standard curve of gallic acid. All tests were done in triplicate.

Determination of Antioxidant Activity

The *Diphenylpicryl-hydrazyl* (DPPH) radical scavenging assay was used for determination using the method described by Wu et al., 2005 [27]. The solution of DPPH radicals was prepared in methanol (81.2 mM in methanol). The sample solution (1 mL) was mixed with 5 mL of DPPH solution. The mixtures were vigorously shaken and left for 30 minutes in the dark. The absorbance was measured at 517 nm using methanol as a blank. 5 mL of DPPH solution in 5 mL of methanol was used as a control.

Percent inhibition = $[(A \text{ control} - A \text{ sample}) / A \text{ control}] \times 100$. Where A control is the absorbance of only DPPH radical solution, A sample is the absorbance of a sample mixed with DPPH radical solution. The sample concentration providing 50% inhibition (IC₅₀) was calculated from the graph of inhibition percentage against sample concentration.

Test of the efficiency of extracts to inhibit bacterial growth by agar-well diffusion

Staphylococcus aureus, *Pseudomonas aeruginosa*, and *Escherichia coli* were cultured in Muller-Hinton broth and incubated at 37 °C [28], [29]. Then the bacteria were used to adjust the turbidity of the solution with McFarland No. 0.5 using a sterile cotton swab. Infect food with Muller Hinton agar and use a cork borer to drill holes. And add 100 µl of extract into the test hole. The food plate was incubated at 37 °C for 24 hours. By measuring the clear diameter of the inhibition.

Minimal Inhibitory Concentration, MIC, and Minimum Bactericidal Concentration, MBC

Dilute the two-fold serial dilution with Muller Hinton broth. Food and culture media tested in MHB broth were incubated at 37 °C for 18–24 hours. Then adjust the turbidity of the bacteria. With the McFarland No. 0.5 solution. After that, add the test fuel to the test tube. And was incubated at 37 °C for 18–24 hours. Then, the lowest concentration of the extract was able to inhibit the growth of the test bacteria. By looking at the turbidity of the test in vitro compared to the control tube that did not detect the growth of the infection, And the experiment tube was incubated at 37 °C for 18–24 hours without turbidity or growth of streak plates on MHA agar. 99.99% sterilization test.

Identification of the metal ions in the samples

The LuL and LuA dried samples were sent for checking for metal ions at the Central Laboratories (Thailand) Co., Ltd. By the in-house method based on EPA 3052 and the ICP-OES technique.

3. Result

3.1 Microscopic Identification. The Microscopic characteristic of *Luffa leave* is as **fig.1, fig. 2, fig.3** and **fig.4**

The diagnostic characters are:

1. In surface view, the fragments of the lamina in the upper polygonal epidermis and lower epidermis are wavy in outline. Anomocytic stomata were also present on both surfaces.
2. Palisade mesophyll is usually found in surface view; it is composed of cells with thin walls, circular in outline, containing abundant chloroplasts.
3. The fragments of spongy mesophyll show thin-walled parenchyma containing moderately large chloroplasts with large intercellular spaces and air chambers.

4. The vascular strand is found in various sizes and views, some of which are associated with spongy mesophyll.
5. The fragments of spiral and reticulated vessels in longitudinal view are not very frequent.
6. The occasional fibers could be found in groups or solitary.
7. The occasional glandular trichome appeared as whole trichomes with stalk and head, or fragments of them.
8. The very occasional tracheid fragments in longitudinal view
9. Starch grains are seldom found and accumulate in parenchymatous tissue.

UNDER PEER REVIEW




Fig.1 Powder drug of *Luffa cylindrica* leaf; 1-6 upper epidermis showing stoma, 7-8 lower epidermis showing stoma and wavy epidermis, 10 upper epidermis over vein with stoma, 11-16 upper epidermis with palisade underneath, 17-18 epidermis in sectional view




Fig.2 Powder drug of *Luffa* sp. Leaf (continuee) ; 19-21 palisade in surface view, 22-24 spongy mesophyll, 25-27 polygonal epidermis, 28-29 vascular strand of mesophyll, 30-32 vascular bundle, 33-34 spiral vessel associated with chlorenchyma, 35-36 fragments of vessel




Fig.3 Powder drug of *Luffa cylindrica* Leaf (continuee) ; 37-38 spiral vessel, 39 reticulated vessel, 40 xylem element showing reticulated vessel and xylem fiber, 41-42 tracheid, 43 group of fiber, 44-45 solitary fiber, 46-48 trichome stalk, 49-52 capitate stalked trichome in surface view, 53 capitate stalked trichome head, 54 starch grains

3.2 Physico-chemical Identification

The samples were tested as in the Thai Herbal Pharmacopeia. The physico-chemical examinations were as follows: Loss on drying, total ash, ethanol-soluble extractive value, and chloroform water extractive value were presented in Table 1 along with their mean values. TLC is shown as in Fig. 4, Fig. 5, Table 2, and Table 3. From the result, the optimum system used in quality control of raw materials, LuL, should be: Solvent system: Hexane: ethyl acetate (6:4), with a UV 366 detector.

Table 1 Pharmacogenetic characteristic of LuL

Specification	Content (%)
Loss on drying	13.01±0.20
Total ash	21.9144±0.59
Ethanol-soluble extractive value	11.7900±0.17
Chloroform water extractive value	18.3633±0.22

Thai Herbal Pharmacopoeia 1995 Volume 1 pp.123, 126




Fig.4 Sample before (1) and after (2) detected with UV 254 detector, with UV 366 detector (3) and after spraying with Anisaldehyde – sulfuric acid (AS) (4). Solvent system: Hexane: Ethyl acetate (6:4)




Fig.5 Sample before (1) and after (2) detected with UV 254 detector, with UV 366 detector (3) and after spraying with Anisaldehyde – sulfuric acid (AS) (4). Solvent system: Dichloromethane: Ethyl acetate (9:1)

Table 2 TLC of Luffa leaf extract in Hexane: Ethyl acetate (6:4)

Rf value	Visual inspection	Spot color UV254	Spot color UV366	Spraying with Anisaldehyde – sulfuric acid
0.14	-	-	orange	-
0.26	opaque	opaque	orange	-
0.36	-	-	orange	purple-gray
0.48	opaque	opaque	orange	-
0.50	opaque	opaque	orange	-
0.60	-	-	orange	-

Table 3 TLC of Luffa leaf extract in Dichloromethane: Ethyl acetate (9:1)

Rf value	Visual inspection	Spot color UV254	Spot color UV254	Spraying with Anisaldehyde – sulfuric acid
0.26	-	-	orange	purple-gray
0.30	-	-	orange	-
0.44	opaque	opaque	orange	-
0.84	-	-	orange	-
0.90	-	-	orange	-
0.98	-	-	orange	-

3.3 Quantitative Phytochemical Determination

Phytochemical screening of LuL and LuA demonstrated the presence of cardiac glycosides, alkaloids flavonoids, phenolics and triterpenoids as showed in Table 4.

Table 4 Contents bioactive compounds reference with their standards

Extract No.	Name of the extract	Cardiac glycosides		Alkaloids		Phenolics		Flavonoids		Triterpenoids	
		mg digoxin /g extract	SD	mg berberine /g extract	SD	mg gallic acid /g extract	SD	mg Rutin /g extract	SD	mg Ursolic acid/g extract	SD
1	LuA	22.3	0.17	10.7	0.59	16.7	0.24	20.3	1.02	6.9	0.68
2	LuL	63.7	0.75	20.1	0.95	59.0	0.98	47.8	0.31	46.1	0.34

LuA contained secondary metabolites as follows: cardiac glycosides 22.3±0.17 (mg digoxin/g extract), flavonoids 20.3±1.02 (mg rutin/g extract), phenolics 16.7±0.24 (mg gallic acid/g extract), alkaloids 10.7±0.59 (mg berberine/g extract), terpenoids 6.9±0.68 (mg ursolic acid/g extract). LuL contained secondary metabolites as follows: cardiac glycosides 63.7±0.75 (mg digoxin/g extract), phenolics 59.0±0.98 (mg gallic acid/g extract), flavonoids 47.8±0.31 (mg rutin/g extract), triterpenoids 46.1±0.34 (mg ursolic acid/g extract), alkaloids 20.1±0.95 mg (berberine/g extract). It was shown that LuA still contained all active compounds but in a lower quantity than sample LuL; this appeared because the sample was burned out into ash incompletely.

3.4 Antioxidant activity of LuA and LuL extract

The DPPH radical scavenging activities of LuL extract showed an IC₅₀ value of 87.63 ± 10.00 mg/ml. or LuL 87.63 mg/ml can remove 50 % of free radical, DPPH. Whereas LuA presented % inhibition 3.17±0.69 (mg/mL) or LuA at a concentration of 1 mg/ml reduce can remove 3.7 % of DPPH ($p < 0.05$). This phenomenon showed the importance of phenolic compounds in DPPH's radical scavenging activities.

3.5 Antibacterial Activity of *Luffa* Leaf (LuL) Extract and *Luffa* Ash (LuA) Extract

In this study, the inhibitory activities of leaf extract were investigated for the minimal inhibitory concentration (MIC) and minimal bactericidal concentration (MBC) against pathogenic bacteria, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, and *Escherichia coli*. by broth dilution method. The result demonstrated that LuL extract revealed an inhibitory effect on all tested bacteria. Moreover, LuL extract showed the lowest MIC and MBC values of 125 mg/ml against *Staphylococcus aureus* and *Pseudomonas aeruginosa*, followed by *Escherichia coli* with MIC and MBC values of 250 mg/ml (Table 5). But LuA extract showed the lowest MIC and MBC values of 125 mg/ml against *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Escherichia coli*. This revealed that LuA extract possessed more effective in antibacterial, *E. coli* than LuL.

Table 5 MIC and MBC values of LuL extract and LuA extract against pathogenic bacteria

sample	MIC and MBC (mg/ml)		
	<i>S. aureus</i>	<i>Ps. aeruginosa</i>	<i>E. coli</i>

	MIC	MBC	MIC	MBC	MIC	MBC
LuL extract	125± 0.0	125± 0.0	125± 0.0	125± 0.0	250± 0.0	250± 0.0
LuA	125± 0.0	125± 0.0	125± 0.0	125± 0.0	125± 0.0	125± 0.0

Data represents mean values of three replicates ± SD

Therefore, LuL and LuA extracts showed strong antimicrobial activity against pathogenic bacteria, which are usually presented on the skin. In addition, the extract demonstrated a high content of phenolics and flavonoids that served the antimicrobial activity. Flavonoids are effective both in directly damaging the envelope of Gram-negative and Gram-positive bacteria [30]. Thus, this study implied that the LuL and LuA extracts contained both bacteriostatic and bactericidal effects.




Fig. 6 LuA extract at 500 mg/ml in DMSO



Fig. 7 LuL extract at 500 mg/ml dissolve in DMSO

3.6 Metal ion in the LuA and LuL

The results of the metals in the sample were as presented. The minerals in LuL were high in iron, zinc, and copper, also in LuA but in higher quantities. The research finding reviewed that a high-level supplement of trace metal not only improved growth performance but also reduced footpad lesions by improving the wound healing process via promotion of collagen synthesis, decomposition and organization, cell migration, matrix remodeling, angiogenesis, and regulation

of inflammation [31]. The role of each mineral in wound healing should be the subject of future studies.

Table 6 Metals in LuL and LuA by Inhouse Method Based on EPA 3052, by ICP-OES Technique

Test Item	Result		Unit
	LuL	LuA	
Arsenic (As)	0.9	2.04	mg/kg
Cadmium (Cd)	0.15	0.29	mg/kg
Copper (Cu)	7.25	25.34	mg/kg
Iron (Fe)	148	1332	mg/kg
Lead (Pb)	1.56	8.83	mg/kg
Mercury (Hg)	Not detected	Not detected	mg/kg
Zinc (Zn)	43.28	189	mg/kg

Discussion

Herbal medicines have become a popular form of therapy in developing countries. They are believed to be nontoxic, with little side effects compared to modern drugs. This is in accordance with the common use of *Luffa cylindrica* (L.) Roem in several countries worldwide for the traditional management of diseases. There was evidence from a few pharmacological investigations that *Luffa cylindrica* possessed anti-inflammatory, analgesic, antipyretic, hypoglycemic, antibacterial, antifungal, antiviral, anthelmintic, antioxidant, anticancer, hepatoprotective, antiemetic, wound healing, immunological, bronchodilation, reproductive effect, and in the treatment of cataract [13], [20], [22], [25].

Furthermore, based on our study, in order to compare the chemical composition, antibacterial, antioxidant, and wound healing activity of extract from dry leaves (LuL) compared with the ash dry leaves (LuA) of *Luffa cylindrica* (L.) Roem, it was found that LuL extract and LuA extract showed the physical-chemical effect, phytochemical effect, antioxidant activity, and antibacterial activity, including metal ion content. However, in comparison between the LuL extract and LuA extract of *Luffa cylindrica* (L.) Roem, there was the most quantitative phytochemical determination, such as the cardiac glycosides, alkaloids, phenolics, flavonoids, and triterpenoids effects presented in the LuL extract because the LuA sample was burned out into ash. This is in accordance with the study by [22], [31].

Further, the result of the antioxidant activity of LuA and LuL extracts was that the LuL had the most significant effect on the DPPH (1,1-diphenyl-2-picrylhydrazyl) radical scavenging activities, in accordance with preview studies by [32][24]. Free radicals are known to play a definite role in a wide variety of pathological manifestations. Antioxidants fight against free radicals and protect us from various diseases. They exert their action either by scavenging the reactive oxygen species or protecting the antioxidant defense mechanisms [21]. The electron donation ability of natural products can be measured by 2,2'-diphenyl-1-picrylhydrazyl radical (DPPH) purple-colored solution bleaching [8]. The method is based on scavenging of DPPH through the addition of a radical species or antioxidant that decolorizes the DPPH solution. The degree of color change is proportional to the concentration and potency of the antioxidants. A large decrease in the absorbance of the reaction mixture indicates significant free radical scavenging activity of the compound under test [34]. In the present study among all the fractions tested, *n*-butanol, chloroform and ethyl acetate showed significantly higher inhibition percentage and positively correlated with total phenolic content. Results of this study suggest that the plant extract contain phytochemical constituents that are capable of donating hydrogen to a free radical to scavenge the potential damage.

Antibacterial Activity of *Luffa* Leaf (LuL) Extract and *Luffa* Ash (LuA) Extract: It was revealed that LuA extract was more effective in antibacterial activity than LuL; this result had a similarity with studies by [17], [19], [21], and because the extract demonstrated a high content of phenolics and flavonoids that served the antimicrobial activity.

Furthermore, in this study, we have shown that a high-level supplement of trace metals not only improved growth performance but also reduced footpad lesions by improving the wound healing process via promotion of collagen synthesis, decomposition and organization, cell migration, matrix remodeling, angiogenesis, and regulation of inflammation. And in LuA, more metal ion items were detected, such as arsenic (As), cadmium (Cd), copper (Cu), iron (Fe), lead (Pb), mercury (Hg), and zinc (Zn). This is in accordance with the preview study by [24], [31], [33] The role of each mineral in wound healing should be the subject of future studies.

Conclusion

The results of the present study clearly indicated that the crude methanol extract of *Luffa cylindrica* did produce strong antimicrobial activity against pathogenic bacteria on the skin. In addition, the extract demonstrated a high content of phenolics and flavonoids that served as antimicrobial agents. Thus, this study implied that the activity of extracts from dry leaves (LuL) and ash-dry leaves (LuA) contained both bacteriostatic and bactericidal effects.

References

- [1] P. K. Singh, B. R. Choudhary, R. Singh, D. R. Bhardwaj, and M. Rai, "Developing sponge gourd [*Luffa cylindrica* (Roem.) L.] populations with a very high proportion of pistillate flowers," *Acta Hort.*, vol. 938, no. June 2014, pp. 491–493, 2012, doi: 10.17660/actahortic.2012.938.64.
- [2] I. O. Oboh and E. O. Aluyor, "*Luffa cylindrica*" vol. 4, no. August, pp. 684–688, 2009.
- [3] W. Pootakham *et al.*, "De novo assemblies of *Luffa acutangula* and *Luffa cylindrica* genomes reveal an expansion associated with substantial accumulation of transposable elements," *Mol. Ecol. Resour.*, vol. 21, no. 1, pp. 212–225, 2021, doi: 10.1111/1755-0998.13240.
- [4] O. Folake Lucy and O. Babajide Abidemi, "Food Value and Phytochemical Composition of *Luffa cylindrica* Seed Flour," *Am. J. Biochem.*, vol. 2, no. 6, pp. 98–103, 2012, doi: 10.5923/j.ajb.20120206.02.
- [5] M. W. K. P. Silva, R. H. G. Ranil, and R. M. Fonseka, "*Luffa cylindrica* (L.) M. Roemer (Sponge Gourd-Niyan wetakolu): An Emerging High Potential Underutilized Cucurbit," *Trop. Agric. Res.*, vol. 24, no. 2, p. 186, 2015, doi: 10.4038/tar.v24i2.8004.
- [6] J. M. Davis, "Luffa Sponge Gourd Production Practices for Temperate Climates," vol. 29, no. 4, pp. 263–266, 2016.
- [7] A. Vandkar, R. Shaikh, R. Thombare, and A. Chivate, "Formulation and Analysis of Herbal Face Wash Using *Luffa Cylindrica* Seeds Oil Extract As A Soap Base," vol. 8, no. May, 2018.
- [8] M. Bisiriyu and R. Salau, "Assessment of The Conversion of Luffa Cylindrical Seed Oil into Biodiesel using Cao/Metakaolin as Heterogenous Catalyst," 2021, [Online]. Available: <http://repository.futminna.edu.ng:8080/xmlui/handle/123456789/7755>
- [9] S. Partap, A. Kumar, K. Sharma, and K. K. Jha, "*Luffa Cylindrica* : An important

- medicinal plant,” *J. Nat. Prod. Plant Resour*, vol. 2012, no. 1, pp. 127–134, 2012, [Online]. Available: <http://scholarsresearchlibrary.com/archive.html>
- [10] C. Nat, “1 Supplementary Table,” vol. 56, pp. 1–6, 2022.
- [11] I. Aroma *et al.*, “Volatile Compounds Governed by Single Recessive Gene,” 2022.
- [12] R. E. View and P. A. Per, “zyxwvu zyxwv,” vol. 9, pp. 159–168, 1995.
- [13] M. A. Azeez, O. S. Bello, and A. O. Adedeji, “Journal of Medicinal Plants Studies Traditional and medicinal uses of *Luffa cylindrica* : a Review,” vol. 1, pp. 102–111, 2013.
- [14] P. K. Kokkula, R. Imadi, P. Garrepally, and S. Nikitha, “Anthelmintic Activity of Some Medicinal Plants : A Short Review,” vol. 9, pp. 605–611, 2019.
- [15] A. E. Al-snafi, “Traditional uses of Iraqi medicinal plants (part 2),” vol. 02, no. June, pp. 22–41, 2021.
- [16] P. Raut, S. Dhawale, D. Kulkarni, S. Pekamwar, and S. Shelke, “Pharmacodynamic findings for the usefulness of *Luffa cylindrica* (L .) leaves in atherosclerosis therapy with supporting antioxidant potential,” vol. 8, 2021.
- [17] N. Agarwal and A. Jindal, “Herbal Components as an Advantageous Remedy for Pimple and Acne in Face-Wash : A Systemic Review,” vol. 13, no. 01, pp. 1–20, 2023, doi: 10.24092/CRPS.2023.130101.
- [18] J. Sainkhediya and C. Chouhan, “Study of Effect of Temperature , Water , Light and Darkness on Seed Germination in *Luffa aegyptiaca*,” vol. 5, no. 1, pp. 1–5, 2023.
- [19] R. G. Polytechnic, O. State, and O. State, “Comparative Study of Antimicrobial Potency and Phytochemical Analysis of Methanolic Extracts of the Leaf and Flower of *Luffa cylindrica*,” vol. 4, no. 8, pp. 7–11, 2014.
- [20] I. T. Oyeyemi, O. M. Yekeen, P. O. Odusina, and T. M. Ologun, “Genotoxicity and antigenotoxicity study of aqueous and hydro-methanol extracts of *Spondias mombin* L ., *Nymphaea lotus* L . and *Luffa cylindrical* L . using animal bioassays,” vol. 8, no. 4, pp. 184–192, 2015, doi: 10.1515/intox-2015-0028.
- [21] A. E. Al-snafi, “Constituents and pharmacology of *Luffa cylindrica*-A review Constituents and pharmacology of *Luffa cylindrica* - A review,” no. September, 2019.
- [22] Al-Snafi AE, “Constituents and pharmacology of *Luffa cylindrica*- A review,” *IOSR J. Pharm.*, vol. 9, no. 9, pp. 68–79, 2019, [Online]. Available: https://www.researchgate.net/profile/Ali-Al-Snafi/publication/338902020_Constituents_and_pharmacology_of_Fumaria_officinalis-A_review/links/5e31e6e3a6fdccd9657672a7/Constituents-and-pharmacology-of-Fumaria-officinalis-A-review.pdf
- [23] H. Thi, H. Truong, T. T. Phan, H. Thi, and K. Le, “CORRELATIONS BETWEEN FRUIT ’ S YIELD WITH HORTICULTURAL TRAITS AND INHERITANCE OF (LUFFA CYLINDRICA),” vol. 128, no. 3, pp. 17–26, 2019, doi: 10.26459/hueuni-jard.v128i3B.4973.
- [24] A. S. Abitogun, A. O. Ashogbon, R. G. Polytechnic, O. State, and O. State, “Nutritional Assessment and Chemical Composition of Raw and Defatted *Luffa cylindrica* Seed Flour,” no. Ologhobo 1980, 2010.
- [25] U. M. Dhanalekshmi, G. Poovi, N. Kishore, M. D. Raja, and P. N. Reddy, “Enhancement of Cutaneous Wound Healing by Ethanolic Extract of *Luffa cylindrica* linn . Leaf and Flower in Wistar rat model” vol. 2, no. 3, pp. 85–91, 2010.
- [26] L. Pontoni *et al.*, “Modified Sample Preparation Approach for the Determination of the Phenolic and Humic-Like Substances in Natural Organic Materials by the Folin Ciocalteu Method,” *J. Agric. Food Chem.*, vol. 65, no. 48, pp. 10666–10672, 2017, doi: 10.1021/acs.jafc.7b04942.
- [27] Y. Wu and D. Wang, “Structural characterization and DPPH radical scavenging activity of

- an arabinoglucogalactan from *Panax notoginseng* root,” *J. Nat. Prod.*, vol. 71, no. 2, pp. 241–245, 2008, doi: 10.1021/np070323+.
- [28] M. Silva, N. V Jacobus, C. Deneke, and S. L. Gorbachl, “Silva, M., Jacobus, N. V., Deneke, C., & Gorbach, S. L. (1987). Antimicrobial substance from a human *Lactobacillus* strain. *Antimicrobial Agents and Chemotherapy*, 31(8), 1231–1233. doi:10.1128/aac.31.8.123,” vol. 31, no. 8, pp. 1231–1233, 1987.
- [29] I. Bhattacharjee, S. K. Chatterjee, S. Chatterjee, and G. Chandra, “Antibacterial potentiality of *Argemone mexicana* solvent extracts against some pathogenic bacteria,” *Mem. Inst. Oswaldo Cruz*, vol. 101, no. 6, pp. 645–648, 2006, doi: 10.1590/S0074-02762006000600011.
- [30] Donadio G, Mensitieri F, Santoro V, Parisi V, Bellone ML, De Tommasi N, Izzo V, Dal Piaz F. Interactions with Microbial Proteins Driving the Antibacterial Activity of Flavonoids. *Pharmaceutics*. 2021 May 5;13(5):660. doi: 10.3390/pharmaceutics13050660. PMID: 34062983; PMCID: PMC8147964.
- [31] Juxing Chen, Guillermo Tellez, Jeffery Escobar & Mercedes Vazquez-Anon Impact of Trace Minerals on Wound Healing of Footpad Dermatitis in Broilers. *Scientific Reports*, 7: 1894, | DOI:10.1038/s41598-017-02026-2 1
- [32] L. L. Roem, “Role of some nutrients on in vitro pollen germination and tube development of *Luffa cylindrica* (L.) Roem.” *Ann. Plant Sci.*, no. 3, 2014.
- [33] T. Zhang *et al.*, “Long-read sequencing and de novo assembly of the *Luffa cylindrica* (L.) Roem. genome,” *Mol. Ecol. Resour.*, vol. 20, no. 2, pp. 511–519, 2020, doi: 10.1111/1755-0998.13129.
- [34] A. Stephen and R. Suresh, “Nutritive and Therapeutic Values of Vegetables from the Markets of Chennai, Tamil Nadu , India.” vol. 3, no. 11, pp. 546–561, 2015.