

Management strategies for Charcoal rot of Sesame: A Review

Abstract

Sesame (*Sesamum indicum* L.) commonly known as *til* is one of the important oilseed crop having good nutritional, biomedical and religious value. It is also called as “Queen of oilseeds” and among the oilseed crops, sesame ranks first for its higher oil content with a dietary energy of 6335kcal/kg in seeds. The crop is attacked by several pathogens causing serious diseases and act as major damaging factor to sesame crop cultivated in the whole world with severe losses of 7million tones yearly. The important diseases of sesame include charcoal rot (*Macrophomina phaseolina*, Fusarium wilt (*Fusarium oxysporum*), Phytophthora blight (*Phytophthora parasitica*) and phyllody (phytoplasma). Charcoal rot is caused by *Macrophomina phaseolina* has been a major threat to the successful cultivation of sesame in Haryana which causes about 5-100% loss. The pathogen being facultative in nature survives as microsclerotia in soil and infested plant debris that serve as the primary source of inoculum and have been found to persist within the soil up to three years. The pathogen not only persists in soil as saprophyte associated with other soil microorganisms but also transmitted through seed. Hence, it requires different management approaches to overcome this menace. So, keeping in view the present scenario ecofriendly management must be done by the use of botanicals and bio agents.

Key Words: sesame, charcoal rot, pathogen, disease, *Macrophomina phaseolina*

Introduction

Charcoal rot caused by *Macrophomina phaseolina* (Tassi) Goid by is one of the most important diseases of sesame in Haryana. The pathogen is widely distributed and highly destructive right from the sowing to all stages of crop growth ((Dinakaran and Mohammed, 2001).) The disease takes a heavy toll resulting in root and stems rot, loss of plant population and subsequently yield loss. Scanty systematic information is available on eco-friendly management of this disease in Haryana, hence the present review have a direct or indirect bearing on the investigation carried out in the studies.

The Host

Sesame (*Sesamum indicum* L.) is an important annual oilseed crop belongs to the family Pedaliaceae. Sesame is grown mainly for seed purpose which has a good quality food, nutrition, bio-medicine, health care and religious value. Due to potent medicinal properties in leaves, oil and seeds of sesame are used to cure hemorrhoids, ulcers, cough, asthma and many other diseases. Sesame is used in various industries like pharmaceutical industry for making pharmaceutical products, cosmetics industry for making soaps, sunscreen cream and insecticide industries for enhancing efficacy of pyrethrin and make it more effective to combat against insects. Sesame is a traditional and high valued crop due to its nutritional and medicinal properties. Despite of high nutritive, economic value and large acreage; production and productivity of crop is low in state as well as in the country.

Occurrence and economic importance of disease

Charcoal rot is one of the most common, widely distributed and destructive root and stem rot disease. In India, charcoal rot of sesame incited by *Macrophomina phaseolina* (Tassi) Goidanich was first reported from Uttar Pradesh (Mehta, 1951). Jain and Kulkarni (1965)

reported this disease from Jabalpur and Gwalior divisions of Madhya Pradesh. Now it is widely distributed in all sesame growing areas of Uttar Pradesh, Madhya Pradesh, Gujarat, Maharashtra, Haryana, Punjab, Bihar, West Bengal, Orissa, Tamilnadu, Karnataka and Kerala. Generally, pathogen of charcoal rot appears to be non-specific in nature and causes diseases in sesame, maize, sorghum, soybean, sunflower and other economically important crops and responsible for huge losses every year in India (Sinclair and Gray, 1972; Khare *et al.*, 1973). Generally, about 25% yield losses by charcoal rot in United States, Uruguay, Spain and Soviet Union has been observed but under favourable weather conditions for the growth and development of pathogen, total crop failure in specific areas have also been recorded (Tikhonov *et al.*, 1976). Murugesan *et al.*, (1978) reported that 1.8kg/ha sesame yield losses at every one per cent increase in charcoal rot disease intensity. The disease was reported from North and South America, Asia, Africa and Europe but more prevalent in subtropical and tropical countries with a semi-arid climate. Yu and Park (1980) found that *M. phaseolina* causes severe reduction in seed germination and seedling stand. Vyas (1981) observed charcoal rot is very serious and destructive disease in sesame growing areas and causes 5-100% yield losses. Hoes (1985) observed that this disease is a serious threat for crops especially in arid regions of the world. Maiti *et al.*, (1988) estimated yield loss of 57% at 40% of disease incidence. Dinakaran and Mohammed (2001) found charcoal rot caused by *M. phaseolina* is the most devastating disease among diseases of sesame. Chattopadhyay and Kalpana (2002) reported about 50% incidence of charcoal rot resulting in heavy yield losses in India. Deepthi *et al.*, (2014) estimated yield losses in sesame due to *M. phaseolina* at capsule formation stage and observed plant protected with fungicides had more number of capsules per plant and test weight of healthy than infected capsule. Min and Toyoto (2019) reported that the incidence of sesame charcoal rot was 10-30% which causes 10-75% yield losses in Myanmar.

The pathogen

Macrophomina phaseolina (Tassi) Goidanich is destructive, omnipresent and non-specific pathogen with vast host range. Tassi (1901) first identified the pycnidial stage of the fungus as *M. phaseolina*. Pathogen is best known as the cause of a disease aptly called "charcoal rot". Taubehaus (1913) discovered sclerotial stage *i.e.* *Sclerotium bataticola* (Taub) and identified the fungus as causal agent for charcoal rot of sweet potato in USA. Petrak (1923) established genus *Macrophomina* which was parasitic on sesame. Charcoal rot fungus is referred as *M. phaseolina* due to sclerotial-bearing mycelial stage and causes characteristic root rot disease. In India, Butler (1925) identified a similar sclerotial-bearing fungus and compared with isolates of Taubehaus and named as *Rhizoctonia bataticola* (Taub.) Butler; subsequently it was transferred to *Macrophomina* by Goidanich (1947). *M. phaseolina* (Tassi) Goidanich belongs to phylum *Ascomycota*, class *Dothideomycetes*, order *Botryosphaeriales*, family *Botryosphaeriaceae*, genus *Macrophomina* and species *phaseolina*. The hyphae of pathogen are branched at right angle with constriction and a septum just after constriction (Jakhar, 1997). Chaudhary *et al.*, (2001) reported that the pathogen continuously changes its nature and rapidly resistant cultivars become susceptible. The absence of known teleomorph has stalled its taxonomy for many years (Crous *et al.*, 2006). The severity of disease is directly related to the presence of viable sclerotia in the soil (Khan, 2007). Sapru and Mahajan, (2010) reported that it is a facultative parasite in nature. It is both externally and internally seed borne pathogen. It is considered to be seed, soil and stubble borne in nature which can survive for more than ten months under dry soil conditions in the form of sclerotia. The hyphae are septate and filiform. Initially hypha is hyaline

afterwards becoming grey to black and producing jet black oval to round microsclerotia of size 80-90 μ m in diameter (Mahmoud and Budak, 2011). Akhtar *et al.*, (2011) proved the necrotrophic behavior of pathogen in sesame and found seed infection efficiency of *M. phaseolina* was 100% with significant reduction in seed index. Chowdhury *et al.*, (2017) reported that infection stages of charcoal rot fungus *M. phaseolina* in sesame revealed a transition phase from biotrophy via BNS (biotrophy-to-necrotrophy switch) to necrotrophy.

Disease Symptoms

The most common symptom of charcoal rot is the sudden wilting of plants from top to downwards throughout crop growth period. Irregular and deep necrotic lesions tend toward hypocotyls and root surfaces were found in soybean (Ammon and Wyllie, 1972). Lesions coalesce to form larger patches on branches or other plant parts, leads to premature senescence and death of plant (Dhingra and Sinclair, 1973). Toxins production and sclerotia formation by pathogen in xylem play an important role in the dehydration of adult plant leads to wilting. *M. phaseolina* infected plants dry up and roots are decayed in a shredded appearance (Ilyas and Sinclair, 1974). Seedling blight, root rot and stem rot are the salient symptoms of charcoal rot of soybean (Sinclair, 1982). Irregular and deep necrotic lesions in hypocotyls and root surfaces were observed in chickpea (Singh and Mehrotra, 1982) and sorghum (Pedgaonkar and Mayee, 1990). Colonization of epidermal and cortical cells is followed by colonization of vascular cambium and phloem cells in chickpea (Singh *et al.*, 1990) in maize (Singh and Kaiser, 1994). Microsclerotia germinate and produce hypha which penetrate the epidermal cells and grows intercellular. This infection leads to cellular collapse, epidermal and cortical cells necrosis in the roots and hypocotyls of common bean (Mayek- Perez *et al.*, 2002). Fungal sclerotia overwinter on weeds and attack plant roots and causing decaying of fibrous roots and blackening of the stems (El-Fiki *et al.*, 2004). The fungus interrupts the function of xylem vessels causing wilting and premature death of plant (Gupta and Chauhan, 2005). *M. phaseolina* is worldwide distributed and infects different parts of plant and causes root rot, stem rot and seedling blight (Moi and Bhattacharyya, 2008). Pathogen mainly attacks at basal region of the plant and causes lesions on roots, stems, pods and seeds (Kumar *et al.*, 2011). The plant undergoes several morphological changes after attack of the pathogen including irregular lesions having grey centers with dark brown borders, death of nodes and finally wilting (Khalili *et al.*, 2016).

Evaluation of phytoextracts/botanicals

Management of charcoal rot pathogen is very difficult due to its long-term survival and wide host range. Minimum incidence of charcoal rot (3.3%) was found by the treatment of rhubarb followed by neem (6.7%) as compared to control(23.3%) under screen house conditions (El-Fiki *et al.* 2004). Tandel *et al.*, (2010) tried phytoextracts of eleven plant species against *M. phaseolina* of green gram and revealed that the onion bulb extract produced maximum inhibition of mycelia growth of root rot fungus. Kumar *et al.*, (2011) studied the integrated management of *Jatropha* root rot caused by *Rhizoctonia bataticola* by using different phytoextracts and revealed that neem extract produced the maximum growth inhibition of the pathogen at 20% concentration by 55.6%. Murugapriya *et al.*, (2011) tested some botanicals against *M. phaseolina* and observed that growth was inhibited with extracts of *Allium* spp. Management of charcoal rot

pathogen is very difficult due to its long-term survival and wide host range. It persists in the soil as saprophytes, associated with other soil organisms and transmitted from seed. Dhingani *et al.*, (2013) tested the bio-efficacy of phytoextracts of thirteen plant species against *Macrophomina phaseolina* causing root rot disease of chickpea under *in vitro* condition by using poison food technique and observed that maximum per cent inhibition was done by *Allium sativum* (73%) followed by *Curcuma longa*, 63.98 per cent inhibition. Hussain *et al.*, (2014) evaluated the efficacy of different plant extracts against the pathogen *Macrophomina phaseolina* causing charcoal rot of sunflower under *in vitro* conditions and revealed that *Allium sativum* produced maximum growth inhibition followed by *Parthenium hysterophorus* and *Cassia fistula* and minimum growth inhibition was produced by *Dalbergia sissoo*. Tabassum *et al.*, (2014) tried phytoextracts of different plant species and revealed that the seeds treated with extract of ginger rhizome showed maximum per cent inhibition. Meena *et al.*, (2014) tested the bio-efficacy of various plant extracts against *M. phaseolina* at different concentrations (10%, 15%, 20%) and revealed that rhizome extract of *Zingiber officinale* produced maximum growth inhibition (74.59%) of the pathogen. Iqbal *et al.*, (2014) tested the antifungal potential of twenty antagonistic plants against the pathogen *M. phaseolina* causing charcoal rot in mungbean and found that all the test plants inhibited the growth of *M. phaseolina* significantly to varying levels. The maximum inhibition was observed with *Carum copticum* (83.5%), *Azadirachta indica* (76.1%) and *Nigella sativa* (70.4%) at 10% concentration. The powders of *Olea europaea*, *Cassia angustifolia*, *Ocimum americanum* and *Lawsonia inermis* caused more than 50% reductions in the growth of the fungus. Savaliya *et al.*, (2015) evaluated the efficacy of phytoextracts of nine plant species under *in vitro* conditions using poison food technique against *M. phaseolina* and revealed that maximum growth of mycelium was inhibited by *Allium sativum*. Akanmu *et al.*, (2015) tested the inhibitory potential of four combined botanicals (*Ficus asperifolia*, *Momordica charantia*, *Anacardium occidentals*, *Psidium guajava*) on mycelial growth of *M. phaseolina* of cowpea and revealed that *Momordica charantia* was most effective in botanicals treatment alone and combined treatment of *Ficus asperifolia*, *Momordica charantia* and *Anacardium occidentals* was found most significant. Gojiya *et al.*, (2016) evaluated the different phytoextracts against *M. phaseolina* causing root rot of greengram and revealed that the extract of garlic cloves (*Allium sativum*) was proved excellent with maximum inhibiting (65.68%) mycelial growth of the pathogen. Khamari *et al.*, (2017) tried phytoextracts of thirty plant species against the pathogen causing root rot of sesame using water and methanol as solvents *in vitro* and revealed that the garlic registered maximum per cent mycelial inhibition followed by onion at all concentrations in methanol as well as in aqueous extract. Lakhran and Ahir (2018) tested the phytoextracts against the pathogen causing root rot in chickpea and observed that the garlic extract was found most effective in reducing root rot incidence followed by neem leaf extract. Khaire *et al.*, (2018) evaluated the efficacy of different phytoextracts of different plant species reduced mycelial growth of pathogen causing charcoal rot. Gwande *et al.*, (2018) tested bio-efficacy of botanicals against *Macrophomina phaseolina* causing dry root rot of safflower under *in vitro* conditions and observed that the maximum inhibition was done by *Allium sativum* (83.57%) followed by *Allium cepa* (72.09%) and *Vitex negundo* (65.85%). Thombre and Kohire (2018) evaluated the bioefficacy of different botanicals against *M. phaseolina* under *in vitro* conditions and observed that all the botanicals were found effective in reducing per cent mycelial growth of *M. phaseolina*.

Evaluation of antagonists

***In vitro* evaluation**

Raguchander *et al.*, (1997) tested the antagonistic activity of biocontrol agent isolates by using talc as a carrier against root rot of mungbean and revealed that maximum growth inhibition was produced by *Trichoderma viride*-III isolate. Dinakaran and Marimuthu (1998) studied the antagonistic efficacy of bio- agents against *Rhizoctonia bataticola* and observed that *Trichoderma viride* exhibited the highest *in vitro* inhibition of mycelial growth (54.4%) and sclerotial germination (75.8%). Bashar and Khatum (1999) observed that *Trichoderma hamatum* and *Trichoderma viride* were most potent antagonists against *Rhizoctonia bataticola*. Gupta *et al.*, (2002) evaluated the antagonistic efficacy of *Pseudomonas* strain biocontrol agent and observed that this strain showed a strong antagonistic effect against *M. phaseolina* .

Salunke *et al.*, (2008) tested the efficacy of bio agents against *Rhizoctonia bataticola* and observed that *Trichoderma viride* was found most effective against the pathogen followed by *T. harzianum* and *Pseudomonas fluorescens*. Rani *et al.*, (2009) reported the inhibitory action of bioagents viz., *Trichoderma viride*, *T. harzianum* and *Bacillus subtilis* against *Macrophomina phaseolina*. Rajput *et al.*, (2010) evaluated the efficacy of biological control agents against *Rhizoctonia bataticola* under *in vitro* conditions and revealed that *Trichoderma harzianum* caused maximum radial growth inhibition (69.62%) of pathogen. Jaiman and Jain (2010) tested the antagonistic activity of five bio-agents viz., *Trichoderma viride*, *T. harzianum*, *Pseudomonas fluorescens*, *Bacillus subtilis* and *Gliocladium virens* against the pathogen and observed that maximum inhibition (69.62%) was showed by *T. viride* followed by *T. harzianum* under *in-vitro* conditions. Kumar *et al.*, (2011) studied the integrated management of root rot disease of Jatropha caused by *Rhizoctonia bataticola* by using different biocontrol agents and revealed that mximum inhibition was produced by *Trichoderma harzianum*. Sreedevi *et al.*, (2011) evaluated the efficacy of five species of *Trichoderma* for biocontrol of *Macrophomina* causing root rot of groundnut under *in vitro* conditions by dual culture and bioassays methods and they revealed that among the five isolates *T. harzianum* (T3) and *T. viride* (T1) had maximum antifungal activity against the pathogen. *Trichoderma viride* and *T. harzianum* reduced the mycelial growth by 61.1% and 64.4% respectively.

Kumari *et al.*, (2012) studied the integrated management of root rot of mungbean incited by *M. phaseolina* by using biocontrol agents and reported that among the tested biocontrol agents *Trichoderma harzianum* was found most effective against the pathogen under *in vitro* conditions followed by *T. viride* and *T. polysporum*, *Pseudomonas fluorescens* was found least effective in reducing root rot incidence. Manjunatha *et al.*, (2013) evaluated the activity of isolates of bio-control agents against *Macrophomina phaseolina* caused dry root rot of chickpea and revealed that *T. viride* and *T. harzianum* were effective against the pathogen , as Tv-R isolate of *T. viride* was found to be most effective. Pf-4 isolate of *Pseudomonas fluorescens* was also found to be more effective than the other isolates. Arshad *et al.*, (2014) evaluated the antagonistic efficacy of seven species of *Trichoderma* named as *T. pseudokoningii*, *T. harzianum*, *T. reesi*, *T. koningii*, *T. hanatus*, *T. viride*, *T. aureoviride* against the pathogen and observed that maximum per cent inhibition was done by *T. harzianum* and overall reduction in mycelium growth was 45-65%. *Trichoderma harzianum* was found to be most effective against the pathogen followed by *T. aureoviride* and *T. hanatum*. Tetali *et al.*, (2015) studied the management of disease by using biocontrol agents and revealed that the best antagonistic action was shown by *Trichoderma viride*. The integrated management approach by using different *Trichoderma* spp. against charcoal rot of *vigna mungo* caused by *M. phaseolina* and among the tested bioagents, *Trichoderma viride* showed maximum mycelial growth inhibition of (77.77%) reported by Karthikeyan *et al.*, (2015). Meena and Pandey (2016) tested the antagonistic property of bio-agents

viz., *Trichoderma viride*, *T. harzianum*, *T. virens* and *Pseudomonas fluorescens* against *M. phaseolina* causing root rot of mungbean and observed that maximum inhibition was shown by *T. viride*. Lakhran and Ahir (2018) evaluated the efficacy of bio-control agents under *in-vivo* conditions and observed that *Trichoderma viride* was most effective against *Macrophomina phaseolina* followed by *T. harzianum*. Satpathi and Gohel (2018) tested the antagonistic potential of eight biocontrol agents against *M. phaseolina* causing root rot of sesame and revealed the *Trichoderma atroviride* showed strong antagonistic effect against the pathogen with highest growth inhibition (60.00%).

Thombre and Kohire (2018) evaluated the bioefficacy of bioagents against *M. phaseolina* under *in vitro* conditions and observed that all the bioagents exhibited fungistatic activity against *M. phaseolina* and inhibited mycelial growth of pathogen. Brahmabhatt (2018) studied the management of root and collar rot of okra caused by *Macrophomina phaseolina* by using different bio agents and observed that *Trichoderma viride* was the most effective with highest growth inhibition (73.06%) followed by *Trichoderma harzianum* (68.89%).

Effect of seed treatment with antagonists

Gupta *et al.*, (2002) evaluated the antagonistic efficacy of *Pseudomonas* strain biocontrol agent and observed that this strain showed a strong antagonistic effect against *M. phaseolina* and seed treatment with this strain increased the seed germination besides enhancing early seedling growth. Indra and Gayathri (2003) tested the efficacy of biocontrol agents in different carriers against root rot of blackgram caused by *M. phaseolina* and observed that the disease incidence was significantly reduced by 50% when treated with *Trichoderma* spp. alone or in combination with biofertilizer. The root length, shoot length, grain yield and nodulation significantly increased with the *T. harzianum* + *Rhizobium* treated seeds (22.26 cm), *T. viride* + *Rhizobium* treated seeds (36.93 cm), *T. harzianum* (gypsum formulation) + *Rhizobium* (661.66 kg/ha) and *T. viride* + *Rhizobium* (22.33 nodules/plant) respectively. Ramesh and Korikanthimath (2006) studied the management of root rot of groundnut caused by *M. phaseolina* by using bio control agents (*Trichoderma viride* and *Pseudomonas fluorescens*) and observed that seed treatment with biocontrol agents increased the germination percentage 11-23% and 54-82%, and reduced disease incidence significantly 40-58 and 55-77%. Kumari *et al.*, (2012) studied the integrated management of root rot of mungbean incited by *Macrophomina phaseolina* by using biocontrol agents and reported that among the tested biocontrol agents *Trichoderma harzianum* was found most effective against the pathogen in pots conditions followed by *T. viride* and *T. polysporum*, *Pseudomonas fluorescens* was found least effective in reducing root rot incidence.

Iqbal *et al.*, (2014) tested the antifungal potential of twenty antagonistic plants against the pathogen *M. phaseolina* causing charcoal rot in mungbean and found that the maximum seedling emergence was observed when the seeds were treated with *C. copticum* (83.3%) followed by *A. indica* (80.0%) at 10%. Khalili *et al.*, (2016) evaluated the antagonistic potential of *Trichoderma* isolates (T_{12} , T_2 , T_{10}) against *M. phaseolina* causing charcoal rot of soybean and observed that maximum growth inhibition was done by T_{12} isolate by (72.31%) and production assays by 63.36%. T_{12} isolate was also found most effective under *in vivo* conditions by seed treatment. Recently, effective integrated management practice for management of charcoal rot in sesame by treating the seeds with *T. viride* (5g/kg seed) was reported by studied the integrated management of charcoal rot of castor caused by *M. phaseolina* by using different bio-agents viz., *T. viride*, *T.*

harzianum and *P. fluorescens* and found that seed treatment of *T. harzianum* was superior in disease control with 26.71% disease incidence. **Adhikary et al., (2019)** studied the integrated management of root rot of sesame using biocontrol agents and reported that disease incidence can be checked by seed treatment with *T. viride* + *P. fluorescens* @ 10g/kg + soil application of *P. fluorescens* @ 2.5kg/ha + *T. viride* @ 2.5kg/ha enriched in 100 kg of FYM + neem cake @ 250kg/ha at sowing.

Effect of soil application

Khalili et al., (2016) evaluated the antagonistic potential of *Trichoderma* isolates (T₁₂, T₂, T₁₀) against *M. phaseolina* causing charcoal rot of soybean and observed that maximum growth inhibition was done by T₁₂ isolate by (72.31%) and production assays by 63.36%. T₁₂ isolate was also found most effective under *in vivo* conditions by soil inoculation. Gupta et al., (2018) studied the effective integrated management practice of charcoal rot in sesame by soil application of *T. viride* @ 2.5kg/hac. Dhawan et al., (2019) evaluated the effect of different biocontrol agents against dry root rot of clusterbean caused by *Rhizoctonia bataticola* and observed that soil application with *T. viride* @ 10g/kg seed was found most effective against the disease.

Effect of volatile and non-volatile compounds of bio agents

Dinakaran and Marimuthu (1997) tested the antifungal activity of nine mutants of *T. viride* against *M. phaseolina* and revealed that cell free culture filtrate of mutant M1 showed the highest *in vitro* inhibition of mycelial growth (54.4%) and sclerotial germination (75.8%) of *M. phaseolina* followed by M3. The mutants M3 and M8 produced maximum volatiles *in vitro*. Sreedevi et al., (2011) evaluated the efficacy of different isolates of bio-control agents against *M. phaseolina* causing root rot of groundnut and revealed that among all the isolates *T. harzianum* (T3) and *T. viride* (T1) showed best results in mycelial inhibition. The inhibition varied depending on the *Trichoderma* sp. Producing metabolites, *T. viride* inhibited fungal growth up to 69% and *T. harzianum* upto 79.7% in non-volatile and 47%, 64.7% in volatile metabolites, respectively.

Conclusion

The pathogen not only persists in soil as saprophyte associated with other soil microorganisms but also transmitted through seed. Hence, it requires different management approaches to overcome this menace. Moreover, the control of plant diseases using pesticides raises serious concerns about food and environmental safety and pesticide resistance, which have dictated the need for alternative disease management techniques. Charcoal rot disease in sesame used to appear every year in patches at farmers field at all growth stages in Haryana and usually go unnoticed and causes full amount loss to the crop at a time as the control of disease is not economical and feasible. The disease is inherited by minor genes, hence sources of resistance are not available to utilize in vertical resistance breeding programs. Moreover, the use of expensive and excessive pesticides that causes environmental pollution has become concern to public awareness for health hazard to all useful organisms including humans and animals. Keeping in view the importance of the disease in recent years in Haryana due to build up of high inoculum in soil and to avoid soil pollution through chemicals it has become necessary to test botanicals and bio-agents for their effectiveness against this menace

Conference disclaimer:

Some part of this manuscript was previously presented in the conference: 6th International Conference on Strategies and Challenges in Agricultural and Life Science for Food Security and Sustainable Environment (SCALFE-2023) on April 28-30, 2023 in Himachal Pradesh University, Summer Hill, Shimla, HP, India. Web Link of the proceeding: <https://www.shobhituniversity.ac.in/pdf/Souvenir-Abstract%20Book-Shimla-HPU-SCALFE-2023.pdf>

References

- Adhikary, N. K., Chowdhury, M. D., Begum, T., and Mallick, R. (2019). Integrated management of stem and root rot of sesame (*Sesamum indicum* L.) caused by *Macrophomina phaseolina* (Tassi) Goid. *International Journal of Current Microbiology and Applied Sciences.*, **8**(4), 22-24.
- Akanmu, A. O., Olawuyi, O. J., Bello, O. B., Akinbode, O. A., Aroge, T., Oyewole, B., & Odebode, A. C. (2015). Genotypic variations in the inhibitory potentials of four combined botanicals on mycelia growth of *Macrophomina phaseolina* of cowpea [*Vigna unguiculata* (L) Walp.]. *Tropical Plant Research.*, **2**(3):257-263.
- Akhtar, K. P., Sarwar, G. and Arshad, H.M.I. (2011). Temperature response, pathogenicity, seed infection and mutant evaluation against *Macrophomina phaseolina* causing charcoal rot disease of sesame. *Archives of Phytopathology and Plant Protection.*, **44**(4):320-330.
- Ammon, V. and Wyllie, T.D. (1972). Penetration and host-parasite relationships of *Macrophomina phaseolina* on *Glycine max*. *Phytopathology*, **62**: 743-744.
- Arshad, J., Laiba, A., Anila, B. and Amna, S. (2014). *In vitro* screening of *Trichoderma* species against *Macrophomina phaseolina* and *Fusarium oxysporum* f. sp. *lycopersici*. *Pakistan Journal of Phytopathology.*, **26**(1):39-43.
- Bashar, M. A. and Khatun, M. T. (1999). Rhizosphere and Rhizoplane microflora of jute (*Corchorus capsularis* L.) and their antagonistic potential against *Macrophomina phaseolina* (Tassi) Goid. *Bangladesh Journal of Botany.*, **28**:69-77.
- Brahmbhatt, A. T. D. A. (2018). Management of root and collar rot (*Macrophomina phaseolina* (Tassi) Goid.) of Okra (*Abelmoschus esculentus* (L.) Moench) through bioagents, oil cakes and fungicides. *Journal of Pharmacognosy and Phytochemistry.*, **7**(4):631-635.
- Butler, E.J. (1925). Identification of the fungi. *Technical and Scientific Service Bulletin (Botanical Section)*, **49**: 64-65.
- Chattopadhyay, C. and Kalpna, S. (2002). Combining viable disease control tools for management of sesame stem root rot caused by *Macrophomina phaseolina* (Tassi) Goid. *Indian Journal of Plant Protection*, **30**(2): 132-138.

- Chaudhary, M.H.Z., Sarwar, N. and Chaughati, F.A. (2001). Biochemical changes in chickpea plant after induction treatment with simple chemical for systemic resistance against *Ascochyta* blight in the field. *Journal of Chemical Society of Pakistan*, **23**:182-186.
- Chowdhury, S., Basu, A. and Kundu, S. (2017). Biotrophy-necrotrophy switch in pathogen evoke differential response in resistant and susceptible sesame involving multiple signaling pathways at different phases. *Nature Scientific Reports*, **7**(1):1-17.
- Crous, P.W., Slippers, B., Wingfield, M.J., Rheeder, J., Marasas, W.F.O., Philip, A.J.L., Alves, S.A., Burgess, T., Barber, P. and Groenewald, J.L. (2006). Phylogenetic lineages in the *Botryosphaeriaceae*. *Studies in Mycology*, **55**: 235-253.
- Deepthi, P., Shukla, C.S., Verma, K.P. and Reddy, E.S.S. (2014). Identification of charcoal rot resistant lines of *Sesamum indicum* and chemical management of *Macrophomina phaseolina*. *Medicinal Plants*, **6**(1):36-42.
- Dhawan, A., Kumar, S., Sharma, P. K., Chugh, R. K., Jain, L. K., Parewa, H. P., & Sood, S.(2019). Effect of different fungicides, organic amendments and bio-control agents on dry root rot of cluster bean [*Cyamopsis tetragonoloba* L. Taub] Caused by *Rhizoctonia bataticola* (Taub.) Butler.
- Dhingani, J. C., Solanky, K. U., and Kansara, S. S. (2013). Management of root rot disease [*Macrophomina phaseolina* (Tassi.) Goid] of chickpea through botanicals and oil cakes. *The Bioscan.*, **8**(3):739-742.
- Dhingra, O.D. and Sinclair, J.B. (1973). Location of *Macrophomina phaseolina* on soybean plants related to culture characteristics and virulence. *Phytopathology*, **63**: 934-936.
- Dhinkaran, D. and Marimuthu, T. (1998). Inhibition of *Macrophomina phaseolina* (Tassi) Goid by mutant of *Trichoderma viride*. *Journal of Oilseed Research.*, **12**:262-263.
- Dinakaran, D. and Mohammed, N. (2001). Identification of resistant sources to root rot of sesame caused by *Macrophomina phaseolina* (Tassi.) Goid. *Sesame and Safflower Newsletter*, **16**: 68- 71.
- El-Fikki All, El-Deep A. A., Mohmmad F. G., Khalifa M.M.A (2004). Controlling sesame charcoal rot by *Macrophomina phaseolina* under field conditions by using resistant cultivars, seed and soil. *Egypt Journal of Phytopathology.*, **32**(1-2):103-118.
- Gawande, P. N., Gholve, V. M., & Ghuge, S. B. (2018). *In vitro* Bio-Efficacy of Botanicals and Organic Amendments against Dry Root Rot of Safflower. *International Journal of Current Microbiology and Applied Sciences.*, **7**(11):2048-2062.
- Goidanich, G. (1947). A revision of the genus *Macrophomina phaseolina* petrak type species: *Macrophomina phaseolina*, **3**: 449-461.
- Gojiya, S., Akbari, L., and Chudasama, M. (2016). Evaluation of Phytoextracts Against *Macrophomina phaseolina* (Tassi) Goid Causing Root Rot of Greengram. *Advances*, 500.

- Gupta, C. P., Dubey, R. C. and Maheshwari, D. K. (2002). Plant growth enhancement and suppression of *Macrophomina phaseolina* causing charcoal rot of peanut by fluorescent *Pseudomonas*. *Biology and Fertility of Soils.*, **35**:399-405.
- Gupta, G.K. and Chauhan, G.S. (2005). Symptoms, identification and management of soybean diseases. *Technical bulletin*, **10**: 89-92.
- Gupta, K. N., Naik, K. R. and Bisen. R. (2018). Status of sesame diseases and their integrated management using *indigenous practices*. *International journal of chemical studies.*, **6**(2):1945-1952.
- Hoes, J.A. (1985). *Macrophomina phaseolina* causal agent of charcoal rot of sunflower and other crops. Agricultural Canada Research Statistics, Modern Manitoba.
- Hussain, M., Ghazanfar, M. U., Hamid, M. I., Zahid, M. A., Hussain, T., Raza, M., & Haq, A. U. (2014). *In vitro* evaluation of fungicides and plant extracts for suppressing mycelial growth of *Macrophomina phaseolina* causes charcoal rot of Sunflower. *International Journal of Agricultural Applied Sciences.*, Vol. **6**(2).
- Ilyas, M.B. and Sinclair, I.B. (1974). Effects of plant age upon development of necrosis and occurrence of intraxylem sclerotia in soybean infected with *Macrophomina phaseolina*. *Phytopathology*, **54**: 156-157.
- Indira, N., & Gayatri, S. (2003). Management of blackgram root rot caused by *Macrophomina phaseolina* by antagonistic microorganisms. *Madras Agric. J*, **90**, 490-494.
- Iqbal, U., Mukhtar, T., and Muhammad Iqbal, S. (2014). *In vitro* and *in vivo* evaluation of antifungal activities of some antagonistic plants against charcoal rot causing fungus *Macrophomina phaseolina*. *Pakistan Journal of Agricultural Sciences.*, **51**(3).
- Jaiman, R. K., Jain, S. C., & Pankaj, S. (2010). Field evaluation of fungicides, bioagents and soil amendments against root rot caused by *Macrophomina phaseolina* in cluster bean. *Journal of Mycology and Plant Pathology.*, **39**(1):74.
- Jakhar, S.S. (1997). Occurrence and control of Rhizoctonia species causing root rot of cotton (*Gossypium* sp.). Ph.d. Thesis, Haryana Agricultural University, Hisar, India, pp.25.
- Khaire, P. B., Hingole, D. G. and Padvi, S. A. (2018). Efficacy of different phytoextracts against *Macrophomina phaseolina*. *Journal of Pharmacognosy and Phytochemistry.*, **7**(3):1124-1126.
- Khalili, E., Javed, M. A., Huyop, F., Rayatpanah, S., Jamshidi, S. and Wahab, R. A. (2016). Evaluation of *Trichoderma* isolates as potential biological control agent against soybean charcoal rot disease caused by *Macrophomina phaseolina*. *Biotechnology and Biotechnological Equipment*, **30**(3):479-488.

- Khamari, B., Beura, S., Roy, A., and Ranasingh, N. (2017). Assessment of antifungal potency of some plant extracts against *Macrophomina phaseolina*, causing stem and root rot of sesame. *The Bioscan.*, **12**(4):2047-2051.
- Khan, S. N. (2007). *Macrophomina phaseolina* as causal agent for charcoal rot of sunflower. *Mycopathology.*, **5**(2):111-118.
- Khare, M.N., Sharma, H.C., Kumar, S.M. and Chaurasia, R.K. (1973). Current plant pathological problems of soyabean and their control. *Proceedings at the fourth All Indian Workshop on Soyabeans, GBPUAT, Pantnagar*, pp. 55-67.
- Kumar S., Aeron A., Pandey P., Maheshwari D. K. (2011). Ecofriendly management of charcoal rot and wilt diseases in sesame (*Sesamum indicum* L.) In: D. K. Maheshwari (ed.) *Bacteria in Agrobiolgy: Crop Ecosystems*, pp. 387-405.
- Kumar, S., Sharma, S., Pathak. D. V. and Beniwal. J. (2011). Integrated management of Jatropha root rot caused by *Rhizoctonia bataticola*. *Journal of Tropical Forest Science.*, **23**(1):35-41.
- Kumari, R., Shekhawat, K. S., Gupta, R. and Khokhar, M. K. (2012). Integrated management against root rot of mungbean incited by *Macrophomina Phaseolina*. *Journal of Plant Pathology and Microbiology.*, **3**:5.
- Lakhran, L., and Ahir, R. R. (2020). *In vivo* evaluation of different fungicides, plant extracts, biocontrol agents and organics amendmets for management of dry root rot of chickpea caused by *Macrophomina phaseolina*. *Legume Research-An International Journal.*, **43**(1):140-145.
- Mahmoud, A. and Budak, H. (2011). First report of charcoal rot caused by *Macrophomina phaseolina* in sunflower in Turkey. *Plant Diseases.*, **95**(2):223.
- Maiti, S., Hedge, M. R. and Chattopadhyay, S. B. (1988). *Handbook of Annual Oilseed Crop*. Oxford and IBH Publ. Co. Pvt. Ltd., New Delhi.
- Manjunatha, S. V., Naik, M. K., Khan, M. F. R., and Goswami, R. S. (2013). Evaluation of bio-control agents for management of dry root rot of chickpea caused by *Macrophomina phaseolina*. *Crop protection.*, **45**:147-150.
- Mayek-Perez, N., Garcia-Espinosa, R., Lopez-Castaneda, C., Acosta-Gallegos, J.A. and Simpson, J. (2002). Water relations, histopathology and growth of common bean (*Phaseolus vulgaris* L.) during pathogenesis of *Macrophomina phaseolina* under drought stress. *Physiological and Molecular Plant Pathology*, **60**(4): 185-195.
- Meena, B. and Pandey, R. N. (2016). *In vitro* evaluation of bio-control agents against *Macrophomina phaseolina* and *Rhizoctonia solani*. *Trends in Biosciences.*, **8**(3):669-671.

- Meena, P. N., Tripathi, A. N., Gotyal, B. S., and Satpathy, S. (2014). Bio-efficacy of phytoextracts and oil cakes on *Macrophomina phaseolina* (Tassi) causing stem rot disease of Jute, *Corchorus* spp. *Journal of Applied and Natural Science.*, **6**(2):530-533.
- Mehta, P. R. (1951). Observation on new and known diseases of crop plants. *Plant protection Bulletin.*, **3**:7-11.
- Min, Y.Y. and Toyota, K. (2019). Occurrence of different kinds of diseases in sesame cultivation in Myanmar and their impact to sesame yield. *Journal of Experimental Agriculture International*, **38**(4):1-9.
- Moi, S. and Bhattacharya, P. (2008). Influence of biocontrol agents on sesame root rot. *Journal of Mycopathological Research.*, **46**(1):97-100.
- Murugapriya, E., Alice. D. and Jayamani, P. (2011). Antifungal activity of botanicals and micro nutrients against *Macrophomina* lead blight in mungbean. *Journal of Food Legumes.*, **24**(2): 113-116.
- Murugesan, M., Shanmugam, N., Menon, P.P.V., Arokiaraj, A., Dharmo, K.P. and Kochubabu, M. (1978). Statistical arrangement of yield loss of sesame due to insect pests and diseases. *Madras Agriculture Journal*, **65**:290-295.
- Pedgaonkar, S.M. and Mayee, C.D. (1990). Stalk water potential in relation to charcoal rot of sorghum.
- Petrak, F. (1923). Mykologische Notizen VI. *Annales Mycologici*, **21**: 314-315. *Phytopathology*, **59**: 19.
- Raguchander, T., Rajappan, K., & Samiappan, R. (1997). Evaluating methods of application of biocontrol agent in the control of mungbean root rot. *Indian Phytopathology.*, **50**(2):229-234.
- Rajput, V. A., Konde, S. A. and Thakur, M. R. (2010). Evaluation of bio agents against chickpea wilt complex. *Journal of Soils and Crops.*, **20**(1):155-158
- Ramesh, R., & Korikanthimath, V. S. (2006). Management of groundnut root rot by *Trichoderma viride* and *Pseudomonass fluorescens* under rainfed conditions. *Indian Journal of Plant Protection.*, **34**(2): 239.
- Rani S. U. Udaykumar, R. and Christopher, D. J. (2009). Bio-efficacy of plant extracts and bio-control agents against *Macrophomina phaseolina*, *Annals of Plant Protection Sciences.*, **17**(2):389-393.
- Salunke, V. N. Armarkar, S. and Ingle, R. W. (2008). Efficacy of fungicides and antagonistic effect of bio-agents against *Rhizoctonia bataticola* isolates. *Annals of Plant Physiology.*, **22**(1):134-137.

- Sapru, T. and Mahajan, S.K. (2010). Production of phenolics by *Rhizoctonia bataticola* (Taub.) Butler during pathogenesis. *Environment Conservation Journal.*, **11**(1):31-36.
- Satpathi, A. K., & Gohel, N. M. (2018). Evaluation of Antagonistic Potential of Biocontrol Agents against *Macrophomina Phaseolina* (tassi) Goid. causing Stem and Root Rot of Sesame [*Sesamum Indicum* L.] under *In vitro* and *In vivo*. *International Journal of Current Microbiology and Applied Sciences.*, **7**(12):1748-1754.
- Savaliya, V. A., Bhaliya, C. M, Marviyaand, P. B. and Akbari, L. F. (2015). Evaluation of phytoextracts against *Macrophomina phaseolina* (Tassi) Goid. Causing root rot of sesame. *Journal of Biopesticides.*, **8**(2):116-119.
- Sinclair, J.B. (1982). Compendium of soybean diseases 2nd edition *American Phytopathology Society: St. Paul*, pp. 104.
- Sinclair, J.B. and Gray, L.E. (1972). Three fungi that can reduce yields. *Dillinois Research.*, **14**:5.
- Singh, P.J. and Mehrotra, R.S. (1982). Penetration and invasion of gram roots by *Rhizoctonia bataticola*. *Indian Phytopathology*, **35**: 336-338.
- Singh, R.D.N. and Kaiser, S.A.K.M. (1994). Histopathological study of roots and stalk of maize plant invaded by the charcoal rot pathogen *Macrophomina phaseolina*. *Advances in Plant Science*, **7**:125-130.
- Singh, S.K., Nene, Y.L. and Reddy, M.V. (1990). Influence of cropping systems on *Macrophomina phaseolina* population in soil. *Plant Disease*, **74**: 812-814.
- Sreedevi, B., Charitha Devi, M., & Saigopal, D. V. R. (2011). Isolation and screening of effective *Trichoderma* spp. against the root rot pathogen *Macrophomina phaseolina*. *Journal of Agricultural Technology.*, **7**(3):623-635.
- Tabassum, A., Bhale, M. S., Pandey, A. K., and Rangnatha, A. R. G. (2014). Efficacy of phytoextracts in the management of sesame seeds associated *Macrophomina phaseolina*. *Review Paper Advances in the cultivation technology of tropical mushrooms in India.*, 120-135, 184.
- Tandel, D. H., Sabalpara, A. N., Pandaya, H. V. and Naik, R. M. (2010). Efficacy of phytoextracts against *Macrophomina phaseolina* causing leaf blight of green gram. *International Journal of Pharma and Bio Science.*, **1**(2) p1.
- Tassi, A. (1901). *Novae Micromycetum Species*, Series III Bulletin, Laboratory Orto Botanical University, Siena, **4**: 7-12.
- Taubenhaus, J. J. (1913). The black rot of sweet potato. *Phytopathology*, **3**: 161-164.

Tetali, S., Karpagavalli, S. and Pavani, S.L. (2015). Management of dry root rot of blackgram caused by *Macrophomina phaseolina* (Tassi) Goid. using bio agent. *Plant Archives.*, **15**(2):647-650.

Thombre, B. B. and Kohire, O. D. (2018). *In vitro* efficacy of bio agents and botanicals against *Macrophomina* blight of mungbean caused by *Macrophomina phaseolina* (Tassi) Goid. *International Journal of Chemical Studie.s*, **6**(2):3063-3066.

Tikhonov, O.I., Nedelko, O.K. and Persestova, T.A. (1976). Methods for pathogenicity tests for seed borne *Macrophomina phaseolina* isolated from different hosts. *Phytopathology*, **88**(3): 234- 237.

Vyas, S. C. (1981). Diseases in *Sesamum* in India and their control. *Pesticides*, **15**:9-10.

Yu, S. and Park, J.S. (1980). *Macrophomina phaseolina* detected in seeds of *Sesamum indicum* and its pathogenicity. *Korean Journal of Plant Protection*, **19**: 135-140.