

Assessment of heavy metals in chicken feeds sold in Ijebu-Jesa, Osun state.

Comment [MF1]: The title must be unique for increase paper citation ; I suggest to be; An evaluation of the presence of heavy metals in the poultry feeds marketed in Ijebu-Jesa, Osun state.

ABSTRACT

Poultry farming is one of the most important aspects of agriculture with commercial layers and broilers contributing tremendously in meeting the upward protein demand of the increasing population through eggs and meats. Supplementation of some essential metals such as copper (Cu), zinc (Zn) and manganese (Mn) in chickens' diets is of great importance. Four brands (starter, grower, layer and finisher) of two commercial feeds (Animal care and Top feeds) commonly used by poultry farmers in Osun state were analysed for cadmium (Cd), Cobalt (Co), Copper (Cu), Iron (Fe), Manganese (Mn), Nickel (Ni), Lead (Pb), Chromium (Cr), Zinc (Zn). The feeds were purchased from different retail stores at Ijebu-Jesa in Osun State. The feeds were ashed, digested, and subsequently analyzed for metal contamination using Atomic Absorption Spectrometer (AAS). The starter feed type consistently had the lowest concentration of the essential elements. The essential elements (zinc, iron, manganese and copper) were very low in the feed. Therefore, the nutritive values of the feed as estimated from the concentrations of the essential elements were very low. This shows that supplements were not added to the feeds as should have been expected. However there were very high concentrations of lead in the feed samples. This could be attributed to anthropogenic sources of lead pollution in the environment especially fossil fuels. There is a great need to adopt the alternative renewable energy sources such as biodiesel and bioethanol.

Comment [MF2]: Better to be: One of the most crucial areas of agriculture is poultry production, with commercial layers and broilers making a significant contribution to supplying the rising need for protein from the growing population through eggs and meats. It's crucial that hens have enough of certain necessary metals like copper (Cu), zinc (Zn), and manganese (Mn) in their meals. We examined the levels of cadmium (Cd), cobalt (Co), copper (Cu), iron (Fe), manganese (Mn), nickel (Ni), lead (Pb), chromium (Cr), and zinc (Zn) in four brands of two commercial feeds (Animal care and Top feeds) that are often used by chicken farmers in Osun state.

Comment [MF3]: Better to be: The meals were bought at Ijebu-Jesa, Osun State, from several retail establishments. Atomic Absorption Spectrometer (AAS) analysis was done after the feeds had been ashed, digested, and examined for metal contamination. The concentration of the key components was consistently lowest in starting feed. The feed had relatively little of the necessary components (copper, zinc, iron, and manganese). Because of this, the projected nutritional values of the feed based on the concentrations of the necessary components were extremely low. This demonstrates that supplements were not provided to the diets as was reasonable to expect. The feed samples, however, had very high levels of lead.

Comment [MF4]: Anthropogenic sources of lead contamination in the environment, particularly fossil fuels, may be to blame for this. Adoption of alternative renewable energy sources like biodiesel and bioethanol is highly necessary.

Keywords

Poultry, Grower, Heavy metals, chicken feeds, contamination

Introduction

Poultry farming is one of the major aspects of agriculture with commercial layers and broilers contributing tremendously in meeting the upward protein demand of the increasing population through eggs and meats. Supplementation of some essential metals such as copper (Cu), zinc (Zn) and manganese (Mn) in chickens' diets is of great importance. Copper prevents anaemia,

while Zn and Mn act as catalysts in many enzymatic and hormonal reactions that are related with growth, immunity and skeletal integrity (McDowell, 1992). Supplementation of Cu, Zn and Mn at 8, 40 and 60 ppm ($\mu\text{g/ml}$) respectively was recommended in broiler diets by NRC (1994) majorly in term of growth. However, Cu deficiency in birds can lead to rupture of the aorta. Diets deficient in Zn cause retarded growth, shortening and thickening of leg bones and enlargement of the hock joint, poor feathering, anorexia and mortality. Chicks hatched from Zn-deficient hens are weak, while a deficiency of Mn in the diet of chickens is one of the causes of perosis. Chickens are one of the main sources of protein for the south east population especially in IjebuJesa, Osun state where there are varieties of poultry farms and abundant market. The chickens are nourished by the feed they consume which as a result of heavy metal pollution in the country might be poisonous and therefore detrimental to the health of the human population that consumes the chickens. There is insufficient data as to the nutritive content of feeds consumed by chickens in the country and possible contamination of the feeds by the nutritive elements. So far no work has been carried out to cover the south west. Therefore this study was carried out to determine the concentration of zinc, iron, manganese, copper, lead, cadmium, nickel and cobalt in chicken feeds obtained in the south western part of Nigeria.

Environmental contamination and pollution with heavy metals, has taken a global dimension. Heavy metals have been reported in water; soil; food and even in the atmosphere at varying concentrations: at trace and /or toxic levels, in different parts of the world. Heavy metals occur in all foods as natural or inherent component of plant and animal tissues and fluid along cycles of food chain and also may be present as a result of contamination or deliberate addition. Nickel is a transition metal with an atomic structure very similar to that of cobalt. Although it is not normally added to chicken diets, but nickel has been detected in liver, kidney and muscle of broilers (Coleman *et al.*, 1992). Evidence of nickel deficiencies in chicks has been reported (NAS, 1980) and dietary nickel levels of 0.1 – 0.3 parts/106 or $\mu\text{g/ml}$ dry weight are considered adequate in poultry diets (Puls, 1988).

Heavy metals are ubiquitous and are being released continuously from man-made sources into the aquatic and terrestrial ecosystems, threatening the health of man and animals (Aschner, 2002; Abulude *et al.* 2006). They are potentially dangerous due to their toxicity, bioaccumulation and biomagnification abilities when found within living tissue (Ayciceket *et al.*, 2008), and are stored more quickly than they are excreted. The increase in urbanization, industrialization and agricultural activities has been shown to release heavy metals into the environment (Falandysz *et al.*, 2005). In early 2010, there was an incidence of heavy metal poisoning in Zamfara State, Nigeria due to indiscriminate mining by the locals (WWPPR, 2010).

Toxic metals are possible environmental pollutants domestic layer have the capacity to cause health problems in human selected poultry farm. Many studies showed that elements used in making poultry feed such as small grains (wheat, rice, feed deposit in body and cause harmful impacts. In case maize, millet, sorghum barley, dried sea food, blood of poultry industry deposition of heavy metals in body of animals, broken pulses and different vitamins) and heavy

Comment [MF5]: One of the most important areas of agriculture is poultry production, with commercial layers and broilers making a significant contribution to satisfying the rising protein needs of the expanding population through eggs and meats. It is crucial that certain critical metals, such as copper (Cu), zinc (Zn), and manganese (Mn), be supplemented in hens' diets. While Zn and Mn serve as catalysts in several enzymatic and hormonal processes associated with development, immunity, and skeletal integrity, copper prevents anemia (McDowell, 1992).

Comment [MF6]: According to NRC (1994), broiler diets should contain supplements of Cu, Zn, and Mn at concentrations of 8, 40, and 60 ppm (g/ml), respectively. But in birds, aortic rupture can result from a copper shortage. Zn deficiency in the diet slows development, thickens and shortens the leg bones, enlarges the hock joint, causes poor feathering, anorexia, and even leads to death...

Comment [MF7]: Because of the heavy metal pollution in the nation, the feed that the hens eat may be hazardous and thus harmful to the health of the human population that eats the chickens. The nutrient composition of the feeds used by hens in the nation and any potential nutrient contamination of the feeds are not well known. The south west hasn't been covered...

Comment [MF8]: All meals include heavy metals since they are a natural or ingrained part of plant and animal tissues and fluid along the food chain cycles. They can also be present due to contamination or intentional inclusion. The atomic structure of nickel, a transition metal, is strikingly similar to that of cobalt. Nickel has been found in broilers' liver, kidney, and muscle, despite the fact that it is not normally added to chicken diets...

Comment [MF9]: Evidence of nickel deficiency in chicks has been observed (NAS, 1980), and dietary nickel levels of 0.1 to 0.3 parts/106 or g/ml dry weight are regarded sufficient in chicken diets (Puls, 1988).

Comment [MF10]: The health of people and animals is at risk due to the pervasiveness of heavy metals and their ongoing release from man-made sources into aquatic and terrestrial ecosystems (Aschner, 2002; Abulude *et al.*, 2006). They are potentially harmful because they are stored more quickly than they are eliminated and have the ability to bioaccumulate and biomagnify when...

broiler were result of their excessive use in poultry feed. Poultry feed is the combination of foodstuff and concentration of these toxic metals were also higher in Pre-mixers. Pre-mixers are combination of cadmium, zinc, copper, Lead, Chromium, Nickel, Barium, different heavy metals manufactured locally on demand Cobalt, Strontium, Titanium, Mercury and Silver were and their concentrations were not known to the poultry conducted. Samples of broiler feed (foodstuff + Pre-mixers) were with high concentrations of these heavy metals than and tissue of muscles, liver and skin were collected from the National Hygienic Standards. Similarly litter samples were amount of aluminum and arsenic concentrations were collected from each poultry farm. So, contamination of poultry products with heavy metals may result from the feed and water consumed by the chickens. The present study is therefore design to ascertain the possibility of contamination of chicken feeds with heavy metals.

Materials and Methods

Study Area

The study was conducted in Ijebu-jesa the capital of Oriade Local Government area in Osun State of Nigeria. It is a commuter city with connections to Ekiti State on one side, Ondo State on another and it has a border with the famous Ilesa the surrounding town's area Iwoye-jesa, Iloko-jesa, Ere and Ijeda. This city also serves as the main route to the Osun State College of Technology in Esa-Oke. This city also has a private Polytechnic approved by the National Board for Technical Education (NBTE) Nigeria.

Ijebu-jesa which hitherto before now was known and called IjebuEgboro is the ancient historic town in the east of Osun State, Nigeria occupies a strategic position in Ijesaland. In Ijesa division, it is the next most important town politically and in term of history. Her Oba is the next to the OwaObokun of Ijesaland.

The town is situated eight kilometers north of Ilesa and about 128 kilometers east of Ibadan. It lies approximately on latitude 7.45 degrees north within the rain forest belt and so offers opportunity for farming on a large scale. The people are the core of Ijesa and are noted for their dogged industry.

Sample Collection

Comment [MF11]: Toxic metals are potential environmental contaminants that can have a negative impact on human health in poultry farms that raise only humane chickens. Numerous studies have demonstrated the detrimental effects of ingredients used in poultry feed, such as small grains (wheat, rice, and feed), on the body. When heavy metals from the poultry industry's blood are deposited in the bodies of animals, broken pulses and other vitamins) and heavy broilers are the result of their excessive usage in poultry feed. Poultry feed is a mixture of foodstuffs and pre-mixers had greater concentrations of these hazardous metals.

Comment [MF12]: Cadmium, zinc, copper, lead, chromium, nickel, barium, and other heavy metals are combined to make pre-mixers, which are produced locally and on demand. The poultry experiment included cobalt, strontium, titanium, mercury, and silver, although its amounts were unknown. High quantities of these heavy metals were found in samples of broiler feed (food + pre-mixers), which were also obtained from the National Hygienic Standards along with muscle, liver, and skin tissue. From each chicken farm, samples of litter with similar quantities of aluminum and arsenic were taken. Therefore, the water and feed that the birds ingest may contaminate poultry products with heavy metals. Determining if heavy metal contamination of chicken feeds is a possibility is the purpose of the current investigation.

Comment [MF13]: Re-write to reduce the plagiarism

Comment [MF14]: Re-write

All the plastic and glass wares were washed with detergent and rinsed with water before immersion in 10% nitric acid solution. They were finally rinsed with deionized water. Analytical grade reagents and deionized water were used throughout the analysis. Sampling Four brands (Starter, Grower, Layer and Finisher) of two feeds (Animal care and Vital feed coded as A, B, C, D, E, F, G and H respectively) available in IjebuJesa Metropolis were purchased from different locations within the metropolis.

Comment [MF15]: Re-write too

Sample Preparation

Dried sample weighed 2.0g was placed in crucibles. Conc. nitric acid (1cm³) was added as ashing aid and then pre-ashed by placing the crucible on a heater until the content charred. The pre-ashed samples were then transferred into a muffle furnace at a temperature of 480°C for 2-3hrs until a constant weight is obtained, after which they were allowed to cool. The cooled samples were dissolved using 5cm³ of 30% HCl and then filtered using Whatman filter papers. The filtrates were individually poured into 50cm³ volumetric flask and made up to the mark with deionized water. The sample solution was then kept in sample bottles for further analysis (Okoye et al., 2011).

Sample Analysis

Each brand of the same feed sample obtained from different locations, mixed, homogenized thoroughly, and 1g each of chick mash, growers mash, layers mash, broiler starter and broiler finisher was weighed using Mettler weighing balance. Each sample was then transferred into a 50ml Kjeldahl digestion flask, 10ml of concentrated nitric acid were added and then 2ml of perchloric acid were also added. The content was swirled gently and digested at about 3700 C heat first in a digestion block. Increasing the heat slowly to about 4500 C for 15min, after the appearance of white fumes, the digested samples were allowed to cool. The samples were dissolved using 10ml of distilled water and then filtered using Whatman filter papers. The filtrates were poured individually into 50ml prewashed sample bottles; the samples were analyzed using Atomic Absorption Spectrophotometry (AAS).

RESULT

Table 1: Heavy Metals of analysis of animal Feed sample A (Animal care). Concentration of Heavy Metals in animal feed was presented in (mg/kg).

Sample	Cd	Co	Cu	Fe	Mn	Ni	Pb	Cr	Zn
Starter	2.13±0.02	1.67±0.01	3.38±0.01	13.32±0.06	25.01±0.01	1.37±0.01	0.27±0.06	1.42±0.01	16.30±0.01
Grower	1.60±0.07	1.67±0.01	2.70±0.01	08.79±0.06	25.01±0.01	1.37±0.01	0.80±0.01	0.47±0.01	32.61±0.05
Layer	1.60±0.07	1.67±0.01	3.38±0.01	10.96±0.01	18.88±0.06	1.37±0.01	0.53±0.06	1.42±0.01	38.04±0.06
Finisher	2.13±0.02	2.51±0.12	2.70±0.01	10.96±0.01	18.88±0.06	2.06±0.01	0.27±0.06	1.42±0.01	21.74±0.01

Table 2: Heavy Metals of analysis of animal Feed sample B(Vital feed). Concentration of Heavy Metals in animal feed was presented in (mg/kg).

Sample	Cd	Co	Cu	Fe	Mn	Ni	Pb	Cr	Zn
Starter	2.13±0.02	1.67±0.01	3.38±0.01	13.32±0.06	25.01±0.01	1.37±0.01	0.27±0.06	1.42±0.01	16.30±0.01
Grower	1.60±0.07	1.67±0.01	2.70±0.01	08.79±0.06	25.01±0.01	1.37±0.01	0.80±0.01	0.47±0.01	32.61±0.05
Layer	1.60±0.07	1.67±0.01	3.38±0.01	10.96±0.01	18.88±0.06	1.37±0.01	0.53±0.06	1.42±0.01	38.04±0.06
Finisher	2.13±0.02	2.51±0.12	2.70±0.01	10.96±0.01	18.88±0.06	2.06±0.01	0.27±0.06	1.42±0.01	21.74±0.01

DISCUSSION

In poultry feed reference standard prepared by the Standard Organization of Nigeria (SON), there are requirements of some heavy metals mentioned as micro-nutrient but no standard as a contaminant in terms of maximum acceptable limit for the heavy metals was mentioned. Copper, zinc, manganese and iron which are heavy metals are included in the requirement as micro-nutrient. Zinc was mentioned as a nutrient at 40- 55mg/kg in starter, grower and finisher and 30-40mg/kg in layer feed. With the exception of feed A (Table 1) layer sample, all the other samples contained zinc below the micro-nutrient requirement. However, these were comparably lower than 54.3-482.2mg/kg obtained by Mahesaret *al.* (2010) and within the range of 33.945-49.950mg/kg obtained by Okoyeet *al.* (2011) in their analysis of poultry feed.

Cadmium levels (Table 1 – 2) was found in all samples to exceed the permissible limits of FAO/WHO which both are 1mg/kg.

Lead concentration level in all the samples was below the permissible limit of 1mg/kg in the United Kingdom (Nicholson, 1999), however it is also lower than the maximum acceptance limit of 5mg/kg by FAO/WHO. The values obtained in this study were lower than 1.10-7.85mg/kg and 23.2- 32.6mg/kg obtained by Okeoyeet *al.* (2011) and Mahesaret *al.* (2010) in their analysis of poultry feed respectively. Chromium was also detected in all the feed samples and was found to be above the maximum acceptable limit of 0.3mg/kg (Act No. 21 NRC, 2006) where Nickel was detected but found to be below the maximum acceptable limit of 4.05mg/kg (Act No. 21, NRC 2006). However, comparing the values obtained with that of Okoyeet *al.* (2011) of 2.250-4.875mg/kg, the values were found to be lower. Iron was found in all samples but below the permissible level of 45-80mg/kg as stipulated by FAO and comparing with that of SON (90-95mg/kg) for starter; grower and finisher while 50- 60mg/kg for layer, it was also observed that all the values obtained in this study are below requirement. Feed A grower and starter contained lowest content of Zinc (08.787mg/kg) which will not suffice the nutritional requirement of the poultry. In the case of Cobalt, comparing the values obtained with the maximum acceptable limit of 1mg/kg as stipulated by FAO/WHO (2010), only five (5) of the feed samples are below the limit. Manganese and copper being part of the essential trace minerals are also detected in the entire feed samples. Comparing the values obtained for manganese with that of the maximum acceptance limit of 20-60mg/kg as stipulated by FAO, it was found that all the feeds are below with the exception of eight (8) of them, which are within the limit of 20-60mg/kg while

Manganese which is also a micro-nutrient mentioned by SON (2012) at 55-60mg/kg for starter and finisher, 30-40mg/kg for grower and 50- 60mg/kg for layer. It was found that, all the samples are below requirement. Copper was referred to as micro-nutrient at (0.0-10mg/kg) level in starter and (9-10mg/kg) in grower, layer and finisher. In the samples, all the feeds for starter are within the ranged but for the grower; layer and finisher are found to be below the ranged. Statistically, significant difference is observed in all the metals ($p < 0.05$).

The study revealed the presence of Pb and Cd in chicken feeds at non toxic levels. To maintain the safety of food chain and to further minimize the heavy metals contamination therefore, it is mandatory for the nation's chicken feed producers to always observe and maintain these standards for heavy metals in chicken feeds, then any activity allowing heavy metals to gain access into food chain beyond those limits can automatically be regarded as unsafe and will be prescribed for alleviation strategies. On the other hand, zinc, copper and manganese appeared in all the feed samples at a level below what is required by chickens. Therefore, efforts should be made to increase supplementation of these micro-elements in chicken feeds.

Comment [MF16]: Re-write to reduce the plagiarism

CONCLUSION

The essential elements zinc, iron, manganese and copper were found to be low in the feed and the nutritive values of the feeds are estimated from the concentration level of the essential elements. Some of the heavy metals are found to be above the maximum acceptable limit whereas majority fall below the limit. The metals in the whole feeds were found to be statistically significant (p is less than 0.05)

Therefore, a definite standard for heavy metals should be provided as a contaminant so as to maintain the food chain safe from heavy metals and subsequent consequences. Extra care need to be carried out by manufacturers in order eliminate/reduce heavy metal content in the feed and there is also need for the manufacturers to increase the quantity of supplements been added so as to increase its nutritional value.

The starter feed type consistently had the lowest concentration of the essential elements. The essential elements (zinc, iron, manganese and copper) were very low in the feed. Therefore, the nutritive values of the feed as estimated from the concentrations of the essential elements were very low. This shows that supplements were not added to the feeds as should have been expected. However there were very high concentrations of lead in the feed samples. This could be attributed to anthropogenic sources of lead pollution in the environment especially fossils fuels. There is a great need to adopt the alternative renewable energy sources such as biodiesel and bioethanol.

Further studies should be done to cover more locations/states in western part of the country. Also, other poultry feeds available in the region should be screened for heavy metal contamination. Feed companies should periodically carry out heavy metal assessment of their feed products so as to keep them at a safe level.

Comment [MF17]: Re-write too

UNDER PEER REVIEW

REFERENCES

Abulude FO, Akinjaagunla YS & Omoniyi A (2006a). An investigation into the effect of vehicular exhaust fumes on the levels of some heavy metals in cows' blood. *Research Journal of Biological Sciences*, 1(1-4): 9-11.

Abulude FO, Eluyode SO & Jegede A (2006b). An investigation into the effect of traffic pollution on the levels of some heavy metals in goats' urine samples. *Journal of Animal and Veterinary Advances*, 5(2): 132- 134

Ahmad, F., Muhammad, T.J., Mansor, A.S. and Kausar, R., (2004). Effects of higher levels of chromium and copper on broiler health and performance during tropical summer season. *Vet. Arch* 74(5): 395-408.

Aschner M (2002). Neurotoxic mechanism of fishborne methylmercury. *Environmental Toxicology and Pharmacology*, 12(2): 101- 102.

Aycicek M, Kaplan O & Yaman M (2008). Effect of cadmium on germination, seedling growth and metal contents of sunflower (*Helianthus annuus* L.). *Asian Journal of Chemistry*, 20(4): 2663-2672.

Baath, E., 1989. Effects of heavy metals in soil on A.M. Gallis, 2009. Determination of heavy metals microbial processes and populations. *Water Air content of meat and meat by-products by using Soil Pollut.*, 47(3-4): 335-379.

Cang, L., Wang, Y.J., Zhou, D.M. and Dong, H.Y., (2004). Heavy metal pollution in poultry and livestock feeds and manures under intensive farming in Hiongsu province China. *Journal of Environmental Science*. 16:371-374.

Coleman ME, Elder RS & Basu, P (1992). Trace metals in edible tissues of livestock and poultry. *Journal of the Association of Official Analytical Chemists International*, 75(4): 615-625.

Costa (2000). Trace elements; Aluminum, Cadmium, Arsenic and Nickel. In: *Environmental Metal Toxicants, Human Exposure and their Health effect*, 2^{le}. John Wiley and sons, Inc. New York City. Pp 811-850.

Delgado Anoyo, M., Mrialles de Imperial, H.R., Peralla, F.A. and Almestre, R.C., (2014). Heavy metal concentration in soil, plant, earth worm and leachate from poultry manure applied to agricultural land in Spain. *Rev int. contam. Ambio*. 30(1): 45 – 50.

Demirezen, O. and Uruc, K., (2006). Comparative study of trace elements in certain fish, meat and meat products. *Food chemistry*, 32: 215-222.

Dudka, S. and Miller, W.P., (1999). Accumulation of potentially toxic element in plants and their transfer to human food chain. *Journal of Environmental Science Health*. B34(4): 681-708.

European Commission; Health and Consumer Protection Directorate - General, Belgium (2003). Opinion of the Scientific Committee on Animal Nutrition on Undesirable Substances in Feed. http://ec.europa.eu/food/fs/sc/scan/out126_bis_en.pdf, retrieved 10-04-2015.

Falandysz J, Szymczyk-Kobrzynska K, Brzostowski A, Zalewski K &Zasadowski A (2005). Concentrations of heavy metals in the tissue of red deer (*Cervuselaphus*) from the region of Warmia and Mazury, Poland. *Food Additives and Contaminants*, 22(2): 141-149.

Gerberding J.L., (2005). Toxicological profile for zinc. Atlanta Georgia. 2005, pp 22 and 23.

Gerberding J.L., (2004). Toxicological profile for zinc. Atlanta Georgia. 2004, pp22 and 26.

Hassan, A.R., Saleh, M., Sobih, M., Wilson, S. and Reddy, P., (1998). Effects of some heavy metals pollutants on the performance and immune system of chicks. *Poultry Science* 77(1):24-30.

Iwegbue, C.M.A., G.E. Nwajei and E. Iyoha, 2008. pp: 165. Heavy metal residues of chicken meat and gizzard 29. Davis, K., 1998. Federal Codes of Regulations: and turkey meat consumed in southern Nigeria. *Bulg. Animals and Animal Products. Consumer Report., J. Vet. Med.*, 11(4): 275-280

J.H. Koplan, Atlanta Georgia, 2000a, 26p.

Kaplan, O., Yildirim, N.C. and Cumen, M., (2010). Toxic elements in animal products and environmental health. *Asian J. Animal Vet. Adv.* 18: 1623 – 1627.

Kingery, W.L. Wood, C.W., Williams, J.C and Mullins, G.L. (1994). Impact on environmentally related soil properties. *J. Environ. Qual.* 23: 139 – 147.

Mariam, I.S. and Nagre, S., (2004). Distribution of some trace and macro minerals in beef, mutton and poultry. *International Journal of Agric and Biology*, 6:816-820.

Mariam, I., S. Iqbal and S.A. Nagra, 2004. Distribution of some trace and macro minerals in beef, mutton and 26. Anonymous, 1980. Mineral Tolerance of Domestic poultry. IJAB., 6: 816-820

Maqbool, A., M.N. Sarwar and K. Bakhsh, 2005. S.C. Kestin, 2000. The behaviour of broiler chickens Moderanization of poultry farming. Economic and its modification by lameness. Appl. Anim. Business Review. Pakistan Vet. J., 27(1): 25-28. Behav. Sci., 67: 111-25.

Musa, B., and Abudullahi, M.S., (2013). The toxicological effects of cadmium and some other heavy metals in plants and human. Journal of Environmental Science and water Res. 2(8): 245-249.

National Research Council (1994). Nutrient requirements of poultry. 9th Rev. edn, National Academy Press, Washington D.C, pp. 84-85.

Okoye, C.O.B, Ibeto, C.N and Ihedioha, J.N (2011). Assessment of heavy metals in chicken feeds sold in south eastern, Nigeria. Advance in Applied science Research, 2(3):63-68.

Okoye, P.A.C., Nwude, D.O. and Babayemi, J.C., (2010). Heavy metal levels in animal tissue. A case study of Nigerian raised cattle. Research Journal of Applied Science 5(2): 146-150.

Onwuachu, V.J., (2012). Assessment of heavy metal pollution in local and exotic breeds of chicken raised in Port Harcourt Metropolis and its environs. Ph.D Thesis submitted to Pure and Industrial Chemistry Dept., University of Port Harcourt, Nigeria. p118.

Puls R (1988). Mineral levels in animal health: Diagnostic Data, (Second edition.). Sherpa International, Clearbrook, British Colombia, Canada. Pp. 356.

Raloff, Janet (2003). Food for thought: Global Food Trends. Science new online. May 31,2003.

Sanotra, G.S., C. Berg and J.D. Lund, 2003. A of Heavy Metals in Poultry Feed Production in comparison between leg problems in Danish and Bangladesh J. Sci. Ind. Res., 42(4): 465-474. Swedish Broiler Production. Anim. Welfare

Santhi, D., V. Balakrishnan, A. Kalaikannan and K.T. Radhakrishnan, (2008). Presence of heavy metals in pork products in Chennai (India). Am J. food Technol., 3(3):192-199.

SCAN (2003), Scientific Committee on animal Nutrition: Opinion of the undesirable substance in feed. Retrieved from: http://Europe.eu.int/comm./food/fs/sc/scan/out_126_bis_en.pdf.

Scheinberg, J.H (1991). Copper in Merian E. (ed). Metals and their compounds in the environment. New York p. 893-908

Shah, T.G., J.A.K. Baig, H.I. Afridi, G.A. Kandhro, Council, (NRC), National Academy Press. M.B. Arain, N.F. Kolachi and S.K. Wadhwa, 2010. 27. Anonymous, 1992. World Health Organization Determination of total mercury in chicken feed, its (WHO), Environmental Health Criteria, Geneva, translocation to different tissues of chicken and their pp: 134. manure using cold vapor atomic absorption 28. Anonymous, 1995. World Health Organization spectrometer. FCT., 48(8-9): 2357-2360.

Smeda, 2002. Small and Medium Enterprise 23. Wijewardena, J.D.H. and S.P. Gunaratne, 2004. Heavy Development Authority Government of metal contents in commonly used animal manure. J. Pakistan Pre-Feasibility Study Broiler farm. PREF., Soil Sci. Soc. Sri Lanka., 6: 245-253. 1: 3-10

Zang, M.F., (1995). A summary about pollution and prevention from animal husbandry production around the world. J. Ecology of Domestic animal, 16(3): 45 – 49.