

Case study

Neurologic Complications After Repeat Epidural Blood Patch in Recurrent Spontaneous Intracranial Hypotension : A Case Report

Abstract

A 61-year-old male patient presented to Guro Korea University Medical Center with a one-month history of increasingly severe headaches that were aggravated by an upright position and alleviated by a supine position. The patient had no other specific underlying disease. Epidural blood patch was repeatedly performed several times because headache was not fully subsided, 10 minutes after fourth EBP, the patient showed a decrease in motor grade 1 in both lower limbs, Lt. Upper limbs and Grade 3 for Rt. upper limbs. Urgent surgical evacuation of spinal epidural hematoma was conducted. After operation, patient recovered from neurologic deficit.

While epidural blood patches are generally considered safe and effective for treating SIH, like any medical procedure, there can be potential complications. It's important to note that complications are relatively rare, and the benefits of the procedure often outweigh the risks. When we repeatedly perform epidural blood patches, more caution is required. Careful evaluation of absolute indications is necessary to prevent additional complications and possible complications should be seriously considered.

Keywords

Epidural blood patch, quadriplegia, neurologic deficit, spontaneous intracranial hypotension, CSF leakage

Introduction

Epidural blood patch (EBP) is an effective treatment for spontaneous cerebrospinal fluid (CSF) leak syndrome patients which is characterized by occipital or frontal headaches that improve when lying down and worsen when in an upright position [1]. Usually, about 20 mL of autologous blood that is drawn and injected into the epidural space forms a clot that can

seal the leak and help restore normal CSF pressure[2, 3].The normal range for CSF is reported differently in various sources, with most reporting a normal range of 7-18 cmH₂O in adults[4]. While an EBP (Epidural Blood Patch) is considered effective treatment for CSF leakage, unexpected complications may occur. Common complications may include headaches, neck pain, and back pain. Although rare, serious complications such as chronic adhesive arachnoiditis, subdural or spinal hematoma, seizures, infections, and intracerebral hemorrhage can also occur[5]. In most cases, these complications can be managed through conservative treatments, and in the instance of severe complications, prompt intervention is often instrumental in preventing the development of significant issues.

Case report

A 61-year-old male patient presented to Guro Korea University Medical Center with a one-month history of worsening severe headaches that were aggravated by an upright position and alleviated by a supine position. The patient had no other specific underlying disease and recent trauma history.

Diagnostic workup included brain Magnetic Resonance Imaging (MRI). Brain MRI revealed diffuse subdural hematoma (SDH) and dural enhancements along the both cerebral convexity, falx and tentorium with suspicious dilatation of venous sinuses and sagging of the midbrain. A Computed Tomography (CT) myelography was performed to differentiate CSF leakage, revealing an extensive CSF leakage extending from C2/3 to T7/8. Based on diagnostic imaging, an EBP was performed at the T1/2 level. The patient's headache gradually improved. The patient was prescribed naproxen and discharged.

Nine days after the first procedure, the patient visited the outpatient clinic and expressed a recurrence of headache. The headache was sharp, stabbing with a Numeric Rating Scale (NRS) of 4 and exacerbated by movement. The follow-up CT showed increased in extent of diffuse chronic stage subdural hematoma (SDH) in both cerebral convexities. Consequently, we conducted a second EBP at the T1/2 level, which was at the same location as the first EBP, two weeks after the first EBP. Following the second EBP, a marked alleviation of the headache was noted, and the patient was discharged.

One month after the second procedure, a follow-up brain CT exhibited increased densities of SDH at the bilateral cerebral convexities. As a result, a third EBP performed at the C3-4 level.

If CSF leakage was not seen on CT myelography after the third procedure, burr hole operation was planned for SDH. However, after the third EBP, although the headache was improved, the findings on CT myelography still showed contrast leakage at ventral aspect of epidural space, C2/3 and C6/7 to mid-thoracic level (Figure 1). Therefore, we decided to postpone the Burr hole operation and proceed with a fourth EBP.

3minutes after fourth EBP (Figure 2), the patient showed a decrease in motor grade 1 in both lower limbs, 5 minutes later, decrease in motor grade 1 in Lt. upper limbs and grade 3 in Rt. Upper limb, and 7 minutes later, his speech began to slur. 10 minutes after EBP, the motor grade was grade 1 for both lower limbs and Lt. upper limbs and Grade 3 for Rt upper limbs. The patient's vital sign remained stable and the patient was alert. His pupils were isocoric, measuring 3 millimeters in diameter on both sides (3P/3P).

Spine MRI was taken immediately, and we found epidural hematoma with air bubbles from C2 to T4 posterior epidural space, compressing the spinal cord (Figure 3). Dexamethasone 10mg was administered to patient 1 hour after EBP for neuroprotection. Just before entering the operating room, the patient's motor function recovered to Grade 4, subjectively indicating an approximate 80% recovery. However, considering the lack of complete recovery of neurological deficit and significant cord compression, it was decided to proceed with the surgery in consultation with the neurosurgical team. An emergency operation was undertaken 2 hours after paralysis began. Urgent surgical evacuation of spinal epidural hematoma was conducted. Motor Evoked Potentials (MEP) and Somatosensory Evoked Potentials (SSEP) was monitored during the surgery. Hematoma was originated from externally injected autologous blood component, not hemorrhage from dura. Epidural hematoma was identified from C6 to T2. Hematoma was successfully evacuated without causing damage to the dura mater. Distal MEP was recovered after hemilaminectomy (C6-T2, Right) and hemipartial laminectomy (C5 Right, T3 Right). Proximal MEP showed signs of recovery after the evacuation of the epidural hematoma. CSF leakage from the cervical dura was suspected, but the exact leakage site could not be located. After surgery, the patient's motor function showed full recovery as a Grade 5 in all four limbs and sensory function was also fully recovered. Two weeks after the surgery, the patient's headache completely improved, and a follow-up brain CT showed that the SDH was slightly resolved. The patient was discharged.

Discussion

Spontaneous Intracranial Hypotension (SIH), also known as a spontaneous CSF leak syndrome, is a medical condition characterized by a low CSF pressure within the skull. SIH usually occurs when there is a leakage of CSF from the spinal canal, reducing the pressure of CSF around the brain. This condition often causes a variety of symptoms that can vary in severity [6].

Conservative management may include bed rest, increased fluid intake, and pain medications. In more severe cases, targeted interventions such as epidural blood patches, fibrin glue injections, or surgical repairs may be considered [7, 8].

Some SIH patients might have an underlying connective tissue disorder or other factors that predispose them to developing multiple leaks. One of the initial treatment modalities, epidural blood patch (EBP), does not always provide a permanent solution, and may require repeated intervention [9]. Repeated EBPs may be more frequently required in patients with SIH [10].

Moreover, in some cases, as described in this report, complications may arise, including inadequate spread of blood components within the epidural space due to repeated EBPs. The autologous blood component, which did not spread and accumulated, compressed the spinal cord as mass effect, and changes in neurologic abnormality appeared rapidly.

Prolonged INR and CSF leakage can be associated with repeated EBPs in patient with SIH [10]. However, in this case, no abnormal bleeding tendency were found in the blood coagulation test, it is more likely that autologous transplanted blood which was loculated and caused myelopathy as a mass effect.

The inadequate spread of blood components within the epidural space may arise due to multiple reasons. Patient-specific anatomical variations, such as spinal curvature and epidural adhesions, could impede the uniform distribution of the injected blood. Furthermore, the rate of injection might influence the ability of the blood to diffuse and cover the necessary area for effective sealing. It is also possible that repeated EBP procedures may have resulted in fibrosis or scarring within the epidural space, rendering it less amenable to blood spread.

In the case report by Iga K et al. (2019) [11], published in JA Clinical Reports, the authors present a noteworthy case involving a patient who underwent multiple EBP procedures due to

unintentional dural puncture during combined spinal-epidural anesthesia for a scheduled cesarean section. After the repeated EBPs at the site of dural puncture, she reported neurological complications, such as radicular pains. The procedure led to neurological deficits due to issues related to blood component distribution and subsequent spinal cord compression. This case emphasizes the necessity for precise procedural considerations and vigilance when contemplating repeated EBPs, especially in the context of unintentional dural puncture. While our case and Iga K case both involve complications related to repeated EBP procedures, the underlying reason for EBPs was different. And in the Iga K case, conservative treatment like oral analgesic therapies performed, and the neurologic symptoms were alleviated compared to the first time, so the patient declined further radiologic studies at the time and was discharged, which is different from our case where immediate emergency treatment was performed through surgical intervention.

The other case report by Carlswärd C presented the occurrence of chronic adhesive arachnoiditis following the administration of repeat EBPs [12]. A healthy, 29-year-old woman underwent two epidural blood patches due to severe PDPH occurring after epidural analgesia for delivery. And she experienced lumbar pain and radiculopathy in both legs as well as pain radiating to the upper thoracic region. MRI showed changes with arachnoiditis and an intradural haematoma extending from T12 to S2. Conservative analgesic therapy with anti-inflammatory drugs, corticosteroids, and physiotherapy improved her condition slightly but during the following weeks her symptoms worsened without significant relief. In the end, Follow-up MRI two years after the incident did not show further improvement and her physical condition remains unchanged.

The Carlswärd C case is an unfortunate case in which the patient's condition became chronic and the symptoms could not be relieved, affecting the patient's life. Compared to our case, Both cases, Iga K and Carlswärd C, have something in common: complications that occurred after repeated epidural blood patches were performed, but the difference is that the symptoms patients were complaining about were not fully recovered. In addition, there is a difference in that PDPH occurred after spinal or epidural anesthesia was the reason for performing a blood patch in both cases. Two epidural blood patches were performed in the above two cases, but in our case, a total of four times was performed, which may have resulted in more severe adhesion.

In our case, quadriplegia occurred due to compression of the spinal cord by the autologous

blood components after 4 times epidural blood patches. And the patient was recovered completely without sequelae after decompressive laminectomy. This is because, through prompt detection and response, surgery was decided and carried out immediately after rapid multidisciplinary consultation. Since this has not been reported before, we believe it is a good case that emphasizes the importance of close monitoring and rapid detection of patients after epidural blood patch, and the need for a multidisciplinary approach.

To minimize the risk of complications, it is necessary to meticulously assess patient's anatomy through imaging studies, such as MRI or CT before proceeding with EBP procedures. Choosing an appropriate injection site and technique can help optimize the spread of blood components. Additionally, consideration should be given to the volume and rate of injection, ensuring a controlled and uniform distribution of blood. In cases where repeated EBPs are being contemplated, careful monitoring of the patient's response and potential complications is essential.

The case report's successful outcome can be attributed, in large part, to the vigilant neurological monitoring implemented post-EBP procedures. Continuous monitoring of the patient's neurological status allowed for the early detection of any emerging deficits, enabling rapid intervention before irreversible damage occurred. The ability to identify subtle neurological changes underscores the necessity for a multidisciplinary approach involving anesthesiologists, neurologists, and neurosurgeons to ensure comprehensive patient care. The decision to proceed with emergency surgery reflects the agility and expertise of the medical team involved. In cases where neurological deficits develop following interventions such as EBPs, prompt decision-making is crucial to mitigate potential complications. The surgical approach chosen likely depended on the patient's presentation, available imaging data, and the expertise of the surgical team. The case report should ideally elaborate on the factors considered when opting for surgical intervention.

Conclusion

This case report highlights a rare but serious complication of EBP procedures in the management of recurrent spontaneous intracranial hypotension. An understanding of the underlying anatomy, pathophysiology, and factors influencing EBP success is crucial for preventing adverse outcomes. Close cooperation between anesthesiologists, neurologists,

neurosurgeons, and neuroradiologists is essential to develop optimal strategies for patient selection, procedure execution, and post-procedural monitoring, with the goal of minimizing the risk of spinal cord compression and associated neurologic deficits. Especially after the procedure, it is important to observe the patient's condition through careful monitoring to quickly recognize changes and prepare to respond.

INFORMED CONSENT

Written consent obtained from the patient has been collected and preserved by the authors.

ETHICAL APPROVAL

This article does not contain any studies with human participants or animals performed by any of the authors.

Reference

- [1] Phillip B. Gaukroger, Peter Brownridge, Epidural blood patch in the treatment of spontaneous low CSF pressure headache, PAIN®, Volume 29, Issue 1, 1987, Pages 119-122, ISSN 0304-3959, [https://doi.org/10.1016/0304-3959\(87\)90185-0](https://doi.org/10.1016/0304-3959(87)90185-0).
- [2] Shin HY. Recent update on epidural blood patch. *Anesth Pain Med (Seoul)*. 2022 Jan;17(1):12-23. doi: 10.17085/apm.21113. Epub 2022 Jan 18. PMID: 35038855; PMCID: PMC8841262.
- [3] Paul J. Zetlaoui, Thomas Buchheit, Dan Benhamou, Epidural blood patch: A narrative review, *Anaesthesia Critical Care & Pain Medicine*, Volume 41, Issue 5, 2022, 101138, ISSN 2352-5568, <https://doi.org/10.1016/j.accpm.2022.101138>.
- [4] Reichman E F, Polglaze K, Euerle B. Neurological and Neurosurgical Procedures: Lumbar Puncture. In: *Emergency Medicine Procedures*. McGraw Hill; 2013:747-761.
- [5] Carlswärd C, Darvish B, Tunelli J, Irestedt L. Chronic adhesive arachnoiditis after repeat epidural blood patch. *Int J Obstet Anesth*. 2015 Aug;24(3):280-3. doi: 10.1016/j.ijoa.2015.04.005. Epub 2015 Apr 29. PMID: 26119259.

- [6] Massey TH, Robertson NP. Spontaneous intracranial hypotension: features, diagnosis and management. *J Neurol*. 2021 Apr;268(4):1555-1557. doi: 10.1007/s00415-021-10500-1. Epub 2021 Mar 8. PMID: 33686508; PMCID: PMC7990821.
- [7] Lin JP, Zhang SD, He FF, Liu MJ, Ma XX. The status of diagnosis and treatment to intracranial hypotension, including SIH. *J Headache Pain*. 2017 Dec;18(1):4. doi: 10.1186/s10194-016-0708-8. Epub 2017 Jan 13. PMID: 28091819; PMCID: PMC5236046.
- [8] Shin HY. Recent update on epidural blood patch. *Anesth Pain Med (Seoul)*. 2022 Jan;17(1):12-23. doi: 10.17085/apm.21113. Epub 2022 Jan 18. PMID: 35038855; PMCID: PMC8841262.
- [9] Hazama A, Awawdeh F, Braley A, Loree J, Swarnkar A, Chin LS, Krishnamurthy S. Recurrent Spontaneous Intracranial Hypotension (SIH) and the Durability of Repeat Epidural Blood Patch (EBP). *Cureus*. 2023 Jul 6;15(7):e41457. doi: 10.7759/cureus.41457. PMID: 37546124; PMCID: PMC10404115.
- [10] Oh AR, Park J, Jeong JS, Lee JY, Choi JW, Kim H, Sim WS. Risk factors associated with repeated epidural blood patches using autologous blood. *Korean J Pain*. 2022 Apr 1;35(2):224-230. doi: 10.3344/kjp.2022.35.2.224. PMID: 35354685; PMCID: PMC8977201.
- [11] Iga K, Murakoshi T, Kato A, Kato K, Terada S, Konno H, Irikoma S, Suzuki T, Matsushita M, Toba Y. Repeat epidural blood patch at the level of unintentional dural puncture and its neurologic complications: a case report. *JA Clin Rep*. 2019 Feb 28;5(1):14. doi: 10.1186/s40981-019-0232-3. PMID: 32025902; PMCID: PMC6966726.
- [12] C. Carlswärd, B. Darvish, J. Tunelli, L. Irestedt, Chronic adhesive arachnoiditis after repeat epidural blood patch, *International Journal of Obstetric Anesthesia*, Volume 24, Issue 3, 2015, Pages 280-283, ISSN 0959-289X, <https://doi.org/10.1016/j.ijoa.2015.04.005>.




Figure 1. An axial computed tomography myelography image at the level of C2/3. This image shows contrast leakage at the epidural space of C2/3. A yellow arrow indicates the epidural space.




Figure 2. C-arm image of contrast injection during epidural blood patch procedure and C-arm image of contrast diluted by autologous blood injection during blood patch.




Figure 3. A parasagittal T2-weighted image and an axial T2-weighted image at the level of C7/T1. The magnetic resonance images show epidural hematoma with air bubbles in C2~T4 posterior epidural space. White arrow indicates the epidural hematoma. These images suggest cord impingement or compression, C2-3~T2-3.