

### Original Research Article

## **Isolation and Identification of Microfungi from Diesel Oil Contaminated Soils and Their Physico-chemical Analysis**

**Abstract-** Diesel oil contaminated soils were collected and analyzed for various Physico-chemical properties that included moisture content, total carbon, total nitrogen and phosphorus using standard protocol and the data compared from normal uncontaminated soils. Analysis revealed that diesel oil contaminated soils possessed higher percent of moisture content, greater amounts of total carbon, nitrogen and phosphorus while these soils were unsuitable for cultivation of agricultural crops. Six species of *Aspergillus* were isolated from diesel contaminated soils using nutrient enrichment cultural technique. These included *Aspergillus luchuensis*, *Aspergillus violaceofuscus*, *Aspergillus niveus*, *Aspergillus terreus*, *Aspergillus japonicus*, *Aspergillus ustus* which grew well on 2% (w/v) agar medium supplemented with 10% diesel oil only which suggest that these species could utilize diesel oil as sole carbon and nitrogen source. The role of carbon, nitrogen and phosphorus in soils has been discussed in this paper.

**Keywords:** Microfungi, Diesel Soil, Soil Analysis, Carbon, Nitrogen, Phosphorus

**Introduction-** Soil, water and air are the essential parts of any ecosystem. Soil harbour approximately 59% of the total species in the biosphere and is the home to 99% of *Enchytraeidae* worms, 90% of fungi, 86% of plants and more than 50% of bacteria but only 3% of mammals live in it [1]. The soil manage the recycling of nutrients, energy, and water to promote the plant growth in the ecosystem. Additionally, the soils play significant role in transfer of carbon between the atmosphere, plant, land, and ocean [2]. Soil microbiology and its impact on ecosystem is required to be studied in detail to understand the functional linkages between different life forms that live in there, as opposed to above ground systems [3]. Soil fungi play significant functional role in the cycling of carbon (C) and other nutrients. They can also exhibit a significant impact on plant growth, either as pathogens that kill plants or as commensal (mycorrhizal associations) that improve nutrient uptake [4]. In addition, prokaryotic functional groups involved in the nitrogen (N) cycling are also directly associated with soil fungi [5]. The most significant chemical components of soil are carbon (C), nitrogen (N) and phosphorus (P), and their relative amounts greatly influence the growth and development of various life forms, nutrient cycling and

energy flow [6]. Plant life depends on mineral elements, which are required for growth, development, and metabolism [7]. Exploring their intrinsic links can help reveal the mechanisms governing the accumulation and transformation of soil C, N, and P fractions because mineral elements and microorganisms have a significant impact on the transformation and transport of these fractions as well [8]. Enzymatic C: N: P stoichiometry has been investigated in detail to understand and evaluate the impact of nutritional limitations of soil microbes [9-11]. Soil microbes secrete extracellular enzymes that break down the complex molecules in the soil, and as a result, the ratios of enzymes reflect relative microbial nutritional constraint [12]. Nutrients can be immobilized and mineralized by soil microbial biomass [13]. Liu *et al.*, [14] opined that soil enzyme activities are connected to "plant-soil enzymes-soil nutrients" and are involved in the biochemical processes of the soil system.

The fungi isolated from petroleum-polluted soil were screened and applied as mono or co-cultures in the treatment of soil-contaminated microcosms in aerobic conditions [15] and evaluated the possibilities of native fungal bioaugmentation as a tactic to encourage the bio remediation of an old, industrially damaged soil that was enhanced with heavy hydrocarbon fractions. Fungal bioaugmentation's bio-degradation efficiency was compared to that of soil bio stimulation (the addition of water and nutrients) and untreated soil as a control. Total petroleum hydrocarbons (TPH) and HMW-PAHs were biodegraded more rapidly by fungal bioaugmentation than by bio stimulation [16]. Oil sludge is produced in vast quantities during the drilling and refinement of petroleum. Utilizing certain microbes for bio remediation is one of the efficient approaches utilized in the waste degradation process [17]. Ten fungal species isolated from polluted soils were assessed for their capacity to break down crude oil samples in solid media based on their rates of growth [18]. According to Sun *et al.* [19], soil's biological and chemical conversion processes directly involve the soil's active organic carbon. Plants and microbes can readily have a considerable impact on the soil's active organic carbon [20]. However, nothing is known about how the amount of soil active organic carbon varies among various vegetation kinds. According to Lino *et al.* [21], soil enzyme activities are connected to "plant-soil enzymes-soil nutrients" and are involved in the biochemical processes of the soil system.

Particularly, the carbon cycle of the soil and the activity of certain enzymes, such as amylases, catalases, ureases, and sucrases, are significant markers of soil fertility [22]. The cornerstone of

modern soils and agricultural management approaches is inorganic chemical-based fertilizers, which are industrially produced compounds that are primarily water-soluble and have high accessible nutrient concentrations [23]. In many terrestrial and aquatic ecosystems, including forests, grasslands, crops, rivers, and wetlands, soil moisture and pH are extensively monitored and assessed [24,25]. According to Neina[26], soil biogeochemical processes like nutrient availability, soil microbial activity, and crop growth and development are all significantly influenced by soil moisture and pH. Given their beneficial impacts on soil C and N nutrient cycles and the growth of the soil biological community underneath the plant species, soil physiochemical characteristics, as well as their spatial pattern and distribution, might generally vary greatly with different plant species [27]. High potential fungal diversity levels in soil can also limit our capacity to draw broad conclusions about how environmental conditions such as global warming will affect various fungal species. To make biological reactions more predictable, phylogenetic methods that assess whether taxa react similarly to environmental change are being used more and more. According to Martiny *et al.*, [28], a finding of phylogenetic conservatism among taxa—close relatives behaving similarly to environmental factors—could improve predictability in response to climate change. Fungal species have great capabilities to degrade diesel oil by producing different amounts and varieties of extracellular secretory enzymes that can break down the complex compounds in nature including hydrocarbons and diesel oil. In fact, the hydrocarbons contaminated soils are not suitable for the production of agricultural crops but if these are re-lamed using microbial strains, they may provide better production of crops.

## **Materials and Methods-**

**Collection of Samples-**The samples were collected from diesel oil-contaminated soils like diesel oil-petrol pumps, workshops having spill of hydrocarbon oils, and parking areas with visible signs of petrol-diesel leakages. Those soils were collected which were exposed to diesel oil spills for a long time. The soil samples were collected 2-3 cm depth with sterile spatula in sterile glass containers, brought to the laboratory immediately and stored at 4°C for further examination.

**Isolation of diesel oil utilizing microfungi:** The enrichment culture techniques was used for the isolation of microfungi for which the soil samples were per-incubated in moist chamber rich with diesel oil at 37°C for 7 days followed by inoculation of 1 mL suspension (1 g soil in 100 mL sterile

distilled water) of per-incubated soil onto Petri plate containing sterile 2% (w/v) agar + 10% diesel oil. The plates were incubated at 37 and 45°C temperature for 3 - 7 days. The fungal colonies showing profuse mycelial growth and/or sporulation were sub-cultured on modified Czapeck's Dox Agar medium (agar 15 g/L; di-potassium hydrogen phosphate 1 g/L, Iron sulfate heptahydrate 0.01 g/L, magnesium sulfate heptahydrate 0.5 g/L, sodium nitrate 3 g /L and the Sucrose was replaced with 10g/L jaggery + 10mL diesel oil), identified using standard reference books [29, 30] and stored at -20°C temperature until further used.

**Physico-chemical Properties of Soil Samples-** The amounts of soil organic matter, readily accessible phosphorus, were measured in accordance with Das *et al.*, [31] explanation.

**Soil Moisture Content-** The moisture contents of the control and diesel contaminated soil were measured by weighing a fresh sample of soil (15g) and dried in the oven maintain at 80 °C overnight till constant weight, cooled down in the desiccators and weighted. Moisture was calculated as per the formula given below.

$$\text{Moisture Content \%} = 100 \times \frac{\text{Initial weight} - \text{Final weight}}{\text{Initial weight}}$$

**pH-** Soil pH was determined by suspending 10g of air-dry soil with 100 mL of deionized in 250 mL Erlenmeyer flask, stirred well, allowed to settle for 10 minutes. pH of the suspension was determined using glass electrode with digital pH meter (SYSTRONICS model  $\mu$  pH System 362).

**Organic Carbon-** For determination of organic carbon, soil sample was air dried, sieved through (0.5 mm sieve size), 1 g sample was taken in 250 mL Erlenmeyer flask, to which 10 mL of 1N solution of potassium di-chromate was added and spread over the soil sample by swirling, 20 ml of concentrated Sulphuric acid was added. The flask was immediately gently stirred until the soil and reagent were combined, then aggressively stirred for one minute. After being spun once again, the flask was left to stand on an asbestos sheet for around 30 minutes. Then 100mL of distilled water was added and allowed to cool once more and the suspension was filtered, 5 mL O- phosphoric acid + three to four drops of 0.5N ferrous sulphate solution were added. The solution initially showed greenish hue as the tip drew closer before turning dark green. At this point, ferrous sulphate was gradually added until maroon color was reflected in light on a white background abruptly changed

from blue to red. However, there was no soil used to standardize the dichromate for the blank determination. The following formula was used to calculate % organic carbon:

$$\% \text{ organic C in soil} = \frac{(\text{K}_2\text{Cr}_2\text{O}_7 - \text{FeSO}_4 \times 0.003 \times 100)(F)}{\% \text{ of air in dry soil}}$$

**Total Nitrogen**-total nitrogen was estimated using Kjeldahl method as described by Anurag and Garg [32] where the soil sample was first digested with concentrated  $\text{H}_2\text{SO}_4$  in presence of catalyst selenium, condensed and then allowed to cool to which boric acid ( $\text{H}_3\text{BO}_3$ ) was added, combined with 40% NaOH. The released  $\text{NH}_3$  was titrated with regular HCl and terminal point's color changed from green to pink .

**Phosphorus**- 1g of air-dried soil was taken into a 15mL centrifuge tube, 7mL of the extraction solution was added, mixed well for 1 minute, and centrifuged at 2,000 rpm for 15 minutes. The supernatant was evaluated Colorimetrically using ascorbic acid molybdate blue as described by Ibrahim et al. [33].

## Results and Discussion –

All 5 samples contaminated with diesel oil exhibited slightly acidic soils when compared with adjoining non-contaminated control soils (Table 1). The acidity of the soils may be due to the degraded products of hydrocarbon contaminated soils. Soil pH has a direct impact on nutrient availability [34, 65]. Schroder et al. [35] found that long-term N fertilizer application can lower soil pH, which may result in a fall in nutrient availability. Long-term inorganic fertilizer application facilitate the nitrification and acidification processes, which results in the drop in soil pH [36,37]. The kind of soil, the source of the manure, manner of its application, and numerous other soil dynamics all affect the soil's pH [38].The presence of carbonate and bicarbonate, as well as carboxyl and phenolic groups, typically results in the rise of soil pH [36]. Soil pH is a key condition with substantial influence on soil biology, chemistry and physical processes which have direct impacts on plant growth and their development. Soil pH and crop productivity are linked together. The United States Department of Agricultural National Resources Conservation Service has categorized soil pH as follows: ultra-acidic (<3.5), extremely acidic (3.5–4.4), very strongly acid (4.5–5.0), strongly acidic (5.1–5.5), moderately acidic (5.6–6.0), slightly acidic (6.1–6.5),

neutral (6.6–7.3), slightly alkaline (7.4–7.8), moderately alkaline (7.9–8.4), strongly alkaline (8.5–9.0) and very strongly alkaline (>9.0) [39,40]. Agricultural crop production is generally conducted within the range of slightly acidic to slightly alkaline which facilitate optimal availability of soil nutrients. In all soils, solubility, mobility and bio availability of trace elements is strongly affected by pH. A full understanding of pH is necessary for optimizing nutrient cycling, soil remediation and plant nutrition, as it affects the entire interacting system.

Mean nitrogen, phosphorus, and carbon concentrations were found higher in hydrocarbon contaminated soil samples when compared with their corresponding non-contaminated normal control soils (Table 1). It shows that hydrocarbon contaminated soils provide additional nitrogen, carbon and phosphorous to the soils, however, the diesel oil contain various other components that are harmful to the biology of the soil which makes the soils unsuitable for cultivation of crops. If the biology of such soils is activated using microbial inoculations, these may be used better fertile soils for cultivation of crops. This paper suggests that infertile hydrocarbon soil can be recovered and soil nutrients may be restored using soil fungi. It has been reported that microfungi like nitrogen-fixers, phosphorus and carbon solubilizers, and arbuscular mycorrhizal fungi affected the growth of agricultural crops [41-43]. The soil that had been colonized with microfungi absorbed more carbon and phosphorus to varied degrees compared to the control. Higher nutrient concentrations were seen in pots contaminated with the concoction of arbuscular fungus, phosphorus and carbon solubilizers, and nitrogen-fixers.

Diesel contaminated soils contained higher amounts of carbon 0.41% (S-1), 20.40% (S-2), 0.38% (S-3), 0.37% (S-4), and 0.39% (S-5), at depths of 0 to 1 cm in comparison to uncontaminated normal control soils that exhibited only 0.3% of total organic carbon. The organic carbon is crucial for the growth and cultivation of crop plants and the increasing amounts of chemical fertilizers have fastly depleted soil organic carbon from 3-4% to 0.3% in Indian soils over the past 70 years and this depletion has been very fast during the last 20 years. At a soil depth of 2 cm, S-1 and S-2 had a greater soil organic carbon(SOC) amount than control soils. The SOC is crucial to soil processes that provide an array of ecosystem services. Abdelhafez et al. [44], found that the organic matter in manure has more stable organic components as it decomposes. SOC levels in normal soils were higher at the soil's top than they were at deeper soil layers in all treatments, indicating a slower translocation of SOC through the soil profile [45]. There is a significant association between soil aggregate stability, soil structure, and SOC content[46]. Present study revealed that diesel soil

treatment resulted in greater soil organic matter (SOM).

The moisture content was also higher in diesel contaminated soils 13.1% (S-1), 13.5 % (S-2), 14.0 % (S-3), 13.9 % (S-4), 12.7 % (S-5) as compared to control soil. The moisture content is important physiological parameter of soil biology as all the minerals and nutrients are absorbed by the plants only in water soluble form. The additional supply of hydrocarbon rich diesel oil may increase the water holding capacity of the soils to retain more amount of water. The water holding capacity of the soil is an important factor of soil ecosystem that determine the rate of photosynthesis, carbon allocation, plant growth, nutrient cycling, microbial activities, soil biodiversity, soil physiology including enzymatic availability for breakdown of different complex compounds in the soil [47].

Table 1: Physico chemical properties of control and diesel contaminated soils

Physico-chemical properties	Control Soil	Diesel Contaminated Soils				
		S-1	S-2	S-3	S-4	S-5
pH	7.2	6.9	6.8	7	6.6	6.9
Organic Carbon	0.3 %	0.41%	0.40%	0.38%	0.37%	0.39%
Total Nitrogen	1.5%	2.9%	2.0%	2.5%	2.8%	2.7%
Phosphorus	0.5%	1.1%	1.0%	0.9%	1.1%	1.3%
Moisture Content	11.8%	13.1 %	13.5%	14.0%	13.9%	12.7%

The total nitrogen (TN) level and total phosphorus (TP) was 2.95% (S-1), 2.0% (S-2), 2.5% (S-3), 2.8% (S-4), 2.7% (S-5) than 1.5% (control soil) showing that total carbon, nitrogen and phosphorus all three were higher in hydrocarbon contaminated diesel oil rich soil. Such soils are dominant in oil rich Arabian countries and our studies reveal that using microbial inoculation of such soils, these


may be converted into fertile soils rich in carbon, nitrogen and phosphorus that may be used for agricultural crop production. Both soil total nitrogen (STN) and soil total phosphorus (STP) responded similarly to rising SOC and soil pH, rising in tandem with SOC and falling in tandem with rising soil pH in both alpine meadows and wetlands. On the Qinghai-Tibetan Plateau, alpine shrub land has shown similar connections between STN and soil pH and SOC [48]. Fang et al. [49], demonstrated that soil organic matter decomposition is a prominent mechanism for soil N buildup, which causes a positive association between STN and SOC. The Three Rivers Source Region's alpine wetland alpine meadows show a positive correlation between STP and SOC because soil P can also be affected by organic matter supply and loss [50]. Both STN and STP demonstrated a notable negative connection with soil pH, in contrast to their favorable relationship with SOC. The facilitation of microbial activity and moderate alkalinity promotes the breakdown of soil organic matter, whereas acidic conditions often slow the turnover of soil organic matter. As a result, raising of soil pH caused a decline in soil organic matter [51].

The mineralization of organic phosphorus is a crucial soil process and only the inorganic phosphorus can be absorbed by the plants which is released into the soil solution only by microbial activity, particularly the microfungi found in soil and on plant roots [42,52]. Numerous soil fungi, mainly belonging to the genus *Aspergillus* have been demonstrated to have the capacity to convert insoluble phosphates into the soluble forms in soils by secreting organic acids [53]. These acids cause the breakdown of bound phosphate forms and lower pH. Several researchers have demonstrated that plants injected with P solubilizing fungi have improved phosphorus uptake and promote the growth [54].


The purpose of the present investigation was to isolate microfungi from diesel contaminated soils using substrate enrichment technique so that we may further evaluate the potential of these soil fungi to degrade the diesel oil in nature for reclamation of such soils using microbial ecosystem. Present study revealed that the species of *Aspergillus* were highly dominant in diesel contaminated soils and we have isolated total *Aspergillus luchuensis*, *Aspergillus violaceofuscus*, *Aspergillus niveus*, *Aspergillus terreus*, *Aspergillus japonicus*, *Aspergillus ustus* (Fig 1) that were actively associated from hydrocarbon contaminated diesel oil soils. These species were isolated using nutrient enrichment technique and were found to grow fast and optimum on 2% agar containing diesel oil as sole carbon and nitrogen source. For identification, colony morphology, mycelial structure, conidiophore formation (size and color), vesicle orientation and morphology of conidia.

The genus *Aspergillus*, family Trichomaceae, and phylum Ascomycota were used to categorize *Aspergillus sp.* A common species that is isolated from diesel soil is *Aspergillus spp.* that have been utilized to produce a variety of significant industrial enzymes, including cellulases, hemicellulases, and proteases, which are employed hydrolysis of lignocarbohydrates in nature and under laboratory conditions. *Aspergillus* is one of the most extensively studied biological control agents for the management of plant-parasitic nematodes [55]. Under *in vitro* conditions, Doilom et al., [56], assessed the phosphate-solubilizing capacity of several isolates of *Aspergillus spp.*, an insoluble tricalcium phosphate, were tested among the fungi. The capacity of *Aspergillus spp.* to solubilize phosphate and nitrogen in soil has been described in scientific literature [29]. It was found that *Aspergillus aculeatus* was able to dissolve rock phosphate. Doilom et al. [56], reported the mineralization of organic phosphates by *A. japonicus* and showed strong acid phosphatase activity in Pikovskaya's broth, indicating that a high percentage of phosphorus was liberated. *A. japonicus* has also been investigated in the past as a rock phosphate solubilizer. By increasing the quantity of soluble phosphorus, *Aspergillus sp.* successfully solubilized a large amount of insoluble rock phosphate [57].

Primary productivity and nitrogen cycling are both constrained by the availability of nitrogen in practically all ecosystems, including those that are agricultural [58]. The rate of photosynthesis can be directly correlated with the availability of nitrogen because a plant requires a relatively significant amount of nitrogen for photosynthesis [59]. Inorganic nitrogenous substances are a kind of nitrogen that plants can absorb. To release an inorganic nitrogen compound that plants may utilize, fixed nitrogen from the atmosphere in terrestrial systems may be bound in organic matter and must first be mineralized [60]. Fungal decomposers in the soil play a significant role in mediating nitrogen conversion [61]. The energy contained in the redox potential of organic nitrogen can be used by a number of soil fungi [62], most fungi have the capacity to release exoenzymes like proteases and peptidases that can degrade organic materials and then can capture molecules that contain nitrogen, creating a direct pathway from organically bound nitrogen in the soil to the plant [55]. Numerous *Aspergillus* species have been found to produce proteases with optimal activity expressed at pH 7. *Aspergillusoryzae* and *A. fumigatus* are two more *Aspergillus* species that have been identified as having the ability to make protease [63,64].



**Figure 1:** Fungal growth in solid CDA media with carbon sources, isolated species were culture in CDA with 1% diesel oil incubated for a growth periods. A) *Aspergillus luchuensis* B) *Aspergillus violaceofuscus* C) *Aspergillus niveus* D) *Aspergillus terreus* E) *Aspergillus japonicus* F) *Aspergillus ustus*.



**Figure 2:** Microscopic characters of species isolated: (A) *Aspergillusluchuensis*(B) *A. violaceofuscus*(C) *A. niveus*(D) *A. terreus*(E) *A. japonicus*(F) *A. ustus*(All under 10 X x 40X ).

**Conclusion-** *Aspergillus luchuensis*, *A. violaceofuscus*, *A. niveus*, *A. terreus*, *A. japonicus*, *A. ustus* were isolated from diesel contaminated soils that utilized diesel oil as sole carbon and nitrogen source when cultured on 2% agar medium containing 10% diesel oil. The soils contaminated with diesel oil possessed higher moisture content, higher percent carbon, nitrogen and phosphorus but were not suitable agricultural crops. The treatment with fungi degraded the oil and enhanced their nutrient quality.

## References-

1. Mark A. A., Bender S.F. and van der Heijden M.G.A. Enumerating soil biodiversity. *Proceedings of National Academy of Sciences* (2023) 120(33) e2304663120 <https://doi.org/10.1073/pnas.2304663120>
2. Babur E. and Dindaroglu T. *Seasonal Changes of Soil Organic Carbon and Microbial Biomass Carbon in Different Forest Ecosystems*. 2020. DOI:<http://dx.doi.org/10.5772/intechopen.90656>.
3. Bach E.M., Ramirez K.S., Fraser T.D., Wall D.H. Soil biodiversity integrates solutions for a sustainable future. *Sustainability*, 2020,12, 2662.<https://doi.org/10.3390/su12072662>
4. Prasad R., Malla R., Bhatnagar K., Das A., Kharakwal H., Verma N., Garg A.P. and Varma A. Beneficial microorganisms: Herbal and Medicinal Plants. In: *microbes for Human Life* (eds. Chauhan, A.K., Harsh, K. and Varma, A.) I.K. International India 2007. vol. 4:49-72.
5. BURKE D.J., CARRINO-KYKER S.R., BURNS J.H.. Is it climate or chemistry? Soil fungal communities respond to soil nutrients in a multi-year high-resolution analysis. *Ecosphere* .2019. 10(10):e02896. 10.1002/ecs2.2896.
6. Bertolet B.L., Corman J.R., Casson N.J., Sebestyen S.D., Kolka R.K., Stanley E.H. Influence of soil temperature and moisture on the dissolved carbon, nitrogen, and phosphorus in organic matter entering lake ecosystems. *Biogeochemistry*.2018.139, 293–305.<https://doi.org/10.1007/s10533-08-0469-3>
7. Hou E., Luo Y., Kuang Y., Chen C., Lu X., Jiang L., Luo X., Wen D.. Global meta-analysis shows pervasive phosphorus limitation of aboveground plant production in terrestrial ecosystems. *Nat. Commun.* 2020, 11, 637.<https://doi.org/10.6084/m9.figshare.8969963>
8. Wu Y. and Yu Y. Soil Carbon, Nitrogen and Phosphorus Fractions and Response to Microorganisms and Mineral Elements in *Zanthoxylum planispinum* ‘Dintanensis’ Plantations at Different Altitudes. *Agronomy* 2023.13, 558. <https://doi.org/10.3390/agronomy13020558>.
9. Feyissa A., Gurmesa G.A., Yang F., Long C., Zhang Q., Cheng X. Soil enzyme activity and stoichiometry in secondary grasslands along a climatic gradient of subtropical China. *Sci. Total Environ.* 2022. 825, 154019 <https://doi.org/10.1016/j.scitotenv.2022.154019>.
10. Xie Y., Ouyang Y., Han S., Se J., Tang S., Yang Y., Ma Q., Wu L. Crop rotation stage has a greater effect than fertilisation on soil microbiome assembly and enzymatic stoichiometry. *Sci. Total Environ.* 2022.815, 152956 <https://doi.org/10.1016/j.scitotenv.2022.152956>.
11. Ye Z., Wang J., Li J., Zhang C., Liu G., Dong Q. Ecoenzymatic stoichiometry reflects the regulation of microbial carbon and nitrogen limitation on soil nitrogen cycling potential in

- arid agriculture ecosystems. *J. Soils Sediments*. 2022. 22, 1228–1241. <https://doi.org/10.1007/s11368-022-03142-x>.
12. Mori T., Rosinger C., Margenot A.J. Enzymatic C:N:P stoichiometry: Questionable assumptions and inconsistencies to infer soil microbial nutrient limitation.2023. <https://doi.org/10.1016/j.geoderma.2022.116242>.
  13. Liddle K., McGonigle T. and Koiter A. Microbe Biomass in Relation to Organic Carbon and Clay in Soil. 2020.doi:10.3390/soilsystems4030041.
  14. Liu S., Zhang W., Wang K., Pan F., Yang S., & Shu S. . Factors controlling accumulation of soil organic carbon along vegetation succession in a typical karst region in Southwest China. *Science of the Total Environment*, 2015. 521, 52–58. <https://doi.org/10.1016/j.scitotenv.2015.03.074>.
  15. Daâssi D., Almaghrabi F.Q. Petroleum-degradingfungalisolates for the treatment of soil microcosms. *Microorganisms*, 2023. 11, 1351. <https://doi.org/10.3390/microorganisms11051351>.
  16. Medaura M.C., Guivernau M., Moreno-Ventas X., Prenafeta-Boldú F.X. and Viñas M. Bio augmentation of Native Fungi, an Efficient Strategy for the Bio remediation of an Aged Industrially Polluted Soil With Heavy Hydrocarbons. *Front. Microbiol.*2021. 12:626436. doi: 10.3389/fmicb.2021.626436.
  17. NoviantyR., Saryono, Awaluddin A., Pratiwi N.W., Hidayah A., Juliantari E. The diversity of fungi consortium isolated from polluted soil for degrading petroleum hydrocarbon. 2021. DOI: 10.13057/biodiv/d221145.
  18. Khdiar M. Y., Mahmood A. A., Shihab L. A., Hardy G.E.ST.J. Isolation of Fungi from Petroleum-Contaminated Soil and Evaluation of Their Ability to Degrade Crude Oil. *IraqiJ.Appl. Phys.* 2023. 19, No. 1, January-March 2023, pp. 13-20
  19. Sun L., Hu T., Kim J. H., Guo F., Song H., Lv X., & Hu H. The effect of fire disturbance on short-term soil respiration in typical forest of Greater Xing'an Range, China. *Journal of Forestry Research*, 2014. **25**(3), 613–620. <https://doi.org/10.1007/s11676-014-0499-1>
  20. Chen H. L., Zhou J. M.& Xiao B. H. Characterization of dissolved organic matter derived from rice straw at different stages of decay. *Journal of Soils and Sediments*, 2010. **10**(5), 915–922. <https://doi.org/10.1007/s11368-010-0210-x>.
  21. Lino I. A. N., Santos V. M., Escobar I. E. C., Silva D. K. A., Araújo A. S. F., & Maia L. C. Soil enzymatic activity in *Eucalyptus grandis* plantations of different ages. *Land Degradation &Development*, 2015.**27**(1), 77–82. <https://doi.org/10.1002/ldr.2454>.
  22. Wang H., Wu J., Li G., Yan L.. Changes in soil carbon fractions and enzyme activities under different vegetation types of the northern Loess Plateau. 2020. DOI: 10.1002/ece3.6852.
  23. Suhag M. Potential of Bio-fertilizers to Replace Chemical Fertilizers. 2016.DOI: 10.17148/IARJSET.2016.3534.
  24. Ghazali M. F., Wikantika K., Harto A. B., & Kondoh A. Generating soil salinity, soil moisture, soil pH from satellite imagery and its analysis. *Information Processing in Agriculture*, 2020.**7**(2), 294–306. <https://doi.org/10.1016/j.inpa.2019.08.003>
  25. Wang G., Zhang X., Yinglan A., Duan L., Xue B., & Liu T. A spatio-temporal cross comparison framework for the accuracies of remotely sensed soil moisture products in a climate-sensitive grassland region. *Journal of Hydrology*,2021. **597**, 126089. <https://doi.org/10.1016/j.jhydrol.2021.126089>.
  26. Neina D. The role of soil pH in plant nutrition and soil remediation. *Applied and Environmental Soil Science*, 2019.5794869. <https://doi.org/10.1155/2019/5794869>.

27. Wang X., Jian S., Gamage L., Li J. Effect of nitrogen fertilization on central tendency and spatial heterogeneity of soil moisture, pH and dissolved organic carbon and nitrogen in two bioenergy croplands.2022. <https://doi.org/10.1002/jpln.202100311>
28. Martiny J. B. H., Jones S.E., Lennon J.T., Martiny A.C. Microbiomes in light of traits: A phylogenetic perspective. *Science* 2015.350:aac9323.doi: 10.1126/science.aac9323.
29. Raper K.B. and Fennel D.I. *The Genus Aspergillus*. The Williams & Wilkins, Baltimore. 1965.
30. Gams W., Christensen M., Onions A.H., Pitt J.I., Samson R.A. Infrageneric taxa of *Aspergillus*. In: *Advances in Penicillium and Aspergillus systematics*. (Samson RA, Pitt JI, eds.) New York: Plenum Press: 1985.55-62.
31. Das S., Liptzin D., Maharjan B. Long-term manure application improves soil health and stabilizes carbon in continuous maize production system. 2023.<https://doi.org/10.1016/j.geoderma.2023.116338>.
32. Anurag and Garg A.P. Estimation of protein of germinated multi-millet puffs using an advanced Kjeldahl titration method. *The Pharma Innovation Journal* 2023; 12(6): 2621-2626 .
33. Ibrahim M., Tasi'u Y.R., Umma M. and Dangora I.I. The effects of inorganic fertilizer on soil characteristics and production of Egg Plant (*Solanum melongena* L.) in Warawa area of Kano State. *Standard Research Journal of Agricultural Sciences* 2019.2(8): 129-135, (ISSN: 2311-2751) <http://www.standresjournals.org/journals/SRJAS>.
34. Murphy C.J., Baggs E.M., Morley N., Wall D.P., Paterson E. Nitrogen availability alters rhizosphere processes mediating soil organic matter mineralisation. *Plant Soil*.2017. <https://doi.org/10.1007/s11104-017-3275-0>
35. Schroder J.L., Zhang H., Girma K., Raun W.R., Penn C.J., Payton M.E. Soil acidification from long-term use of nitrogen fertilizers on winter wheat. *Soil Sci Soc Am J*. 2011.75(3):957–964. doi:10.2136/sssaj2010.0187.
36. Zhu Q., Liu X., Hao T., Zeng M., Shen J., Zhang F., De Vries W. Modeling soil acidification in typical Chinese cropping systems. *Sci Total Environ*. 2018.613:1339–1348. doi:10.1016/j.scitotenv.2017.06.257.
37. Gautam A., Guzman J., Kovacs P. & Kumar S. Manure and inorganic fertilization impacts on soil nutrients, aggregate stability, and organic carbon and nitrogen in different aggregate fractions. *Archives of Agronomy and Soil Science*, 2021.DOI: 10.1080/03650340.2021.1887480.
38. Liang Q., Chen H., Gong Y., Fan M., Yang H., Lal R., Kuzyakov Y. Effects of 15 years of manure and inorganic fertilizers on soil organic carbon fractions in a wheat-maize system in the North China Plain. *NutrCyclAgroecosyst*. 2012. 92 (1):21–33. doi:10.1007/s10705-011-9469-6.
39. Burt R. (*Soil Survey Staff: Soil Survey Field and Laboratory Methods Manual-Soil Survey Investigations Report*, 2014.Vol. 51. Washington, DC: US Department of Agriculture, 227–234.
40. Msimbira L.A. and Smith D.L. The Roles of Plant Growth Promoting Microbes in Enhancing Plant Tolerance to Acidity and Alkalinity Stresses. *Front. Sustain. Food Syst*.2020. 4:<https://doi.org/10.3389/fsufs.2020.00106>
41. Prasad R., Pham H.G., Kumari R., Singh A., Yadav V., Sachdeva M., A.P. Garg A.P., Peskan T., Hehl S., Sherameti I.,Oelmuller R. and Varma A. *Sebacinaceae* : Culturable mycorrhiza-like endosymbiotic fungi and theirinteractionwith non-transformed and

- transformed roots. *In : Soil Biology* Volume : 4, *In vitro Culture of Mycorrhiza*. (eds.Declerch, D., G. Strullu and A Fortin) Springer-Verlag, Berlin Heidelberg, 2005.p. 291-312.
42. Prasad R., Garg A.P., Abdin M.Z., Ahamad S., Bhatnagar K. and Varma A. Friendly symbiotic fungus promoted plant growth and value added herbal medicines. *Intl. J. Symbiosis*. 2006
  43. Kumari R., Prasad R., Pham G.H., Garg A.P. and Varma A. Biotechnology-An overview. *In : Biotechnological Application of Microbes* (eds Podila and A. Varma). IK International-India, New York and Kulwer Academic Press,Holland 2005.2 : 1-41.
  44. Abdelhafez A.A., Abbas M.H., Attia T.M., El Bably W, Mahrous SE. Mineralization of organic carbon and nitrogen in semi-arid soils under organic and inorganic fertilization. *Environ Technol Innov*. 2018.9:243–253. doi:10.1016/j.eti.2017.12.011.
  45. Sithole N.J., Magwaza L.S., Thibaud G.R. Long-term impact of no-till conservation agriculture and N-fertilizer on soil aggregate stability, infiltration and distribution of C in different size fractions. *Soil Tillage Res*. 2019.190:147–156. doi:10.1016/j.still.2019.03.004.
  46. Annabi M., Le Bissonnais Y., Le Villio-Poitrenaud M., Houot S. Improvement of soil aggregate stability by repeated applications of organic amendments to a cultivated silty loam soil. *Agric Ecosyst Environ*. 2011.144(1):382–389. doi:10.1016/j.agee.2011.07.005.
  47. Zhang Y.W., Wang K.B., Wang J., Liu C. and Shangguan Z.P. Changes in soil water holding capacity and water availability following vegetation restoration on the Chinese Loess Plateau. *Scientific Reports (Nature)* 2021. 11:9692 <https://doi.org/10.1038/s41598-021-88914-0>
  48. Nie X., Wang D., Yang L., and Zhou G. Controlling Soil Total Nitrogen Factors across Shrublands in the Three Rivers Source Region of the Tibetan Plateau. *Forest-Biogeosciences For*. 2020.12,559-565.doi:10.3832/for3533-013.
  49. Fang Z., Li D.D., Jiao F., Yao J., Du H.T. The Latitudinal Patterns of Leaf and Soil C:P:N Stoichiometry in the Loess Plateau of China. *Front. Plant Sci*. 2019. 10,1-10. Doi:10.3389/fpls.2019.00085.
  50. Zhou Y., Boutton T.W. and Wu B.X. Soil C:N:P Stoichiometry Responds to Vegetation Change from Grassland to woodland. *Biogeochemistry* 2018.140,341-351.doi:10.1007/s10533-018-0495-1.
  51. Nie X., Wang D., Zhou G., Xiong F., Ren L., Chen Y., Ma K., Zhong Z., Du Y. Drivers of Soil Nitrogen and Phosphorus Storage in Alpine Wetland Across the Three Rivers Source Region on the Qinghai-Tibetan Plateau. 2022. <https://doi.org/10.3389/fenvs.2022.806771>.
  52. Jacoby R., Peukert M., Succurro A., Koprivova A., Kopriva S. The Role of Soil Microorganisms in Plant Mineral Nutrition—Current Knowledge and Future Directions. 2017.doi: 10.3389/fpls.2017.01617.
  53. Tunlid A., Floudas D., De Beeck M.O., Wang T., Persson P. Decomposition of soil organic matter by ectomycorrhizal fungi: Mechanisms and consequences for organic nitrogen uptake and soil carbon stabilization. 2022. <https://doi.org/10.3389/ffgc.2022.934409>.
  54. Barnaola L. and Stout M.J. The effect of mycorrhizal seed treatments on rice growth, yield, and tolerance to insect herbivores. *Journal of Pest Science* 2021. 94:375–392 <https://doi.org/10.1007/s10340-020-01279-7>
  55. Aziz N.H. and Zainol N. Isolation and identification of soil fungi isolates from forest soil for flooded soil recovery. 2018.doi:10.1088/1757-899X/342/1/012028.
  56. Doilom M., Guo J.W., Phookamsak R., Mortier P.E., Karunarathna S. C., Dong W., Liao C.F., Yan K., Pem D., Suwannarach N., Promputtha I., Lumyong S., Xu J.C. Screening of

- Phosphate-Solubilizing Fungi From Air and Soil in Yunnan, China: Four Novel Species in *Aspergillus*, *Gongronella*, *Penicillium* and *Talaromyces*. 2020. <https://doi.org/10.3389/fmicb.2020.585215>.
57. Ceci A., Pinzari, F., Russo, F., Maggi, O., and Persiani, A. M. Saprotrophic soil fungi to improve phosphorus solubilisation and release: in vitro abilities of several species. *Ambio* 2018. 47, 30–40. doi: 10.1007/s13280-017-0972-0.
  58. Chuppa-Tostain G., Hoarau, J., Watson, M., Adelard, L., Shum Cheong Sing, A., Caro, Y., et al. Production of *Aspergillus niger* biomass on sugarcane distillery wastewater: physiological aspects and potential for biodiesel production. *Fungal Fungal. Biol. Biotechnol.* 2018.5:1. doi: 10.1186/s40694-018-0045-6
  59. Wang G., Schultz P., Tipton A., Zhang J., Zhang F. and Bever J. D. Soil microbiome mediates positive plant diversity-productivity relationships in late successional grassland species.2019. doi: 10.1111/ele.13273.
  60. Liu L., Zhu K., Wurzbürger N., Zhang J. Relationships between plant diversity and soil microbial diversity vary across taxonomic groups and spatial scales. 2020.<https://doi.org/10.1002/ecs2.2999>.
  61. Nicolás C., Martin-Bertelsen T., Floudas D. Bentzer J., Smits M., Johansson T., Troein C., Persson P., Tunlid A. The soil organic matter decomposition mechanisms in ectomycorrhizal fungi are tuned for liberating soil organic nitrogen. *ISME J* 2019.13, 977–988. <https://doi.org/10.1038/s41396-018-0331-6>
  62. Sinsabaugh R.L. Phenol oxidase, peroxidase and organic matter dynamics of soil. 2010.<https://doi.org/10.1016/j.soilbio.2009.10.014>.
  63. de Castro R.J.S. and Sato H.H. Protease from *Aspergillus oryzae*: Biochemical Characterization and Application as a Potential Biocatalyst for Production of Protein Hydrolysates with Antioxidant Activities. 2014. <https://doi.org/10.1155/2014/372352>.
  64. Farnelle E., Rousseau K., Thornton D.J., Bowyer P., Herrick S.E. Expression and secretion of *Aspergillus fumigatus* proteases are regulated in response to different protein substrates.2012. doi: 10.1016/j.funbio.2012.07.004.
  65. Murphy S. M. C. Bio stimulation for Enhanced Bio remediation of Crude Oil and Diesel Fuel by Marine Sediment Communities of Canada's Subarctic: A Microcosm-Simulated Oil Spill Study (Master's thesis, University of Calgary, Calgary, Canada). 2020 <https://prism.ucalgary.ca>. <http://hdl.handle.net/1880/111552>