

## Original Research Article

### **Efficacy of early foliar spray intervention on alteration of physiological efficiencies and seed quality attributes of green gram (*Vigna radiata* L.)**

#### **ABSTRACT**

The present study was focused on the evaluation of foliar spray treatment to arrest the flower drop in order to increase the seed set and seed yield in greengram. The crop of greengram var. CO 8 was raised by adopting the recommended package of practices during *rabi* season during 2019 and 2020 at Department of Seed Science and Technology, TNAU, Coimbatore. Foliar application of different nutrients viz., Nutrigold 0.50%, ZnSO<sub>4</sub> 0.10%, Mono ammonium phosphate (MAP) 2%, as treatments were sprayed at bud initiation (30 days after sowing) and 7 days after flowering stages. Pulse wonder @ 5kg ha<sup>-1</sup> was sprayed during peak flowering stage. Unsprayed plots served as control. During the investigations, data on flower production, arresting of flower drop, seed set, seed filling, seed yield and quality of greengram were recorded and studied.

In the study, significant variations were observed due to the foliar applications. Days to 50 per cent flowering, number of flowers per plant, number of pods per plant, conversion efficiency percentage, pod yield ha<sup>-1</sup> and seed yield ha<sup>-1</sup> were significantly higher with foliar spray of Nutrigold 0.50% applied at bud initiation and 7 days later which was on par with Pulse wonder @ 5kg ha<sup>-1</sup> applied during peak flowering stage. The percentage of flower drop was less in Nutrigold 0.50% (39.69%) which was on par with Pulse wonder @ 5kg ha<sup>-1</sup> (41.04%)

---

when compared to control (52.02%). Seed quality parameters such as seed recovery, 100 seed weight, seed germination and vigour of seedlings were significantly higher in foliar spray of Nutrigold 0.50% when compared to other treatments.

From this study, it could be concluded that the foliar application of Nutrigold 0.50% during bud initiation and 7 days after flowering or Pulse Wonder @ 5kg/ ha during peak flowering could be effectively applied to maximize the flower production, arrest the flower drop and to improve the seed yield and quality in greengram.

**Key words** :Greengram, Flower drop, Conversion efficiency, Foliar application, Nutrigold, Pulse wonder

## 1. INTRODUCTION

Greengram is the third important pulse crop in the country after redgram and blackgram grown in nearly 16 per cent of the total pulse area of the country. It is an excellent source of high quality protein (25%), Riboflavin, Thiamine and Vitamin C with high digestibility. It has been reported that greengram has been cultivated in India since ancient times. It is believed that greengram is a native of India and Central Asia and grown in these regions since prehistoric times. It is widely cultivated throughout the Asia, including India, Pakistan, Bangladesh, Srilanka, Thailand, Laos, Cambodia, Vietnam, Indonesia, Malaysia, south China, and Formosa. Greengram is a protein rich staple food. The biological value improves greatly when wheat or rice is combined with greengram because of the complementary relationship of the essential amino acids [1]. As per the World Health Organization, every man needs 50-60 g of pulses per day and Indian Council of Medical Research suggested, consuming 47 g of protein per day to meet out the daily requirement of body. But at present, the per capita availability of pulses is only less than 30 g per day. It is particularly rich in Leucine, Phenylalanine, Lysine, Valine, Isoleucine, etc. In addition to being an important source of human food and animal feed, greengram also plays an important role in sustaining soil fertility by improving soil physical properties and fixing atmospheric nitrogen. It is a drought resistant crop and suitable for dryland farming and predominantly used as an intercrop with other crops.

India contributes more than 70% of world's greengram production. It is grown in about 4.5 million hectares with the total production of 2.5 million tonnes with a productivity of 548 kg/ha

and contributing 10 % to the total pulse production. The states of Rajasthan (13.99 lakh ha), Karnataka (1.14 lakh ha), Madhya Pradesh (0.25 lakh ha), Haryana (0.16 lakh ha) and Gujarat (0.13 lakh ha) are the major producers of greengram in India. According to Government 3<sup>rd</sup> advance estimates, greengram production in 2022-23 is at 3.74 million tonnes. In Tamilnadu, Salem, Nammakkal, Dharmapuri, Vellore, Thoothukudi, Virudhunagar, Tiruvannmalai and Madurai are major districts contribute greengram production (Source: Directorate of Economics and Statistics, 2022-2023).

The yield potential of greengram is very low because of the fact that the crop is mainly grown in rainfed conditions with poor management practices and also due to various physiological, biochemical as well as inherent factors associated with the crop. Apart from the genetic makeup, the physiological factor *viz.*, insufficient partitioning of assimilates, poor pod setting due to the flower abscission and lack of nutrients during critical stages of crop growth, coupled with a number of diseases and pests [2] were the reasons for the poor yield.

Greengram is basically indeterminate type of flowering and fruiting and there is a continuous competition for available assimilates between vegetative and reproductive sinks throughout the growth period [3]. Since, the source is highly limited in pulses with lowering translocation of assimilates to the growing reproductive sinks, leaf area is an important parameter to obtain higher source in terms of higher assimilate production. Major physiological constraints are flower drop and fruit drop. It has been reported that success of pod set in greengram is only 11-32 per cent [4] due to immature pods are lost at later period because of a large proportion of available assimilates is sequestered in old pods. It might be due to that older pod produces hormones like ABA that promote the abortion of younger fruits.

Most legumes are unable to mobilize their assimilates stored prior to fruiting, so storage reserve formation in the seed depends heavily on assimilates formed during fruiting itself, and

seed yield is likely to be extremely sensitive to adverse environmental influences during filling. As with the cereals, the increased protein content of the developing seed is largely at the expense of nitrogen derived from the senescing mother plant (including the pod), although nitrogen fixation by root nodules during reserve deposition may argue against this supply [5].

Nutrients play a pivotal role in increasing the seed yield in pulses [6]. The soil nutrient content may not be always enough to fulfil crop requirement. Most of micronutrients like Fe and Mn are readily fixed in soil having alkaline pH and plant roots are unable to absorb these nutrients adequately from dry top soil. Similarly, Ca and Mg are not easily translocated to leaves within the plant system [7]. Thus, the application of fertilizer containing macro and micronutrients in the soil may not be meeting the crop requirement for root growth and nutrient use. The alternative approach is to apply these micronutrients as foliar sprays.

Foliar application of micronutrients are 6-20 times more efficient than soil application irrespective of soil type [8]. In legumes, nitrogen is more useful because it is the main component of amino acid as well as protein [9]. Phosphorus is the second main critical plant nutrient but for pulses, it assumes primary importance, owing to its important role in root proliferation and thereby atmospheric nitrogen fixation. The role of phosphorus is to increase root development and pod formation. Potassium acts as a co-factor activating number of important enzymes vitally involved in energy transfer in the translocation of carbohydrates, N metabolism and also protein synthesis [10]. Growth regulating substances /growth regulators are known to influence a wide array of physiological parameters like alteration of plant architecture, assimilate partitioning, promotion of photosynthesis, uptake of nutrients (mineral ions), enhancing nitrogen metabolism, promotion of flowering, uniform pod formation, increased mobilization of assimilates to defined sinks, induction of synchrony in flowering, improved seed quality and delayed senescence of leaves in various crops [11].

Hormones play an important role on the vegetative and reproductive growth of any plant. At present, several synthetic hormones are employed by the farmers to boost the growth in the vegetative phase and subsequently to increase the yield. Several efforts have been made to

increase the yield potential through breeding for high yielding varieties incorporating efficient irrigating methods and application of fertilizers, biocides and hormones. Nutrients such as DAP, Urea,  $\text{FeSO}_4$ ,  $\text{ZnSO}_4$  are known to significantly influence the yield of pulses [12].

Plant growth regulators (PGRs) which can improve the physiological efficiencies of plant could play a significant role in increasing crop yield and quality, when applied as foliar spray at proper crop growth stage in optimum concentration [13]. Growth regulators, when applied as foliar spray at proper crop growth stage in optimum concentration could play a significant role in increasing crop yield and quality of produce in different field crops [14]. These plant growth regulators have been considered as software for plant development and improvement in crop productivity [15]. Zn foliar spray at pre-flowering stage minimized the severity of Zn deficiency on the development of reproductive structure and enhanced the seed nutritional status like seed carbohydrate, storage proteins and also increase pod formation and seed yield [16].

Pulse wonder contains macro and micro nutrients that are water soluble and facilitates complete absorption of nutrients. Foliar application at peak flowering stage, reduces flower drops, tolerance drought and improves the yield by 10-20 per cent [17]. Since, the greengram is predominately growing in marginal lands as rainfed crop in Tamil Nadu, the potential of the crop to express its vigour could not be explored. Besides, there is a continuous competition for availability of nutrients from vegetative to pod formation stage and hence there is an urgent need to supply nutrients through foliar means during pod initiation to flowering stage to reduce the flower drop and also to increase the flower production thereby there will be a chance to get another 15-20 per cent increased yield.

## **2. MATERIALS AND METHODS**

Field experiments were conducted at Department of Seed Science and Technology, Tamil Nadu Agricultural University, Coimbatore with greengram *var.* CO8 during *rabi* season 2019 - 2020. The treatments consisted of foliar spray of different chemicals and plant growth regulators

Nutrigold 0.50%, ZnSO<sub>4</sub> 0.10%, Mono ammonium phosphate (MAP) 2%, as treatments were sprayed at bud initiation (30 days after sowing) and 7 days after flowering stages to study the flower production, flower drop arrest, seed set, seed filling, seed yield and quality of greengram. Pulse wonder @ 5kg ha<sup>-1</sup> was sprayed during peak flowering stage. Unsprayed plots served as control.

**2.1 Nutrigold** : Is an organic plant growth promoter has 20% protein and is rich in nutrients and amino acids. Nutrigold improves growth rate and provides nutrients to the plant for quality production and increase the yield by 15 -20 per cent in field crops and vegetables.

**2.2 .Zinc** is an essential element for plants that act as a metal component of various enzymes or as a functional structural or regulatory cofactor and for protein synthesis, photosynthesis, the synthesis of auxin, cell division, and fertilization. Also, zinc plays a special role in synthesizing proteins, RNA and DNA.

**2.3. Pulse Wonder** : Is a booster with nutrients and growth regulators for Pulses. Recommended to decrease flower shedding, Increase yield up to 20 per cent and Increase drought tolerance.

Yield attributes such as days to first flowering, days to 50 per cent flowering, number of flowers per plant number of pods per plant conversion efficiency percentage, flower drop percentage, pod yield per ha, seed yield per ha seed recovery percentage and seed weight (g) were observed and data were recorded.

Then, the seeds were subjected to seed germination tests using paper media (Between Paper- BP) as per the guidelines prescribed by International Seed Testing Association - ISTA [18]. Four replicates of 100 seeds placed in between paper germination paper and kept in germination room maintained with 25±2<sup>0</sup>C temperature and 95±2% relative humidity in the presence of continuous light. After 10 days, the normal seedlings have been counted and the germination has been recorded and it is expressed in percentage. All the data were subjected to statistical analysis and the results were discussed.

### 3.RESULTS

The investigations in greengram var. CO8 revealed that among different foliar applications, plants sprayed with Nutrigold 0.5% took less number of days for first flowering (32.5), it was on par with Pulse Wonder @5 kg /ha (33.6) and ZnSO<sub>4</sub>0.1% ( 34.03). However, the unsprayed control took 35.1 days for first flowering [Table 1.] Nutrigold 0.5% registered the maximum number of flowers (72.35) followed by other foliar application (Pulse Wonder 67.35 and ZnSO<sub>4</sub>0.1% 61.68 and unsprayed control-55.36). Application of Nutrigold 0.5% significantly improved the number of pods and registered 43.63 pods followed by Pulse Wonder @ 5 kg /ha (37.18) and ZnSO<sub>4</sub> 0.1% (31.1) while the control recorded minimum number of pods per plant (26.56) [Table 1.]. The Conversion efficiency of formation of pods from flowers showed significant variation among different foliar application. Among the nutrients, Nutrigold 0.5% registered the highest conversion efficiency of 60.30 per cent which was on par with Pulse wonder @5 kg (58.95%). The lowest conversion efficiency was recorded in control (48.29%). The percentage of flower drop was less in Nutrigold 0.5% (39.69) which was on par with Pulse wonder @5 kg (40.03). The percentage of flower drop was more in unsprayed control (52.02) [Table 1.]

**Table 1. Influence of foliar spray of nutrients on physiological efficiencies and growth attributes in greengram var. CO8**

Treatments	Number of days to first flowering	Number of days to 50 % flowering	Number of flowers plant <sup>-1</sup>	Number of pods plant <sup>-1</sup>	Conversion efficiency (%)	Flower drop (%)
T <sub>1</sub> - Nutrigold 0.5%	32.5	38.5	72.35	43.63	60.30	39.69
T <sub>2</sub> - ZnSO <sub>4</sub> 0.1%	34.2	39.2	61.68	33.05	54.56	41.04

T <sub>3</sub> - MAP 2%	34.6	41.1	60.58	31.10	50.42	45.44
T <sub>4</sub> - Pulse wonder @5 kg	33.6	38.9	67.35	37.18	58.95	40.03
T <sub>5</sub> - Control	35.1	42.5	55.36	26.56	48.29	52.02
<b>Mean</b>	<b>34.0</b>	<b>40.04</b>	<b>63.46</b>	<b>34.30</b>	<b>54.50</b>	<b>43.64</b>
<b>SEd</b>	<b>0.872*</b>	<b>0.635*</b>	<b>1.558**</b>	<b>0.600**</b>	<b>1.417**</b>	<b>1.066**</b>
<b>CD(P = 0.05)</b>	<b>1.90</b>	<b>1.38</b>	<b>3.38</b>	<b>1.31</b>	<b>3.08</b>	<b>2.32</b>

The results of the seed production and quality of seeds showed that among different foliar application, Nutrigold 0.5% recorded highest pod yield (1511.11 kg ha<sup>-1</sup>) followed by Pulse wonder @5 kg (1444.40 kg ha<sup>-1</sup>) which was on par, while the lowest pod yield was recorded in control(1244.60 kg ha<sup>-1</sup>). Similar to pod yield, seed yield also had greater significance due to the application of Nutrigold 0.5% and registered the maximum seed yield per hectare (713.3 kg ha<sup>-1</sup>) which was on par with Pulse Wonder (705.6 kg ha<sup>-1</sup>) while the lowest seed yield was recorded in control (583.0 kg ha<sup>-1</sup>) [Table 2.].

**Table 2. Influence of foliar spray of nutrients on seed quality parameters in greengram var.CO8**

Treatments	Pod yield ha <sup>-1</sup> (kg)	Seed yield ha <sup>-1</sup> (kg)	Seed recovery (%)	100 seed weight (g)	Germination (%)	Vigour Index
T <sub>1</sub> - Nutrigold0.5%	1511.11	713.3	79.26	5.13	82 (64.87)	2696
T <sub>2</sub> - ZnSO <sub>4</sub> 0.1%	1388.90	681.1	74.89	5.03	78 (62.02)	2304
T <sub>3</sub> - MAP 2%	1344.50	648.9	74.53	4.89	76 (60.66)	2284
T <sub>4</sub> - Pulse wonder @5 kg	1444.40	705.6	78.24	5.14	80 (63.43)	2597
T <sub>5</sub> - Control	1244.60	583.0	72.48	4.51	76 (60.66)	2067
Mean	<b>1386.70</b>	<b>666.38</b>	<b>75.88</b>	<b>4.94</b>	<b>78 (62.33)</b>	<b>2390</b>
SEd	<b>43.341*</b>	<b>13.342*</b>	<b>1.819*</b>	<b>0.137*</b>	<b>0.714*</b>	<b>26.51*</b>
CD(P = 0.05)	<b>94.43</b>	<b>29.07</b>	<b>3.96</b>	<b>0.28</b>	<b>1.56</b>	<b>57.77</b>

(Figures in the parentheses indicates arc-sine values)

Highest seed recovery of 79.26% was recorded in Nutrigold 0.5% which was on par with Pulse wonder @5 kg (78.24) while the lowest in control (72.48%). 100 seed weight differed significantly from each other. It was higher in Pulse wonder @5 kg (5.14 g) which was on par with Nutrigold 0.5% (5.13 g). The 100 seed weight was lower in control (4.51g) [Table 1.].

### 3. DISCUSSION

Among the various factors for low productivity in pulses, physiological problems like flower drop and premature shedding of reproductive structure diminished the number of potential sinks or accumulation of assimilates [19] which seems to be associated with nutrient deficiency and hormonal imbalance and ultimately with reduced translocation of drymatter to

reproductive parts. The poor production potential of pulses is attributed to poor photosynthates of pods and seed setting, which may be improved through foliar application of macro and micro nutrients and growth regulators. Consequently, application of nutrient elements through foliar spray at appropriate stages of growth becomes important for their efficient utilization and better performance of the crop.


Foliar sprays are used for three main purposes. They are (i) to maintain optimum nutrition of a particular nutrient, (ii) to give a crop nutritional boost at a critical junctures of different phenophases and (iii) to correct deficiency disorders [20]. There are three ways of absorption of foliar nutrients; they are (i) penetration through the epicuticular wax and the cuticular membrane and (ii) penetration through the cell wall and plasma membrane.

It has been well established that the fertilizer elements which are absorbed through roots can also be absorbed with equal efficiency through foliage. Foliar application has been found to be favourable in short duration crops than medium and long duration ones where the fertilizer applied may not become fully available before maturity of crop. Keeping these points in view, the present study was taken up to develop crop management technique to increase seed set and filling in order to increase the seed yield and quality.

The investigations in greengram var. CO8 revealed that among different foliar applications, plants sprayed with Nutrigold 0.5% took less number of days for first flowering (32.5), it was on par with Pulse Wonder @5 kg /ha (33.6) and ZnSO<sub>4</sub> 0.1% (34.03). Nutrigold 0.5% registered the maximum number of flowers (72.35) followed by other foliar application (Pulse Wonder 67.35 and ZnSO<sub>4</sub> 0.1% 61.68 and unsprayed control - 55.36). The results of the present study are in agreement with the findings of Suganya (2014) in black gram [21]. The predominant effect of foliar application of nutrients during flower initiation and pod formation stage was seen as reduction in flower shedding percentage. This might have significantly

increased the number flowers per plant [19]. Application of Nutrigold 0.5% significantly improved the number of pods and registered 43.63 pods followed by Pulse Wonder @ 5 kg /ha (37.18) and  $ZnSO_4$  0.1% (31.1) .This results are in agreement with the works of Valenciano *et al.* (2007) in common bean [22], Khurana and Chatterjee (2002) in pea[23], Seifinadergholiet *al.* (2011) in common bean [24] and Pandey *et al.* (2013) in black gram[16]. The higher number of pods per plants might be due to the application of balanced nutrition available in Nutrigold at critical phases from flower induction to pod development stages which may prevent the flower and pod drop. In general, foliar application of DAP 2 % twice at flower initiation and pod formation stages of crop growth significantly influenced the number of pods per plant.

The Conversion efficiency of formation of pods from flowers showed significant variation among different foliar application. Among the nutrients, Nutrigold 0.5% registered the highest conversion efficiency of 60.30 per cent which was on par with Pulse wonder @5 kg (58.95%) [Fig.1].These results are in conformity with the results of Kamaraj and Krishnaswamy (2003) in rice [25], Korzeniowska (2008) in wheat [26] and Liew *et al.* (2012) in rice [27]. This might be due to the absorption of abundant nutrients by the plants as suggested by Sarika *et al.* (2006) in linseed [28].



### Treatments


T<sub>1</sub> Nutrigold 0.50% T<sub>2</sub> Pulse wonder @ 5 kg ha<sup>-1</sup> T<sub>3</sub> ZnSO<sub>4</sub> 0.1% T<sub>4</sub> Control

**Fig 1. Influence of foliar spray of nutrients on number of flowers / plant, Conversion efficiency and flower drop (%) in greengram var. CO8**

During the field investigation, the percentage of flower drop was less in the plots sprayed with Nutrigold 0.50 % which was on par with the plots sprayed with Pulse wonder @5 kg which showed a decrease of 23.70 per cent and 21.10 per cent respectively over unsprayed control [Fig.1]. The results are in conformity with the results of Shetty *et al.* (1992) in Bengalgram [29] and Parimala *et al.* (2013) in chickpea [30]. This might be due to the prolonged assimilatory activity of leaves which would have decreased the flower drop and hence increased the number pods per

plant. Supplementary foliar application during flowering was found to increase seed set and prevent premature abortion of embryos, enrich the source sink ratio thereby improves the seed quality.

The results of the seed production and quality of seeds showed that among different foliar application, Nutrigold 0.5% recorded highest pod yield ( $1511.11 \text{ kg ha}^{-1}$ ) followed by Pulse wonder @5 kg ( $1444.40 \text{ kg ha}^{-1}$ ) which was on par, while the lowest pod yield was recorded in control ( $1244.60 \text{ kg ha}^{-1}$ ) [Fig.2]. Similar to pod yield, seed yield also had greater significance



### Treatments

T<sub>1</sub> Nutrigold 0.50% , T<sub>2</sub> Pulse wonder @ 5 kg ha<sup>-1</sup> T<sub>3</sub> ZnSO<sub>4</sub> 0.1% T<sub>4</sub> Control

**Fig 2. Influence of foliar spray of nutrients on pod yield and seed yield plant<sup>-1</sup> in greengram var.CO8**

due to application of Nutrigold 0.5% and registered the maximum seed yield per hectare (713.3 kg ha<sup>-1</sup>) which was on par with Pulse Wonder (705.6 kg ha<sup>-1</sup>)[Fig.2].The reason for the increased yield might be due to the effective translocation of nutrients from source to sink of the crop. Thus, it could have enhanced the yield attributes like number of pods per plant, number of seeds and 100 seed weight and also the higher dry matter production and nutrient uptake of green gram would be the possible reason for increased yield. This positive effect might be due to the better absorption of nutrients applied through foliage leading to better activity of functional root nodules resulting in more leaf area, dry matter production and uptake of nutrients. This could have led to more flower production and subsequently pod formation and other yield attributing characters. This is in conformity with the results of Abbas *et al.* (1994) in blackgram[31], Solaiappan and Ramaiah (1990) in red gram [32]. It is also reported that the absorption and transportation of micronutrients inside the plant system is much easier when foliar application of nutrients are combined with micronutrients [33].

On seed quality evaluation, seed recovery, 100 seed weight, germination and seedling vigour index were maximum in Nutrigold 0.5% which was on par with Pulse wonder @5 kg. This results are in support with the findings of Kathiresan and Duraisamy (2001) in *Sesbani aculeate*[34], Pradeep and Elamathi (2007) in green gram [35]and Vinothkumar *et al.* (2013) in soybean[36].The increased seed weight due to foliar application of Nutriglod might be due to the increased mobilization of metabolites of nutrients to the reproductive sinks resulting into a greater accumulation of food reserves into the seed resulting in enhanced carbohydrate metabolism and ultimately healthy seed.

#### **4. CONCLUSION**

From this study , it could be concluded that the foliar application of Nutrigold 0.50% during bud initiation and 7 days after flowering or Pulse Wonder @ 5kg/ ha during peak

flowering could be effectively applied to maximize the flower production, arrest the flower drop and to improve the seed yield and quality in greengram.

#### **COMPETING INTERESTS DISCLAIMER:**

**Authors have declared that they have no known competing financial interests OR non-financial interests OR personal relationships that could have appeared to influence the work reported in this paper.**

#### **REFERENCE**

1. Hassan FA, Alshamsi ASM, Alyaefi MAS, Kurup S, Al Busaidi N, Ahmed ZFR. Enhancing germination of ghaf seeds (*Prosopis cineraria* L.) using sulphuric acid scarification and cytokinin. In XXXI International Horticultural Congress (IHC2022): International Symposium on Quality Seeds and Transplants for Horticultural spp.2022 ;1365 :39 – 44.
2. Mahala, C.P.S., Dahheech, R.C. and ulhari, R.K. 2001. Effect of plant growth regulators on growth and yield of black gram (*Vigna mungo* L.) at varying levels of phosphorus. *Crop Res.*, 18(1): 163-165
3. Krishna Surendar, K., S. Vincent, Mallika Vanagamudi, and H. Vijayaraghavan. 2013. Plant Growth Regulators and Nitrogen Responses on Improving Nutrient Content of Black Gram (*Vigna mungo* L.). *Plant Gene and Trait*, 4(12): 66-69.
4. Dilip Kumar, M. 2011. In: *Pulses crop production: Principles and Technologies*. (eds) by Asoke, Ghosh, PHI learning Private Limited, New delhi Pp: 177.
5. Bewley, J.D. and Michal Black. 1943. In: *Seeds: Physiology of development and germination*. Plenum press. New York. London. p: 37.
6. Chandrasekhar, C.N. and U. Bangarusamy. 2003. Maximising the yield of mungbean by foliar application of growth regulating chemicals and nutrients. *Madras Agric. J.*, 90(1-3): 142- 145.

7. Foth, H.D. and B.G. Ellis. 1996. Soil fertility. (2nd Ed.) Lewis Pub. New York.
8. Liew, C.S. 1998. Foliar fertilizers from Uniroyal and their potential in Pakistan. Proceedings of seminar on micronutrient in soils and crops in publication, press, Pak p: 277.
9. Nazirhussain, S., M. Mohammed and H.K. Rehana. 2011. Response of nitrogen and phosphorus on growth and yield attributes of black gram. Res. J. Agric. Sci., 2(2): 334-336.
10. Joshi, M. 1997. Agronomic requirements of hybrid cotton. In: Hybrid cotton in India. Kalyani Publishers, New Delhi.
11. Sharma, P., Sadana, V., and K.S. Sukhvinder. 2013. Dry matter partitioning and source-sink relationship as influenced by foliar sprays in groundnut. Int. J. Life. Sci., 8: 1171-1176.
12. Sujatha, K.B. 2001. Effect of foliar spray of chemicals and bioregulators on growth and yield of green gram (*Vigna radiata* L.). M.Sc. (Ag.) Thesis, Tamil Nadu Agricultural University, Coimbatore.
13. Garai, A.K. and J.K. Datta. 2003. Effect of phosphorus sources and cycocel spray on green gram (*Vigna radiata* L.). Legume Res., 26(1): 15 – 19.
14. Nagasubramaniam, A., G. Pathmanabhan and V. Mallika. 2007. Studies on improving production potential of baby corn with foliar spray of plant growth regulators. Ann. Plant Phsiol., 21: 154-157.
15. Malik, C.P. 1995. Plant growth regulators; software for plant development and crop productivity. Presidential address (botany section) Indian science congress association: 1-5.
16. Pandey, N., B. Gupta and C.P. Grish. 2013. Foliar Application of Zn at flowering stage improves plants performance, yield and yield attributes of black gram. Indian J. Exp. Biol., 51: 548-555.

17. Marimuthu, S. and U. Surendran. 2015. Effect of nutrients and plant growth regulators on growth and yield of black gram in sandy loam soils of Cauvery new delta zone, india. Cogent food and Agric., 1: 10.
18. ISTA International rules for seed testing, Switzerland : The International Seed Testing Association:2015.
19. Ganapathy, M., G. Baradhan and N. Ramesh. 2008. Effect of foliar nutrition on reproductive efficiency and grain yield of rice fallow pulses. Legume Res., 31(2): 142-144.
20. Wittwer, S.H., F.G. Teubner. 1959. Foliar absorption of mineral nutrients. Annu. Rev. Plant Physiol., 10: 13-27.
21. Suganya. R. 2014. Effect of foliar nutrient on seed yield and quality of black gram variety vamban 4. M.Sc. (Ag.) Thesis, Tami Nadu Agricultural University, Coimbatore.
22. Valenciano, J.B., M.M. Miguélez Frade, V. Marcelo, B. Reinoso. 2007. Response of irrigated common bean (*Phaseolus vulgaris* L.) yield to foliar zinc application in Spain. N.Z.J. Crop Horti. Sci., 35: 325-330.
23. Khurana, N. and C. Chatterjee. 2002. Effect of zinc on reproductive physiology of pea (*Pisum sativum* L.). Indian J. Agric. Sci., 72: 57-59.
24. Seifinadergholi, M., M. Yarnia and F. RahimzadeKhoei. 2011. Effect of zinc and manganese and their application method on yield and yield components of common bean. Middle East J. Sci. Res., 8(5): 859-865.
25. Kamaraj, A. and V. Krishnaswamy. 2003. Effect of foliar spraying of nutrients on seed yield, split husk occurrence and seed better quality in hybrid rice ADTRH-1. Madras Agric. J., 90 (4-6): 221-223.
26. Korzeniowska, J. 2008. Response of ten winter wheat cultivars to boron foliar application in a temperate climate (South-West Poland). Agron Res., 6:471 - 476.

27. Liew, Y.A., S.R. Omar, M.H.A. Husni, A.M.A. Zainal and N.P.A. Ashikin. 2012. Effects of foliar applied copper and boron on fungal diseases and rice yield on cultivar MR219. *Pertanika J. Trop. Agric. Sci.*, 35: 339–349.
28. Sarika, V., R.M. Deshpande, V.S. Khawale, P.K. Baviskar and B.P. Gurao. 2006. Effect of phosphorus and sulphur application on growth and yield of linseed. *J. Soils and Crops*, 16(1): 217 – 221.
29. Shetty, R.A., A.S. Channabasavanna and S.P. Patil. 1992. Response of gram (*Cicer arietinum* L.) to the foliar application of diammonium phosphate and single super phosphate. *Indian J. Agron.*, 37(4): 828-829.
30. Parimala, K., G. Anitha and A. Vishnuvardhan Reddy. 2013. Effect of nutrient spray on seedling quality parameters of chickpea (*Cicer arietinum* L.) *Plant Archives.*, 13 (2): 735-737.
31. Abbas, Z. 1994. Growth, yield and quality response of mung bean (*Vigna radiata* L.) to N, P and K fertilizer application. M.Sc. (Agri.) Thesis, Univ. of Agric., Faisalabad, Pakistan.
32. Solaiappan, U. and S. Ramiah. 1990. Effect of seed treatments, soil and foliar fertilization on N and P on yield attributes of pigeonpea grown under rainfed condition. *Indian J. Agron.*, 35(3): 234-237.
33. Ibrahim, M.E., M.A. Bekheta, A. El-Moursi and N.A. Gafar, 2010. Improvement of growth and seed yield quality of *Vicia faba* L. plants as affected by application of some bioregulators. *Aust. J. Basic and Appl. Sci.*, 1(4): 657-666.
34. Kathiresan, G. and K. Duraisamy. 2001. Effect of clipping and diammonium phosphate spray on growth and seed yield of daincha (*Sesbania aculeate* L.). *Indian J. Agron.*, 46(3): 568-572.
35. Pradeep, M.D. and S. Elamathi. 2007. Effect of foliar application of DAP, micronutrients and NAA on growth and yield of green gram. *Legume Res.*, 30(4): 305-307.

36. Vinoth Kumar, C., K. Vaiyapuri, M. Amanullah and G. Gopalswamy. 2013. Influence of Foliar Spray of Nutrients on Yield and Economics of Soybean (*Glycine max* L.). J. Biol. Sci., 13: 563-565.

UNDER PEER REVIEW