

Acute toxicity of commonly used insecticides to pumpkin beetle, *Aulacophora foveicollis* Lucas

ABSTRACT

Acute toxicity of commonly used insecticides viz., fipronil, cyantraniliprole, fluxametamide, phenthoate and thiamethoxam were determined against pumpkin beetle, *Aulacophora foveicollis* collected from the unsprayed pumpkin fields in the orchard of Tamil Nadu Agricultural University, Coimbatore using filter paper disc bioassay method. Based on the LC₅₀ values (ppm) of different insecticides, the order of toxicity was found to be fipronil (6.822) > fluxametamide (11.953) > phenthoate (14.732) > cyantraniliprole (14.793) > thiamethoxam (29.465) at 24 and 48 hours after treatment (HAT). The respective LC₉₅ values were 65.396, 87.077, 42.090, 203.017 and 205.013 against *A. foveicollis* at 24 HAT and 46.585, 54.242, 37.445, 223.682 and 146.267 at 48 HAT for fipronil, cyantraniliprole, fluxametamide, phenthoate and thiamethoxam, respectively. The study results clearly indicated that the phenyl pyrazole insecticide, fipronil was highly effective against the pumpkin beetle, *A. foveicollis*.

Keywords: Pumpkin beetle; insecticides; toxicity; median lethal concentrations

1. INTRODUCTION

Cucurbits are commonly grown vegetable for both food and recreational purpose especially used as salad, desert, fruits and vegetable. Cucurbits belonging to family Cucurbitaceae designated as gourd family comprises of 118 genera and 825 species [1] and wide spread throughout the world. They are rich in nutritive and medicinal values. They possess good market value and hence farmers cultivate gourds in a larger scale. Cucurbits include cucumber, musk melon, water melon, squashes, gourds (bitter gourd, ash gourd, ribbed gourd, snake gourd, bottle gourd, etc) and pumpkin which are commonly grown in different parts of the world.

Numerous types of insects pose a threat to cucurbitaceous crops throughout different growth stages causing varying levels of harm to the crop. Among the pest menace in cucurbits, coleopteran insect red pumpkin beetle, *Aulacophora foveicollis* (Coleoptera: Chrysomelidae) is the predominant pest and is highly destructive causing 30-100 per cent yield loss in various crops [2]. The occurrence of this pest is noticed throughout the year. It is a polyphagous pest that inflicts significant harm to cucurbitaceous crops, particularly during their early growth stages.

Aulacophora foveicollis showed higher preference on pumpkin (*Cucurbita maxima* cv. Sitaphal), bottle gourd (*Lagenaria vulgaris* L. Sicerata cv. Lauki), cucumber (cv. Kerala) and musk melon (*Cucumis melo* cv. Kharbooza), respectively [3]. Adult beetles initiate their feeding on cotyledons, flowers and foliage leading to perforations in very young vine plants while grubs dwelling below the surface, cause root damage. In cases of early-sown cucurbitaceous plants, the damage will be very severe and even it leads to resowing of crop [4]. It causes 35-75% damage during seedling stage and 30-100% yield loss in the field conditions [5]. The grubs feed on roots and directly damage the young seedlings.

Farmers commonly rely on chemical pesticides to manage these pest outbreaks. Indiscriminate use of chemical insecticides leads to insecticide resistance, resurgence and residues. Hence, the present investigation was formulated with the intention of screening an effective insecticide for the management of red pumpkin beetle.

2. MATERIALS AND METHODS

The present study was carried out in the Laboratory of Department of Agricultural Entomology, Tamil Nadu Agricultural University, Coimbatore.

2.1 Test Insects

The pumpkin beetles were collected manually from the unsprayed pumpkin field in the orchard of Tamil Nadu Agricultural University, Coimbatore and used for conducting toxicity bioassays. The uniform-sized adult beetles were utilized for bioassays.

2.2 Test Insecticides

The test insecticides used for the study ~~includes~~ include phenyl pyrazole, diamides, isoxazolines, organophosphates and neonicotinoids. The details of the insecticides used in the present study are given in Table 1. The insecticide formulations were procured locally from a pesticide shop and the required concentrations of insecticides were prepared with water. A range of six test concentrations excluding control were prepared for each insecticides and each concentration was replicated four times. The filter paper treated with water alone served as a control.

2.3 Bioassay method

The filter paper disc bioassay method [6] was adopted to assess the acute toxicity of insecticides commonly used in cucurbit crops to pumpkin beetles. The experiment was conducted in plastic containers with perforations to allow adequate aeration for the beetles. Preliminary range finding test was conducted and the obtained concentrations were used for bioassay. The Whatman No.1 filter paper discs were coated evenly with different insecticidal concentrations using micropipette and allowed to dry. These ~~shade-dried~~ shade-dried filter paper discs were placed in glass Petri plates and the pumpkin beetles were released at the rate of 10 per Petri plate and allowed to stay in contact with treated filter paper disc for one hour. After the exposure period, the beetles were transferred into perforated bioassay containers and provided with fresh uninfected leaves as feed. Mortality rates were observed at specific intervals of 24 and 48 h after treatment.

2.4 Statistical analysis

The mortality counts were corrected for control's mortality, if there was any mortality using Abbott's formula [7]

$$\text{Per cent Percent corrected mortality} = \frac{\text{Per cent test mortality} - \text{Per cent control mortality}}{100 - \text{Per cent control mortality}} \times 100$$

The median lethal concentrations (LC₅₀) of insecticide used were determined by Finney's probit analysis [8] and confirmed in EPA probit analysis version 3.1.

Table 1. Details of test insecticides

Insecticide	Trade Name	Class	Manufacturers
Fipronil 80 WG	Jump	Phenyl pyrazole	M/s. Bayer Crop Science Limited
Cyantraniliprole 10.26 OD	Benevia	Diamide	M/s. FMC India Private Limited
Fluxametamide 10 EC	Gracia	Isoxazoline	M/s. Nissan Chemical Corporation
Phenthoate 50 EC	Phendal	Organophosphate	M/s. Coromandel International Limited
Thiamethoxam 25 WG	Actara	Neonicotinoid	M/s. Biostadt India Limited

3. RESULTS AND DISCUSSION

The acute toxicity of commonly used insecticides against pumpkin beetle, *A. foveicollis* at 24 hour after treatment is presented in Table 2. The LC₅₀ values (ppm) obtained were 6.822, 14.793, 11.953, 29.465 and 14.732 for fipronil, cyantraniliprole, fluxametamide, thiamethoxam and phenthoate,

Table 2. Median lethal concentration of insecticides to *Aulachophora indica* at 24 hours after treatment

Insecticides	LC ₅₀ (ppm)	95 % fiducial limits of LC ₅₀		LC ₉₅ (ppm)	95 % fiducial limits of LC ₉₅		Regression equation	X ² value*
		Lower	Upper		Lower	Upper		
Fipronil	6.822	5.081	8.708	65.396	41.469	137.213	Y = 3.603+1.676±0.225x	8.955
Cyantraniliprole	14.793	11.791	17.918	87.077	61.443	151.237	Y = 2.500+2.137±0.274x	4.859
Fluxametamide	11.953	10.121	13.727	42.090	33.044	61.491	Y = 1.758+3.009±0.381x	5.340
Phenthoate	14.732	11.035	19.230	203.017	122.614	434.236	Y = 3.313+1.444±0.165x	7.634
Thiamethoxam	29.465	22.995	36.818	205.013	128.293	477.556	Y = 2.131+1.952±0.305x	2.468

*All lines are at best fit at P=0.05

Table 3. Median lethal concentration of insecticides to *Aulachophora indica* at 48 hours after treatment

Insecticides	LC ₅₀ (ppm)	95 % fiducial limits of LC ₅₀		LC ₉₅ (ppm)	95 % fiducial limits of LC ₉₅		Regression equation	X ² value*
		Lower	Upper		Lower	Upper		
Fipronil	4.608	3.253	8.708	46.585	28.847	103.430	Y = 3.914+1.637±0.233x	7.426
Cyantraniliprole	10.128	7.762	17.918	54.242	38.643	94.892	Y = 2.731+2.257±0.319x	3.691
Fluxametamide	9.863	8.057	13.727	37.445	28.240	60.610	Y = 2.178+2.839±0.419x	3.113
Phenthoate	10.033	6.985	19.230	223.682	113.968	682.525	Y = 3.778+1.220±0.166x	1.222
Thiamethoxam	20.353	16.052	36.818	205.013	90.982	324.811	Y = 2.487+1.920±0.268x	0.733

*All lines are at best fit at P=0.05

respectively and the corresponding LC₉₅ values were 65.396, 87.077, 42.090, 205.013 and 203.017 ppm, respectively.

At 48 HAT, the LC₅₀ values obtained were 4.608, 10.128, 9.863, 20.353 and 10.033 ppm for fipronil, cyantraniliprole, fluxametamide, thiamethoxam and phenthoate, respectively (Table 3). Among the five different groups of insecticides, the phenyl pyrazole insecticide fipronil exhibited the highest level of toxicity to beetles followed by isoxazoline insecticide, fluxametamide against *A. foveicollis*. Thiamethoxam on the other hand, was found to be the least effective among all the insecticides used with LC₅₀ values of 29.465 and 20.353 ppm at 24 and 48 HAT.

Based on the obtained results (LC₅₀), the order of toxicity of different insecticides at 24 and 48 HAT was as follows: fipronil > fluxametamide > phenthoate > cyantraniliprole > thiamethoxam.

The present study revealed that fipronil was found to be effective against pumpkin beetle and this was in accordance with the findings [9], that stated tolfenpyrad 15% EC at 124.5 and 150 g a.i./ ha as well as fipronil 5% SC at 40 g a.i./ ha reduced the population of red pumpkin beetles, *A. foveicollis* during both *Kharif* and *Rabi* seasons. Fipronil 80 WG at 40 and 50 g a.i./ ha reduced the flea beetle, *Scelodonta strigicollis* (Coleoptera: Chrysomelidae) damage in grapes by 80.50, 80.80 and 83.20, 83.80 per cent and 86.70, 87.10 and 81.60, 82.10 per cent flea beetle reduction, respectively in the two year field trials [10]. Carbaryl 85% (30 g), fipronil 5% SC (20 ml), tolfenpyrad 16% EC (20 ml), cyantraniliprole 10% OD (20 ml) and dinotefuran 10% SL (20 ml) were effective in controlling red cucumber leaf beetle, *Aulacophora indica* and it is in accordance with the present findings.

The second best insecticide is fluxametamide which is a newer insecticide molecule registered for the management of insect pests belonging to Lepidoptera, Coleoptera and Thysanoptera (Gope *et al.*, 2022) [11] and its efficacy was already reported against diamondback moth, *Plutella xylostella*. The effect of phenthoate against pumpkin beetle was in line with the findings of an organophosphorous compound, malathion which at 0.5 per cent was effective in controlling *A. foveicollis* in sweet gourd, bitter gourd and bottle gourd both in field and laboratory conditions [12]. Cyantraniliprole 10 OD at 90 g a.i./ ha was found to be effective against *A. foveicollis* [13].

The least efficacy of thiamethoxam deviated from the findings of Gharde (2019) [14] who stated that thiamethoxam 25 WG at 4 g/l was found highly effective against *A. foveicollis* by reducing the beetle population upto 90 per cent.

CONCLUSION

From the present study, it is concluded that among the five commonly used insecticides in cucurbit ecosystem belonging to five different insecticide groups, fipronil was found to be highly effective against pumpkin beetle, *A. foveicollis* followed by isoxazoline group of compound, fluxametamide and which can be recommended for the management of *A. foveicollis* in cucurbits.

REFERENCES

1. Painkra KL, Archana Kerketta, Madan Kumar Jha. Pest management of cucurbits. *Journal of Entomology and Zoology Studies* 2019; 7(2):1371-1373
2. S.K. Gharde, Mohit Kumar, D. Mal, and A.M.Raut. 2019. Population dynamics and management of pumpkin beetle *Aulacophora foveicollis* Lucas in bottle gourd. *Think India Journal*, 22(34): 1085-1090
3. Jori, R. and Johri, P. K. 2003a. Food preference of red pumpkin beetle, *Aulacophora foveicollis* Lucas at Kanpur in Uttar Pradesh Journal-article Journal-of-Applied-Zoological-Researches Cuttack, India: Applied Zoologists Research Association 14(1): 80-81
4. Atwal, A.S. and Dhaliwal, G.S. 2002. Agricultural Pests of South Asia and their Management, Kalyani publishers, pp 262-263

5. Rashid, M.A., Khan, M.A., Arif, M.J. and N. Javed. 2014. Red Pumpkin Beetle, *Aulacophora foveicollis* Lucas; A Review of Host Susceptibility and Management Practices. *Academic Journal of Entomology* 7 (1): 38-54
6. Stanley, J. and Preetha, G. (2016). Pesticide toxicity to pollinators: Exposure, toxicity and risk assessment methodologies. *Pesticide Toxicity to Non-target Organisms: Exposure, Toxicity and Risk Assessment Methodologies*, 153-228.
7. Abbott, W. S. (1925). A method of computing the effectiveness of an insecticide. *J. econ. Entomol.*, 18(2), 265-267
8. Regupathy A, Dhamu KP. 2001. Statistics workbook for insecticide toxicology. Coimbatore: Softeck Computers. p. 206.
9. Mahato S and Misra HP. 2016. Bioefficacy of a new chemical tolfenpyrad 15% EC against red pumpkin beetle, *Aulacophora foveicollis* (L.). infesting cucumber. In: Extended Abstract of the *National symposium on New Horizon in Pest Management for sustainable Development Goals*, December 23-24, OUAT, Bhubaneswar, pp, 110-111.
10. Prema, M.S., S Sridharan, S Kuttalam and VS Indhumathi. 2019. Efficacy of novel phenyl pyrazole insecticide fipronil 80 WG on flea beetle, *Scelodonta strigicollis* (Motschulsky) in grapes. *International Journal of Chemical Studies* 2019; 7(6): 2594-2602
11. Gope, A., Chakraborty, G., Ghosh, S.M., Sau, S., Mondal, K., Biswas, A., Sarkar, S., Sarkar, P.K. and Roy, D., 2022. Toxicity and sublethal effects of fluxametamide on the key biological parameters and life history traits of diamondback moth *Plutella xylostella* (L.). *Agronomy*, 12(7), p.1656.
12. Hasan MK, Uddin MM and Haque MA. 2011. Efficacy of malathion for controlling red pumpkin beetle, *Aulacophora foveicollis* (Lucas) in cucurbitaceous vegetables. *Progressive Agriculture*, 22(1 & 2): 11-18.
13. Balikai, R. A., & Mallapur, C. P. (2015). Bio-efficacy of cyazypyr 10% OD, a new anthranilic diamide insecticide, against the insect pests of gherkins and its impact on natural enemies and crop. *Journal of Experimental Zoology, India*, 18(1), 89-96.
14. JETIR December 2018, Volume 5, Issue 12 [www.Jetir.org\(ISSN-2349-5162\)](http://www.Jetir.org(ISSN-2349-5162)) JETIRD06181
Journal of Emerging Technologies and innovative Research(JETIR) www.jetir.org 1106
Management of pumpkin beetle in cucurbitaceous crop: A review S.K. Gharde* 1106-1109

UNDER PEER REVIEW