

An outlook on GWAS with a special focus on solanaceous vegetable crops – A Review

ABSTRACT

The commercial solanaceous group of vegetables *viz.*, Tomato, Brinjal, Peppers, and Potatoes hold a prime position among all other vegetables in terms of cultivation and consumption worldwide. These vegetables are consumed fresh by all strata of people and are also used as raw materials in the processing industries. Hence, most plant breeders have a keen interest in the crop improvement of these crops emphasizing yield improvement, quality and nutraceuticals enrichment and imparting resistance to various biotic and abiotic stresses in accordance with climate change. Most of the desirable traits are polygenic in nature which may have a complex inheritance pattern. It is essential to have clear-cut knowledge of the expression frequencies of multiple genes/alleles governing a particular quantitative trait/phenotype, the methods employed to examine and comprehend the candidate genes governing one or various characteristics in plants are quantitative trait loci mapping and genome-wide association studies. In This review, we have discussed GWAS and its future perspectives and made efforts to tabulate the recent past works done in solanaceous vegetable crops using this approach. GWAS is a attractive strategy that constitutes an important advancement in genetic analysis and has unquestionably proven to be a useful tool in identifying candidate genes, which is justified in this article.

Keywords: Genome-wide association study (GWAS), Quantitative trait loci (QTL), Genotyping, Candidate gene, Single nucleotide polymorphism (SNP), Vegetables, Solanaceous

Abbreviations:

QTLs - Quantitative trait loci
RIL - Recombinant inbred line
LD - Linkage disequilibrium
GWAS - Genome-wide association study
SNP - Single-nucleotide polymorphism
WGS - Whole genome sequencing
GS - Genomic selection
NGS - Next-generation sequencing
MAS - Marker-assisted selection
GBS - Genotyping-by-sequencing
GLM – Generalised linear model
SSR – Single sequence repeats
RADseq - Restriction site-associated DNA sequencing
ddRAD-seq - Double digest restriction-site associated DNA sequencing

INTRODUCTION

Vegetable breeders' and researchers' key priorities are to increase yield, quality, and impart resistance to various biotic and abiotic challenges in order to ensure food and nutritional security for the fast-expanding population and to reduce the problem of hidden hunger. Vegetable genetic improvement is hampered by several biotic and abiotic variables, and these factors may also reduce the quantity and quality of vegetables, especially before and after harvest (Rikkerink *et al.* 2007). It is anticipated that enhanced genetic designs, agro-biotechnology, advanced plant breeding methods and genomics will be used to meet the problem. The features like yield and quality in vegetable crops that are economically important and are primarily regulated via multi-allelic or polygenic genes found using quantitative trait loci (QTLs) that interact with one another and the surroundings (Khan, M. A, & Korban, S. S. 2012). To break down complicated variables and find genetic variations linked to these traits, two mapping techniques, linkage mapping/QTL mapping and association mapping are frequently utilized (Zhu *et al.*, 2008, Sukumaran, S, & Yu, J. 2014). Since both approaches lay foundation on the co-inheritance of functional polymorphisms (genes which alter phenotype) and DNA modifications (molecular markers), they can be used to **locate QTL associated with vegetable yield and quality (Devi *et al.*, 2022).**

As previously mentioned, the techniques used to investigate and comprehend various characteristics of plants include quantitative trait loci (QTL) mapping and genome-wide association studies (GWAS). A statistical approach known as QTL mapping makes it easier to link multiple genotypes and phenotypes (Lynch and Walsh, 1998; Falconer and Mackay 1996). Molecular markers including single nucleotide polymorphisms (SNPs) and amplified fragment length polymorphisms (AFLPs) are used to link mapped QTLs to reported phenotypic information. Multiple types of investigations employing QTL mapping have been reported for plant species to examine several attributes associated to production, quality, and various biotic and abiotic stresses (Fan *et al.*, 2015; Mora *et al.*, 2016; Gahlaut *et al.*, 2017). QTL mapping has the benefit that it enables the mapping of trait variation in F₂ RIL populations. The same benefit revealed QTL mapping's shortcomings, wherein only RIL populations can be mapped and allelic variation among parents can be measured at an F₂ hybrid (Korte and Farlow, 2013). GWAS outperforms QTL mapping's shortcomings. In this review, we made an effort to discuss the GWAS approach and the past few years' works were highlighted along with the future perspective of GWAS in solanaceous vegetable crops. We believe that this information will provide an overall outlook on GWAS and help researchers to find out the research gaps in the improvement of these crops.

GENOME-WIDE ASSOCIATION STUDY

A study of genome-wide association, commonly referred to as whole genome association studies, evaluate a collection of genetic changes over the entire genome to see whether any variation is linked to a trait (Manolio, 2010). The focus of GWAS is frequently on the connections among characteristics and SNPs, for instance, DNA from diverse plant species is when related to varied phenotypes for a specific character. GWAS, despite being hypothesis-free, can generate hypotheses. GWAS are used for association result analysis, replication, interpretation, and

follow-up based on designs, genotyping technologies, and statistical principles (Bush and Moore, 2012).

GWAS Design

An effective study needs a consistent approach or design. The DNA of several plants with various phenotypes for a given characteristic is frequently compared in case versus control, or quantitative, GWAS designs (Bush and Moore, 2012). Studies that are trait-focused can employ case vs. control research, whereas studies that are trait-measurement oriented can use quantitative research with a quantitative design. When comparing a case to a control, the variety could be the case itself, another variety like it without any modifiable traits, or a variety with different phenotypes of the same trait. Every variation is often classified as either a case or a control. In some situations, it may be straightforward to quantify a measurable characteristic that is included in a statistical design since the study can take genetic variations into consideration (Bush and Moore, 2012). The efficiency of the GWAS design for gathering SNP (single nucleotide polymorphism) data will determine the outcome of the investigation.

Mapping SNPs

SNPs are the current genetic diversity units evidenced in the genomes of all organisms. They are DNA mutations that affect only one base pair. Utilizing genotyping technologies such as tailored, chip-based microarray technology (Kumar et al., 2015), Illumina (San Diego, California), Affymetrix (Santa Clara, California) (Distefano and Taverna, 2011), or other next-generation sequencing technologies, more than a million SNPs can be detected (Neelapu and Surekha, 2016). Based on GWAS design, these collected SNPs can be examined or linked to phenotypes or traits.

GWAS Assessment

The linkage (collection of genes/alleles) of the heritable unit for a population's character is transmitted from parent to offspring. An assessment of changes in the linkage that can be linked to mutations, linkage breaks, or SNPs is called linkage disequilibrium (LD). Evaluation of SNPs in the genome that affects phenotype is the aim of GWAS. There are two methods for evaluating SNPs that affect phenotypes: both direct and indirect associations (Bush and Moore, 2012). Indirect association happens whenever the SNP is not genotyped directly but is correlated statistically with the phenotype. Direct association occurs when the trait's SNP is precisely genotyped and analytically connected to the SNP affecting trait. (Bush and Moore, 2012). For single-locus or multi-locus analyses, the association of alleles and SNP in a sample is performed. (Padmavathi et al., 2003; Devi et al., 2006, 2007). An analysis of the independent connections between each SNP and phenotype is known as single-locus statistical analysis. When the Each SNP's relationship to a phenotype or trait is investigated, multi-locus analysis takes place (Padmavathi et al., 2003; Devi et al., 2006, 2007; Bush and Moore, 2012).

Logistic regression will be used to analyse traits whenever the quantitative approach is used in the GWAS study. When the GWAS investigation is for a case vs control design, a generalised linear model (GLM) and analysis of variance (ANOVA) will be utilized to assess traits. The null hypotheses or assumptions that govern the three techniques of GLM, ANOVA, and logistic regression when the attribute follows normal distribution, the trait variance within each group is

the same, and the groups are independent (Bush and Moore, 2012). Even while there are benefits to utilising GWAS to relate SNPs to traits or qualities, there are also some disadvantages. Population stratification makes it more likely for SNPs to be linked to traits during GWAS analysis. Typically, genetic analysis acts as a sign of substructure of the population (Bush and Moore, 2012). The segmentation of the population can be avoided by investigating population structures and using PCA (Principal component analysis) to eliminate influences among the information (Bush and Moore, 2012).

GWAS replication

The influence of genetics applies to various groups, and a replication leads to generalization of the link between SNPs and characteristics. To validate the results of the population, repetition experiments must be carried out using separate datasets, as with GWAS, generated from comparable samples. (Bush and Moore, 2012). The link of SNPs associated with characteristics is supported if the findings hold up throughout case studies and replication studies (Challa and Neelapu, 2018).

Meta-analysis of GWAS

Since the first successful publication in 2005, innumerable case studies on GWAS have already been reported. On a common hypothesis, scientific articles typically concur or disagree. Publications employ meta-analysis to clarify this misunderstanding. To explore the similar assumption and determine the usual result of integrated study, meta-analysis is a potent statistical technique that brings together GWAS data from multiple published studies (Bush and Moore, 2012; Challa and Neelapu, 2018). The data from merged studies should be imputed before being used in meta-analysis because they were collected from different genome sequence platforms and in the study, no SNPs were genotyped directly. (Bush and Moore, 2012; Challa and Neelapu, 2018).

This strategy has a number of benefits, such as the ability to generalise the study's findings to a larger group. Techniques can increase precision and accuracy of the estimations. Comparing and contrasting the results of various GWAS reports is possible. To test the hypothesis, the meta-analysis may use, for example, an overview of the findings of a GWAS study (Bush and Moore, 2012).

GWAS IN SOLANACEOUS VEGETABLES

The Solanaceae contain many horticultural species of economic importance known for their food and medicinal values, among which the most common and commercial vegetable crops include tomato (*Solanumlycopersicum*), eggplant (*Solanummelongena*), pepper (*Capsicum spp.*) and potato (*Solanumtuberosum*). All these four vegetables are the most important vegetables throughout the world among different strata of people. Most of the yield and quality characteristics of these vegetables are mostly governed by polygenes. As we discussed earlier genome-wide association studies along with linkage disequilibrium are one of the best techniques to dissect complex traits. Hence for effective crop improvement of these crops many researchers are focusing more on research employing GWAS. So far only countable work has been done in this area concerning these vegetable crops and a lot more is yet to be explored. We attempted to tabulate some recent works done in solanaceous vegetable crops using GWAS approaches are given below:

TOMATO

Sl.No	Aim of study	Techniques used	Results	References
1	To study the genomic diversity and population structure analysis	ddRAD-seq	In the Mediterranean 'da serbo' gene pool, 2297 new genes with harboring SNPs were identified which can be subjected to further study using GWAS	Esposito, S (2020)
2	To find QTLs for eight tomato fruit attributes	SNPs (51K Axiom® tomato array)	A number of traits were linked to two SNP markers on chromosome 1 and 2, demonstrating the pleiotropic effects of QTL. Furthermore, eight fruit attribute potential new QTLs are offered by 16 of 30 MTAs (marker-trait associations).	Kim, M (2021)
3	determining the genetic influences on agronomical qualities in contemporary breeding lines, and assessing their genetic diversity	GWAS, Illumina	<ul style="list-style-type: none"> • Over 11,000 SNPs were discovered. • Along with numerous new loci important for tomato breeding, the genetic architecture and connections for fruit weight, plant development, and earliness were discovered. 	Bauchet, G., (2017a)
4	To identify genomic regions for four fruit characteristics they are fruit weight (FW), soluble solid content (SSC), fruit shape, and fruit color and to find associations between them.	GWAS, 547 InDel markers	A group of 102 markers that were substantially ($p < 1.19-1.30 \times 10^{-4}$) linked with SSC, FW, fruit morphology, and fruit colour was discovered on 11 of the 12 chromosomes. The findings showed that association mapping utilising a small population and a limited range of InDel markers might not only supplement and improve prior QTL data but also uncover new loci enabling marker-assisted selection of tomato fruit characteristics.	Liu, X., (2017)
5	To assess the genetic differentiation and linkage disequilibrium in	GWAS, RADseq	<ul style="list-style-type: none"> • In order to account for population stratification, SNP genotypic data consistently verified the 	Lin, Y. P., (2019)

	<i>Solanum pimpinellifolium</i>		<p>genetic differentiation.</p> <ul style="list-style-type: none"> When markers are saturated in GWAS, the 18-Kb linkage disequilibrium decay does in fact show the possibility for single-gene resolution. 	
6	To detect quantitative trait loci (QTL) for fruit traits in tomato	GWAS	A total of 41 loci associated with the six fruit traits <i>viz.</i> fruit shape, fruit color, pericarp thickness, fruit weight, fruit height, and fruit width were identified	Phan, N. T., (2019)
7	To study the genetic mapping for earliness-related traits in cultivated tomato	Molecular markers (SNPs, InDels, and SSRs)	The study found 23 associations for variables linked to earliness, confirmed some prior QTLs, and found several possibly new loci that regulate earliness in tomato.	Wang, T., (2020)
8	To locate key genetic loci and regions that regulate the acid and volatile content of tomato fruit	GWAS	<ul style="list-style-type: none"> In the study, 10000 SNP markers were detected, and 79 relationships that had significant impacts at high resolution on 13 major and 19 secondary metabolites were discovered. The study also identified potential genes that could alter the metabolite content. 	Bauchet et al., (2017b)
9	To describe the genetics of malate synthesis	GWAS	A quantitative trait locus responsible for variation in fruit malate accumulation was identified <i>i.e.</i> TFM6 (tomato fruit malate 6), which corresponds to AL-ACTIVATED MALATE TRANSPORTER9 (SI-ALMT9 in tomato).	Ye et al., (2017)
10	To locate the QTL in organic agriculture conditions for agronomic, fruit quality, and root architectural traits.	Phenotyping and GWAS (dd- RADseq)	<ul style="list-style-type: none"> High heritability was observed among different characters under study. On chromosome 11, a single member of the SUN family (SLSUN31) was linked to six flower and fruit characteristics. 	Tripodi et al., (2021)

11	To look into the quantitative trait loci (QTL) for resistance to bacterial wilt.	GWAS (51 K Axiom® tomato array)	Three genes pertaining to bacterial wilt resistance were identified viz., <i>Bwr-4</i> , <i>Bwr-12</i> and <i>Bwr-6</i> on chromosome 4, 12 and 6 respectively.	Nguyen et al., (2021)
12	To study the regulation and formation of multicellular trichomes in tomato	GWAS	The study revealed the key gene controlling trichome formation i.e., <i>Hair (H)</i> along with its three alleles.	Chang et al., (2018)
13	To identify the genetic factors that contribute to the buildup of tocochromanol in tomato fruits	GWAS and mapping	The study identified a total of 25 quantitative trait loci pinpointing the chorismate-tyrosine pathway.	Burgos et al., (2021)
14	To recognise the lipase gene and its relationship to flavour volatiles produced from short-chain fatty acids in tomato fruits.	GWAS and mapping	The analysis identified a gene <i>Sl-LIP8</i> , which is connected to the buildup of volatile organic molecules produced from short-chain fatty acids in tomato fruit hence contributing to the flavor of the tomato fruits.	Li et al., (2020)

BRINJAL

Sl.No	Aim of study	Techniques used	Results	References
1	To comprehend the process through which the eggplant fruit peel produces chlorophylls	GWAS	The results showed that the genes <i>APRR2</i> and <i>SmAPRR2</i> were found having association with the fruit peel colour in brinjal	Arrones et al (2022)
2	To determine the QTLs in aubergine that control anthocyanin pigmentation and fruit colour	GWAS	56 SNP locus/phenotype relationships were discovered by the study, and the genomic areas carrying these loci were dispersed over nine of the 12 chromosomes of eggplant.	Cericola et al (2014)
3	To detect the candidate genes for anthocyanin pigmentation in MAGIC population of Brinjal	GWAS, Phenotyping and Genotyping (5k probes SPET platform)	Strong connections between MYB113 and two myeloblastosis (MYB) genes involved in the anthocyanin biosynthesis pathway were found.	Mangino et al (2022)
4	To generate high resolution genetic map and to locate the	GWAS	For each characteristic, the study found 7 to 17 QTLs (at least one significant QTL). And the	Toppino et al (2020)

	QTL for anthocyanin pigmentation and Seed vigour		outcomes showed that the newly established map offers useful information for brinjal crop improvement.	
5	To uncover the genetic structure and diversity by studying the selection based on fruit shape	GWAS, Genotyping	Five related SNPs around the SUN and OVATE homologs were found by a study based on the 219 SNPs, and they had a conserved function in regulating the form of the fruit.	Liu et al (2019)

PEPPERS

Sl.No	Aim of study	Techniques used	Results	References
1	To investigate the population structure, genetic diversity and links among the landrace varieties of pepper (<i>Capsicum spp.</i>) from the Spanish Center of Diversity.	Genotyping by-sequencing (GBS)	<ul style="list-style-type: none"> The population under study was divided into seven clusters by using discriminant analysis of principal components (DAPC) A tight relationship between Mexican and Spanish accessions was revealed via phylogenetic analyses. 	Pereira-Dias et al (2019)
2	To understand the underlying genetics of Chile pepper's yield and agronomic characteristics (<i>Capsicum spp.</i>) by employing multi-locus association mapping.	GWAS	Eight SNP markers spread over six chromosomes were discovered to be associated with a number of traits.	Lozada et al (2022)
3	To investigate the genetic variation and population dynamics of Ethiopian Bell pepper accessions.	GBS	<ul style="list-style-type: none"> Two different genetic populations made up of nine <i>C. frutescens</i> accessions and 132 <i>C. annuum</i> accessions respectively made up the 142 capsicum collections. 509 SNP markers were found by GWAS analysis to be strongly correlated with characteristics related to fruit, stems, and leaves. 	Solomon et al (2019)
4	To identify the potential genes	GWAS & QTL Mapping	The GWAS results revealed that 16 casual genes were	Lee et al (2020)

	responsible for features connected to fruit		believed to be connected to characteristics linked to fruit morphology.	
5	To investigate genetic resistance and generate <i>Phytophthora</i> blight disease markers.	GWAS, GBS	The findings showed an SNP marker termed Chr02-1126, which is positioned at 112 Mb on Chr.02 and exhibits <i>P. capsici</i> resistance with 78.5% accuracy.	Ro et al (2022)
6	To identify genomic loci and alleles responsible for resistance to bacterial wilt	GBS	The QTL analysis showed that the gene <i>pBWR-1</i> was found on chromosome 01 reported to be involved in imparting resistance against bacterial wilt	Chae et al (2022)
7	To investigate the correlations between 36 agronomic traits throughout the genomes of 287 pepper accessions.	GWAS, specific-locus amplified fragment sequencing (SLAF-seq)	The results revealed that 109 correlation peaks were significantly associated with the 26 agronomic characteristics via GWAS, which can be further studied for the crop improvement of pepper	Wu et al (2019)
8	To evaluate the genetic diversity and relationship between the amount of capsaicinoids in Thai chilli landraces	GWAS, Diversity array genotype-by-sequencing (DARTseq)	<ul style="list-style-type: none"> • With a dissimilarity score ranging from 0.0092 to 0.841, cluster analysis separated the chilli landraces into two distinct groups based on their species, <i>Capsicum annuum</i> and <i>C. frutescens</i>. • GWAS analysis revealed significant association between seven SNPs with the capsaicinoids content. 	Kethomet al (2019)
9	To locate and catalogue the gene in mature peppers that regulates anthocyanin production	GWAS, Mapping and Sequencing	Using a map-based cloning method, the anthocyanin biosynthesis gene <i>Ca3GT</i> was precisely mapped in a 110.5 kb region and the strong candidate gene <i>Capana10g001978</i> was confirmed.	Liu et al (2020)
10	To identify and locate the gene responsible for producing light-green immature peppers.	GWAS, BSA (bulked segregant analysis)	The study identified a 35.07 kbp region on chromosome 10 as the LG locus affecting light-green immature pepper fruits. In this region, <i>CaPP2C35</i> , a	Wu et al (2022)

			potent candidate gene, was discovered.	
11	To study the genomic diversity of fruit morphology in capsicum	GWAS	The fruit morphology phenotypes were linked to four unique loci, including a non-synonymous mutation in the Longifolia 1-like (<i>CA03g16080</i>) gene of the pepper plant.	Colonna et al (2019)

POTATO

Sl.No	Aim of study	Techniques used	Results	References
1	To investigate the roles and functions of MYB genes in anthocyanin biosynthesis in potato.	GWAS and transcriptomic analysis	The findings showed that the two MYB TFs (StMYB88 and StMYB89) on chromosome 10 were connected to the development of anthocyanin.	Li et al (2021)
2	To identify potential genes associated with potato late blight resistance	GWAS	According to the whole genome association study, there were 18 candidate genes in lesion diameter and 22 candidate genes for disease resistance grade (late blight).	Wang et al (2020)
3	The study sought to discover genomic areas associated with the morphology of potato starch granules in <i>Solanum tuberosum</i> L.	GWAS	The study demonstrated genetic diversity for starch granule morphology. Significant SNPs were found on chromosomes 2, 4, 7, and 11. With this candidate genes can be found in the future.	Erst et al (2019)
4	The study was carried out to reveal genetic variations and genetic architecture associated with Chip Processing clones	GWAS & Sweeps analysis	The Sweeps analysis and GWAS results identified the related genetic variations in chromosome 10, along with variants under selection throughout the genome.	Jo et al (2022)
5	The study sought to investigate the phenotypic diversity and genetic basis of essential floral traits in potatoes.	GWAS, GBS	The results revealed a total of 15 significant SNPs for stamen length, pistil length, and pistil length above stamen across both years. The identified candidate genes from this study might be useful to the potato breeders.	Zia et al (2022)
6	To investigate genetic	GWAS	The findings indicated nine	Diaz et al

	variation in response to water deprivation in a panel of diploid potatoes and find the QTL that governs this feature.		QTL on chromosomes 2, 3, 5, 8, 10, and 12, as well as three candidate genes with a possible function in water deprivation response.	(2021)
7	To identify genomic regions associated with starch granule morphology and to increase number of known genome loci associated with potato starch yield.	GWAS	The study discovered 53 new SNPs on chromosomes 1, 2, 4, 5, 6, 7, 9, 11, and 12 that are linked to tuber starch preparative yields and starch granules morphology.	Khlestkin et al (2020)
8	To find QTLs and potential candidate genes linked to protein levels in a sample of tetraploid potatoes.	GWAS	The study explored and pinpointed potential genes. QTLs have been discovered on chromosomes 3, 5, 7, and 12.	Klaassen et al (2019)
9	To find loci those are strongly linked to common scab disease resistance in a Canadian potato germplasm.	GWAS	The study found significant SNPs for potato common scab resistance on potato chromosomes 2, 4, and 12.	Yuan et al (2020)
10	To identify novel QTLs for common scab resistance in potato	GWAS	This resulted in the discovery of an unique QTL for common scab resistance that accounted at the most 14.7% of the total variance; this QTL was located on potato chromosome 1 in the 640 kb gap between 0.43 Mb and 1.07 Mb.	Koizumi et al (2021)

Future perspectives of GWAS in vegetable crops

Precision marker design and development for significant vegetable crops with high economic value have proven to be costly for many years because of the significant expense of sequencing and genotyping. In addition, a recent development in sequencing dubbed as "next generation sequencing technology" (NGS) has significantly impacted the costs and efforts associated with whole-genome sequencing (Khan *et al*, 2014; Zahidet *al*, 2022), enabling the creation of GWAS and GS (Genomic Selection). Regarding the accuracy and uniqueness of these two methods, they are going to become important tools for next vegetable breeding programmes as well as effective genetics applications. It will be possible to accurately estimate the genetic value and worth of untested lines and genotypes through the generation of genome-wide marker data that could be advantageous by combining low-cost and high-throughput genotyping techniques. This might be accomplished using genome-wide SNPs with better statistical measures with high precision and

efficacy (Zahidet *al*, 2022). The genetics of vegetable crops are constantly being worked on by researchers using genomics-based breeding and genomics-assisted selection. By analyzing the yield and quality of the vegetables as well as characteristics such as storability, firmness, weight, shape, colour, maturity time, size, nutrient content, and chlorophyll index through GWAS, it is possible to ascertain whether any of the found markers are associated with particular traits (O'Connor *et al*, 2018). Vegetable breeders can create novel cultivars with favorable characteristics using genome-wide molecular assisted selection by creating high-density reference genetic linkage maps using genomic resources from the most recent advancements in whole-genome sequencing, genome-wide SNPs, and conventional SSR markers (Yamamoto, T. &Terakami, S. 2016).

Without a doubt, breeding efforts for crops and vegetables have greatly benefited from genome-wide association analyses. Researchers and vegetable breeders have devised new genetic risk factors for prevalent crop illnesses. Once the entire vegetable genome has been sequenced, it is anticipated that the legitimacy and effectiveness of GWAS in vegetable crops would greatly rise. This association study methodology is noteworthy in that it may measure relationships between genetic variation and a variety of phenotypes without requiring any prior knowledge. Additionally, GWAS can analyze the hitherto untapped genetic variety in vegetable germplasm supplies, revealing fresh genetic variability that can be used in upcoming breeding initiatives. Additionally, GWAS can enable transgenesis and target mutagenesis techniques like genome editing (O'Connor, K. M *et al.*, 2021; Zahidet *al*, 2022) to assess whether discovered genes are responsible for the phenotypic variance of traits of interest after QTL mapping and gene cloning of candidate genes (Zahidet *al*, 2022).

Conclusion

Finally, and perhaps most crucially, collecting phenotypic and genotypic data from vegetable breeders will eventually broaden the possibilities for detecting the potency of GWAS and the accuracy of GS. Furthermore, it is anticipated that the application of NGS technologies would advance yield advantages in all significant vegetable crops. Every major vegetable species is currently the subject of collaborative efforts, and we anticipate the results of these efforts, which will eventually sharpen our viewpoint and support the creation and enhancement of effective routes (Tuberosa *et al.*, 2014; Zahidet *al*, 2022).

REFERENCES

1. Arrones, A., Mangino, G., Alonso, D., Plazas, M., Prohens, J., Portis, E., &Gramazio, P. (2022). Mutations in the SmAPRR2 transcription factor suppressing chlorophyll pigmentation in the eggplant fruit peel are key drivers of a diversified colour palette. *Front. Plant Sci.* 13:1025951. doi: 10.3389/fpls.2022.1025951
2. Bauchet, G., Grenier, S., Samson, N., Bonnet, J., Grivet, L., &Causse, M. (2017a). Use of modern tomato breeding germplasm for deciphering the genetic control of agronomical traits by Genome Wide Association study. *Theoretical and applied genetics*, 130(5), 875-889.
3. Bauchet, G., Grenier, S., Samson, N., Segura, V., Kende, A., Beekwilder, J., &Causse, M. (2017b). Identification of major loci and genomic regions controlling acid and volatile

- content in tomato fruit: implications for flavor improvement. *New Phytologist*, 215(2), 624-641.
4. Burgos, E., Belen De Luca, M., Diouf, I., de Haro, L. A., Albert, E., Sauvage, C., & Carrari, F. (2021). Validated MAGIC and GWAS population mapping reveals the link between vitamin E content and natural variation in chorismate metabolism in tomato. *The Plant Journal*, 105(4), 907-923.
 5. Bush, W.S. and Moore, J.H., 2012. Genome-wide association studies. *PLoS Comput. Biol.* 8 (12), e1002822.
 6. Cericola, F., Portis, E., Lanteri, S., Toppino, L., Barchi, L., Acciarri, N., & Rotino, G. L. (2014). Linkage disequilibrium and genome-wide association analysis for anthocyanin pigmentation and fruit color in eggplant. *BMC genomics*, 15(1), 1-15.
 7. Chae, S. Y., Lee, K., Do, J. W., Hong, S. C., Lee, K. H., Cho, M. C., Yang, E. Y., & Yoon, J. B. (2022). QTL Mapping of resistance to bacterial wilt in pepper plants (*Capsicum annuum*) using genotyping-by-sequencing (GBS). *Horticulturae*, 8(2), 115.
 8. Challa, S., & Neelapu, N. R. (2018). Genome-wide association studies (GWAS) for abiotic stress tolerance in plants. In *Biochemical, physiological and molecular avenues for combating abiotic stress tolerance in plants*. Academic Press. pp. 135-150.
 9. Chang, J., Yu, T., Yang, Q., Li, C., Xiong, C., Gao, S., & Ye, Z. (2018). Hair, encoding a single C₂H₂ zinc-finger protein, regulates multicellular trichome formation in tomato. *The Plant Journal*, 96(1), 90-102.
 10. Colonna, V., D'Agostino, N., Garrison, E., Albrechtsen, A., Meisner, J., Facchiano, A., ...& Tripodi, P. (2019). Genomic diversity and novel genome-wide association with fruit morphology in *Capsicum*, from 746k polymorphic sites. *Scientific reports*, 9(1), 1-14.
 11. Devi, K.U., Reineke, A., Rao, C.U.M., Reddy, N.N.R., Khan, A.P.A., 2007. AFLP and single-strand confirmation polymorphism studies of recombination in the entomopathogenic fungus *Nomuraearileyi*. *Mycol. Res.* 111, 716–725.
 12. Devi, K.U., Reineke, A., Reddy, N.N.R., Rao, C.U.M., Padmavathi, J., 2006. Genetic diversity, reproductive biology and speciation in the entomopathogenic fungus *Beauveria bassiana* (Balsamo) Vuillemin. *Genome* 49 (5), 495–504.
 13. Díaz, P., Sarmiento, F., Mathew, B., Ballvora, A., & Mosquera Vázquez, T. (2021). Genomic regions associated with physiological, biochemical and yield-related responses under water deficit in diploid potato at the tuber initiation stage revealed by GWAS. *PloS one*, 16(11), e0259690.
 14. DiStefano, J.K. and Taverna, D.M. (2011). Technological Issues and Experimental Design of Gene Association Studies. In: DiStefano, J. (eds) *Disease Gene Identification. Methods in Molecular Biology*, vol 700. Humana Press, Totowa, NJ. https://doi.org/10.1007/978-1-61737-954-3_1
 15. Erst, T. V., Rozanova, I. V., Khlestkin, V. K., & Khlestkina, E. K. (2019). Search for genomic regions associated with potato starch granules morphology of *Solanum tuberosum* L. *Current Challenges in Plant Genetics, Genomics, Bioinformatics, and Biotechnology*, 24, 217.
 16. Esposito, S., Cardi, T., Campanelli, G., Sestili, S., Díez, M. J., Soler, S., & Tripodi, P. (2020). ddRAD sequencing-based genotyping for population structure analysis in cultivated tomato provides new insights into the genomic diversity of Mediterranean 'da serbo' type long shelf-life germplasm. *Horticulture research*, 7.

17. Falconer, D.S., Mackay, T.F.C., 1996. Introduction to Quantitative Genetics, fourth ed. Longmans Green, Harlow.
18. Fan, Y., Shabala, S., Ma, Y., Xu, R., Zhou, M., 2015. Using QTL mapping to investigate the relationships between abiotic stress tolerance (drought and salinity) and agronomic and physiological traits. *BMC Genomics* 16 (1), 43. <https://doi.org/10.1186/s12864-015-1243-8>
19. Gahlaut, V., Jaiswal, V., Tyagi, B.S., Singh, G., Sareen, S., Balyan, H.S., Gupta, P.K., 2017. QTL mapping for nine drought-responsive agronomic traits in bread wheat under irrigated and rain-fed environments. *PLoS One* 12 (8), e0182857. <https://doi.org/10.1371/journal.pone.0182857>
20. Jo, K. R., Choi, J. G., Kwon, D. H., Park, Y. E., & Kim, S. J. (2022). Genome-wide association study and selective sweeps reveal genetic variations associated with Chip Processing clones selected over sixty years in global potato breeding programs. *Research Square*. DOI: 10.21203/rs.3.rs-1880525/v1.
21. Kethom, W., Tongyoo, P., & Mongkolporn, O. (2019). Genetic diversity and capsaicinoids content association of Thai chili landraces analyzed by whole genome sequencing-based SNPs. *Scientia Horticulturae*, 249, 401-406.
22. Khan, M. A., & Korban, S. S. (2012). Association mapping in forest trees and fruit crops. *Journal of Experimental Botany*, 63(11), 4045-4060.
23. Khan, M. A., Olsen, K. M., Sovero, V., Kushad, M. M., & Korban, S. S. (2014). Fruit quality traits have played critical roles in domestication of the apple. *The Plant Genome*, 7(3), plantgenome2014-04.
24. Khlestkin, V. K., Erst, T. V., Rozanova, I. V., Efimov, V. M., & Khlestkina, E. K. (2020). Genetic loci determining potato starch yield and granule morphology revealed by genome-wide association study (GWAS). *PeerJ*, 8, e10286.
25. Kim, M., Nguyen, T. T. P., Ahn, J. H., Kim, G. J., & Sim, S. C. (2021). Genome-wide association study identifies QTL for eight fruit traits in cultivated tomato (*Solanum lycopersicum* L.). *Horticulture Research*, 8.
26. Klaassen, M. T., Willemsen, J. H., Vos, P. G., Visser, R. G., van Eck, H. J., Maliepaard, C., & Trindade, L. M. (2019). Genome-wide association analysis in tetraploid potato reveals four QTLs for protein content. *Molecular Breeding*, 39(10), 1-12.
27. Koizumi, E., Igarashi, T., Tsuyama, M., Ogawa, K., Asano, K., Kobayashi, A., ...& Hosaka, K. (2021). Association of genome-wide SNP markers with resistance to common scab of potato. *American Journal of Potato Research*, 98(2), 149-156.
28. Korte, A., Farlow, A., 2013. The advantages and limitations of trait analysis with GWAS: a review. *Plant Methods* 9, 29.
29. Kumar, V., Singh, A., Mithra, S.V., Krishnamurthy, S.L., Parida, S.K., Jain, S., Tiwari, K.K., Kumar, P., Rao, A.R., Sharma, S.K., Khurana, J.P., Singh, N.K., Mohapatra, T., 2015. Genome-wide association mapping of salinity tolerance in rice (*Oryza sativa*). *DNA Res.* 22 (2), 133-145.
30. Lee, H. Y., Ro, N. Y., Patil, A., Lee, J. H., Kwon, J. K., & Kang, B. C. (2020). Uncovering candidate genes controlling major fruit-related traits in pepper via genotype-by-sequencing based QTL mapping and genome-wide association study. *Frontiers in Plant Science*, 11, 1100.
31. Li, X., Tieman, D., Liu, Z., Chen, K., & Klee, H. J. (2020). Identification of a lipase gene with a role in tomato fruit short-chain fatty acid-derived flavor volatiles by genome-wide association. *The Plant Journal*, 104(3), 631-644.

32. Li, Y., Liang, J., Zeng, X., Guo, H., Luo, Y., Kear, P., ...& Zhu, G. (2021). Genome-wide analysis of MYB gene family in potato provides insights into tissue-specific regulation of anthocyanin biosynthesis. *Horticultural Plant Journal*, 7(2), 129-141.
33. Lin, Y. P., Liu, C. Y., & Chen, K. Y. (2019). Assessment of genetic differentiation and linkage disequilibrium in *Solanum pimpinellifolium* using genome-wide high-density SNP markers. *G3: Genes, Genomes, Genetics*, 9(5), 1497-1505.
34. Liu, J., Ai, X., Wang, Y., Lu, Q., Li, T., Wu, L., ...& Shen, H. (2020). Fine mapping of the Ca3GT gene controlling anthocyanin biosynthesis in mature unripe fruit of *Capsicum annum* L. *Theoretical and Applied Genetics*, 133(9), 2729-2742.
35. Liu, W., Qian, Z., Zhang, J., Yang, J., Wu, M., Barchi, L., Zhao, H., Sun, H., Cui, Y., & Wen, C. (2019). Impact of fruit shape selection on genetic structure and diversity uncovered from genome-wide perfect SNPs genotyping in eggplant. *Molecular Breeding*, 39, 140, 1-13.
36. Liu, X., Geng, X., Zhang, H., Shen, H., & Yang, W. (2017). Association and genetic identification of loci for four fruit traits in tomato using InDel markers. *Frontiers in plant science*, 8, 1269.
37. Lozada DN, Barchenger DW, Coon D, Bhatta M, Bosland PW. Multi-locus Association Mapping Uncovers the Genetic Basis of Yield and Agronomic Traits in Chile Pepper (*Capsicum* spp.). *Crop Breed Genet Genom.* 2022; 4(2):e220002. <https://doi.org/10.20900/cbagg20220002>
38. Lynch, M., Walsh, B., 1998. *Genetics and Analysis of Quantitative Traits*. Sinauer Associates, Sunderland.
39. Mangino, G., Arrones, A., Plazas, M., Pook, T., Prohens, J., Gramazio, P., & Vilanova, S. (2022). Newly developed MAGIC population allows identification of strong associations and candidate genes for anthocyanin pigmentation in eggplant. *Frontiers in Plant Science*, 13.
40. Manolio, T.A., 2010. Genomewide association studies and assessment of the risk of disease. *N. Engl. J. Med.* 363 (2), 166–176.
41. Mora, F., Quitral, Y.A., Matus, I., Russell, J., Waugh, R., Del Pozo, A., 2016. SNP-based QTL mapping of 15 complex traits in barley under rain-fed and well-watered conditions by a mixed modeling approach. *Front. Plant Sci.* 7(909).
42. Neelapu, N. R. R. and Surekha, C., 2016. Next-generation sequencing and meta-genomics. In: Wong, K.-C. (Ed.), *Computational Biology and Bioinformatics: Gene Regulation*. CRC Press, Boca Raton, FL, pp. 331–351.
43. Nguyen, T. T., Le, N. T., & Sim, S. C. (2021). Genome-wide association study and marker development for bacterial wilt resistance in tomato (*Solanum lycopersicum* L.). *Scientia Horticulturae*, 289, 110418.
44. O'Connor, K. M., Hayes, B. J., Hardner, C. M., Alam, M., Henry, R. J., & Topp, B. L. (2021). Genomic selection and genetic gain for nut yield in an Australian macadamia breeding population. *BMC genomics*, 22(1), 1-12.
45. O'Connor, K., Hayes, B., & Topp, B. (2018). Prospects for increasing yield in macadamia using component traits and genomics. *Tree genetics & genomes*, 14(1), 1-14.
46. Padmavathi, J., Uma Devi, K., Rao, C.U.M., Reddy, N.N.R., 2003. Telomere fingerprinting for assessing chromosome number, isolating typing and recombination in the entomopathogen *Beauveria bassiana*. *Mycol. Res.* 107 (5), 572–580.
47. Pereira-Dias, L., Vilanova, S., Fita, A., Prohens, J., & Rodríguez-Burruezo, A. (2019). Genetic diversity, population structure, and relationships in a collection of pepper (*Capsicum*

- spp.) landraces from the Spanish centre of diversity revealed by genotyping-by-sequencing (GBS). *Horticulture research*, 6.
48. Phan, N. T., Trinh, L. T., Rho, M. Y., Park, T. S., Kim, O. R., Zhao, J., & Sim, S. C. (2019). Identification of loci associated with fruit traits using genome-wide single nucleotide polymorphisms in a core collection of tomato (*Solanum lycopersicum* L.). *Scientia Horticulturae*, 243, 567-574.
 49. Rikkerink, E. H., Oraguzie, N. C., & Gardiner, S. E. (2007). Prospects of association mapping in perennial horticultural crops. In *Association mapping in plants* (pp. 249-269). Springer, New York, NY.
 50. Ro, N., Haile, M., Hur, O., Geum, B., Rhee, J., Hwang, A., Kim, B., Lee, J., Hahn, B-S., Lee, J., & Kang, B. C. (2022). Genome-Wide Association Study of Resistance to Phytophthoracapsici in the Pepper (*Capsicum* spp.) Collection. *Frontiers in Plant Science*, 13:902464. doi: 10.3389/fpls.2022.902464
 51. Solomon, A. M., Han, K., Lee, J. H., Lee, H. Y., Jang, S., & Kang, B. C. (2019). Genetic diversity and population structure of Ethiopian *Capsicum* germplasms. *PloS one*, 14(5), e0216886.
 52. Sukumaran, S., & Yu, J. (2014). Association mapping of genetic resources: achievements and future perspectives. In *Genomics of plant genetic resources* (pp. 207-235). Springer, Dordrecht.
 53. Toppino, L., Barchi, L., Mercati, F., Acciarri, N., Perrone, D., Martina, M., & Rotino, G. L. (2020). A new intra-specific and high-resolution genetic map of eggplant based on a RIL population, and location of QTLs related to Plant anthocyanin pigmentation and seed vigour. *Genes*, 11(7), 745.
 54. Tripodi, P., Soler, S., Campanelli, G., Díez, M. J., Esposito, S., Sestili, S., & Cardi, T. (2021). Genome wide association mapping for agronomic, fruit quality, and root architectural traits in tomato under organic farming conditions. *BMC plant biology*, 21(1), 1-22.
 55. Tuberosa R, Frison E, Graner A (2014) Genomics of plant genetic resources. *Genom Plant Genet Resour Springer* 11:1–11.
 56. Wang, F., Zou, M., Zhao, L., Li, H., Xia, Z., & Wang, J. (2020). Genome-Wide Association Analysis of Late Blight Resistance Traits in Potato Germplasm Resources.
 57. Wang, T., Zhang, Z., Zhu, H., Zhang, Y., Gao, W., Wang, X., & Zou, Q. (2020). Phenotypic diversity and genome-wide association mapping of earliness-related traits in cultivated tomato (*Solanum Lycopersicum* L.). *Scientia Horticulturae*, 264, 109194.
 58. Wu, L., Wang, H., Liu, S., Liu, M., Liu, J., Wang, Y., ...& Shen, H. (2022). Mapping of CaPP2C35 involved in the formation of light-green immature pepper (*Capsicum annum* L.) fruits via GWAS and BSA. *Theoretical and Applied Genetics*, 135(2), 591-604.
 59. Wu, L., Wang, P., Wang, Y., Cheng, Q., Lu, Q., Liu, J., ...& Shen, H. (2019). Genome-wide correlation of 36 agronomic traits in the 287 pepper (*Capsicum*) accessions obtained from the SLAF-seq-based GWAS. *International journal of molecular sciences*, 20(22), 5675.
 60. Yamamoto, T., & Terakami, S. (2016). Genomics of pear and other Rosaceae fruit trees. *Breeding science*, 66(1), 148-159.
 61. Ye, J., Wang, X., Hu, T., Zhang, F., Wang, B., Li, C., & Ye, Z. (2017). An InDel in the promoter of Al-ACTIVATED MALATE TRANSPORTER9 selected during tomato domestication determines fruit malate contents and aluminum tolerance. *The Plant Cell*, 29(9), 2249-2268.

62. Yuan, J., Bizimungu, B., De Koeyer, D., Rosyara, U., Wen, Z., & Lagüe, M. (2020). Genome-wide association study of resistance to potato common scab. *Potato Research*, 63(2), 253-266.
63. Zahid, G., Aka Kaçar, Y., Dönmez, D., Küden, A., & Giordani, T. (2022). Perspectives and recent progress of genome-wide association studies (GWAS) in fruits. *Molecular Biology Reports*, 49(6), 5341-5352.
64. Zhu, C., Gore, M., Buckler, E. S., & Yu, J. (2008). Status and prospects of association mapping in plants. *The plant genome*, 1(1).
65. Zia, M. A. B., Demirel, U., Nadeem, M. A., Ali, F., Dawood, A., Ijaz, M., & Caliskan, M. E. (2022). Genome-wide association studies (GWAS) revealed a genetic basis associated with floral traits in potato germplasm. *Turkish Journal of Agriculture and Forestry*, 46(1), 90-103.
66. Devi R, Chauhan S and Dhillon TS (2022) Genome editing for vegetable crop improvement: Challenges and future prospects. *Front. Genet.* 13:1037091. doi: 10.3389/fgene.2022.1037091

UNDER PEER REVIEW