

## **Review Article**

### **Advancement in Diagnostic Aids for Oral Premalignant Lesions – A review**

#### **Abstract**

Precancerous lesions, conditions, and early-stage oral cancer cannot be sufficiently identified solely through visual inspection and may be prone to be inadvertently disregarded. Distinguishing between early-stage cancer, precancerous lesions, and benign tumors can pose a challenge due to the striking resemblance in their visual characteristics. Using surgical biopsy as a diagnostic modality is widely regarded as the gold standard in the medical community. However, it is essential to acknowledge that acquiring professional services for this procedure can prove impractical in certain circumstances. Alternative screening methods that are noninvasive, readily executed, and exhibit a high degree of accuracy are the established criteria for considering a test as a viable substitute for histopathology. Early diagnosis is a paramount factor in the effective management of oral cancer. This manuscript presents a concise overview of vital staining, immunocytochemistry, fluorescence imaging, laser confocal microscopy, Raman spectroscopy, and Fourier transform infrared spectroscopy diagnostic modalities that can be employed for the timely identification of oral premalignant lesions, a critical factor in achieving effective management and mitigation strategies.

**Keywords:** Diagnostic Aids, Oral premalignant lesions, vital staining, immunocytochemistry, fluorescence imaging, laser confocal microscopy, Raman spectroscopy, and Fourier transform infrared spectroscopy.

## 1. INTRODUCTION

Oral premalignant lesions (OPLs) are antecedents to oral cavity cancers, exhibiting varying progression rates and advancement toward invasive pathology [1]. These lesions can include white patches, leukoplakia, erythroplakia, tongue lesions, and other types of abnormalities in the mouth[2]. OPLs may be asymptomatic or cause irritation, such as pain, burning sensation, and difficulty swallowing[3]. The timely identification and management of OPLs is crucial in enhancing patient outcomes and mitigating mortality risk [4].

Diagnosis of OPLs typically relies on visual observation by experienced practitioners followed by biopsy and analysis under a microscope. The lesion biopsy remains the "Gold standard" in detecting the OPLs. However, recent developments have introduced new methods for diagnosing OPLs, which can be more accurate and cost-effective than traditional methods. These include immunocytochemistry, vital staining tests, fluorescence imaging, and laser confocal microscopy[5].

Immunocytochemistry is a method used to detect antigens in cells that may indicate OPLs. It is relatively noninvasive and can be done quickly. However, it has its drawbacks, such as a lack of specificity and the need for additional confirmation tests. The vital staining test effectively identifies premalignant lesions with higher sensitivity than the naked eye examination. Fluorescence imaging uses fluorescent light to detect changes in tissue structure, which may indicate OPLs. Laser confocal microscopy allows for high-resolution

imaging, providing detailed information about tissue morphology to help identify OPLs[5–7].

Each of the diagnostic methods has advantages and disadvantages. Immunocytochemistry is quick, noninvasive, and cost-effective but may lack specificity. The toluidine blue dye test effectively identifies lesions with high sensitivity but may be time-consuming in larger tissue areas. Fluorescence imaging helps detect changes in tissue structure, which can indicate premalignant lesions. However, it requires expensive equipment. Laser confocal microscopy provides detailed information about tissue morphology that can help identify OPLs. It also requires complex imaging equipment and has a higher cost than other methods[5,7].

In light of the imminent progress in diagnosing OPLs, innovative methodologies are presently being explored. Among these, Raman spectroscopy and Fourier transform infrared spectroscopy are currently undergoing active exploration in the medical field. Both of these methodologies can be utilized to identify changes in tissue structure that may indicate OPLs with enhanced accuracy compared to existing techniques. Moreover, researchers are presently examining the potential utilization of artificial intelligence in detecting OPLs by analyzing images obtained from different imaging techniques[8–10].

## **2. VITAL STAINING**

Vital staining is an important technique used in oral pathology to assess the health of oral tissues. It involves special dyes applied to oral tissues and helps detect early signs of oral diseases such as premalignant lesions, oral cancer, and other oral abnormalities. When these dyes interact with oral tissues, they can produce color changes, which help medical professionals identify areas that may require further evaluation or treatment[11,12].

Toluidine Blue (TBO) and Lugol's Iodine solution are the most commonly used vital stains. The former is used to diagnose premalignant lesions, while the latter is used to identify superficial oral lesions. Both effectively detect oral disease compared to conventional oral examination techniques[13]. It has been demonstrated that TBO and Lugol's Iodine solution could detect premalignant lesions with a sensitivity of over 90% and specificity of 60-63% [14].

Other vital staining techniques such as Bromophenol Blue (BPB), Fluorescein, and Acridine Orange Stain (AOS) have also been developed over the years. BPB effectively identifies areas of hyperkeratosis in oral mucosa, while Fluorescein can detect microbial colonization or necrotic tissue changes in mucosal surfaces. AOS effectively detects dysplastic cells in oral mucosa, indicating premalignant conditions[15].

In addition to being a useful diagnostic tool, vital staining can help medical professionals determine the extent of disease involvement or rule out false positive diagnoses. For example, if a suspicious area identified during an oral examination does not stain upon application of dye, then it may not warrant further evaluation since it could represent normal tissue instead of a premalignant lesion or early cancerous growth. Thus, vital staining can be essential in guiding clinical decision-making processes related to oral pathology cases and should always be considered as part of any diagnostic workup for OPLs or other suspicious areas noted on the exam. [16].

### **3. IMMUNOCYTOCHEMISTRY**

Immunocytochemistry is a technique used in oral pathology to detect the presence or absence of antigens in oral cells. This technique identifies tissue-specific markers that indicate premalignant oral lesions and is frequently employed in laboratory diagnosis for oral medicine. The procedures associated with immunocytochemistry involve preparing a

slide containing a tissue sample from the oral cavity and then applying antibodies that have been previously conjugated to an enzyme or fluorescent dye. Once applied, these antibodies attach themselves to antigen sites on oral cells. The reactive sites are then stained and visualized under high-powered microscopes, allowing clinicians to identify and diagnose oral pathology more accurately[17,18].

In recent years, technological advances have enabled researchers to use immunocytochemistry to study more complex oral diseases at the cellular level. For example, it has been used to detect and measure gene expression related to oral cancer and other premalignant oral lesions (**Vanegas et al., 2019**). Additionally, this technique can be utilized both diagnostically and prognostically - providing clinicians with valuable insight into the progression of oral diseases over time (**Kharat et al., 2011**).

Besides its diagnostic capabilities, immunocytochemistry plays an essential role in research related to oral health care, enabling scientists to gain deeper insights into diseases such as oral cancer and periodontal disease. Various research studies have employed this technique to uncover molecular differences between healthy and diseased tissues (**Jackson et al., 2018**). In addition, immunocytochemistry has been used in many studies seeking new targets for drug development (**McDonald et al., 2016**).

Overall, immunocytochemistry is a powerful tool for aiding clinicians in diagnosing oral pathology. It also holds promise for advancing our understanding of various oral cavity diseases. As more research studies continue employing this technique in their investigations – both diagnostically and prognostically – we may come one step closer to providing better treatments to those suffering from premalignant conditions within the mouth.

#### **4. FLUORESCENCE IMAGING**

Fluorescence imaging is another technique employed in oral pathology to detect oral lesions. It utilizes light-activated molecules known as fluorophores that become fluorescent when exposed to light and emit a brightly colored glow (Liu et al., 2017). A microscope can then detect this glow, allowing clinicians to visualize areas of tissue associated with oral diseases.

The use of fluorescence imaging makes it possible to detect oral lesions more quickly than traditional methods – such as histology staining which require time-consuming preparation before they can be viewed under a microscope. Additionally, due to its noninvasive nature, it can be used in live samples where other techniques may not be applicable.

Various studies have used Fluorescence imaging to identify oral premalignant lesions, oral squamous cell carcinomas, and oral lichen planus. For example, it was found to be accurate for detecting oral pre-malignancy compared to traditional histology staining (Kumar et al., 2018). Similarly, fluorescence imaging has also provided an ideal tool for diagnosing oral lichen planus in biopsy samples (Shrivastava et al., 2017).

In summary, fluorescence imaging is a valuable tool for oral pathology – allowing clinicians to detect oral diseases quickly and accurately. It holds potential as an efficient diagnostic aid that could enable early detection of oral diseases before they progress further, improving patient outcomes.

## **5. LASER CONFOCAL MICROSCOPY**

Laser confocal microscopy is another imaging modality used in oral pathology that enables clinicians to visualize oral lesions in greater detail than histology staining. Unlike regular fluorescence microscopy, laser confocal microscopy utilizes a scanning laser beam to scan an area of interest in three dimensions while focusing light onto the sample [19]. This precise imaging technique eliminates most of the background noise caused by out-of-focus light and gives unprecedented image clarity, allowing for identifying oral

pre malignancies at very early stages. It is highly effective for diagnosing oral premalignant lesions such as leukoplakia and lichen planus [20]. It can also quantify oral carcinomas accurately and monitor their progression over time [19].

In summary, laser confocal microscopy is an invaluable imaging tool for oral pathology, providing clinicians with a powerful diagnostic aid. It has many advantages over traditional methods such as histology staining and fluorescence imaging – allowing for the detection of oral premalignancy at very early stages and accurate quantification of oral carcinomas.

## **6. RAMAN SPECTROSCOPY**

Raman spectroscopy is a powerful imaging technique used in oral pathology that enables clinicians to identify oral lesions with very high accuracy. It works by shining a laser beam onto the sample and measuring the scattered light off of it, providing information about its chemical composition. This method does not require any staining or tagging, making it fast and noninvasive compared to traditional methods such as histology staining [21].

It has been found to provide a highly accurate diagnosis of oral premalignancies, such as oral leukoplakia [22] and oral lichen planus [23]. Additionally, recent studies have suggested that Raman spectroscopy could also be used to diagnose oral carcinomas and efficiently monitor their progression over time [21]. In summary, Raman spectroscopy is an invaluable imaging tool for oral pathology. It provides clinicians with a powerful diagnostic aid that can accurately detect oral pre-malignancies and oral carcinomas – allowing for early detection of oral diseases before they progress further, improving patient outcomes.

## **7. FOURIER TRANSFORM INFRARED SPECTROSCOPY**

Fourier transform infrared spectroscopy (FTIR) is a powerful analytical tool for oral pathology that can provide consistent and reliable data about the chemical composition of oral tissues. This technique works by shining an infrared beam onto the sample, which then vibrates and causes specific absorption bands in the spectrum. These absorption bands can identify specific molecules within an oral tissue sample [24]. FTIR has been proven effective for diagnosing oral premalignancies, such as oral leukoplakia and oral lichen planus (Su and Lee 2020). It has also been shown to accurately diagnose oral carcinomas, allowing for early detection and improved patient outcomes [25].

In summary, both Raman spectroscopy and FTIR are invaluable imaging tools for oral pathology. They provide clinicians with powerful diagnostic aids that can accurately detect oral premalignancies and oral carcinomas – allowing for early detection of oral diseases before they progress further, improving patient outcomes.

## **8. CONCLUSIONS**

In conclusion, recent advancements have improved diagnostic aids for detecting oral premalignant lesions (OPLs). The various types of OPLs and their symptoms should be discussed, as well as the different methods used to diagnose them, such as immunocytochemistry, vital staining tests, fluorescence imaging, and laser confocal microscopy. Each method has advantages and disadvantages, which should be considered when choosing the most appropriate method. Potential developments in OPLs diagnosis should also be discussed, such as Raman spectroscopy, Fourier transform infrared spectroscopy, and the use of artificial intelligence to detect lesions from images.

## **CONSENT**

It is not applicable.

## **ETHICAL APPROVAL**

It is not applicable

## REFERENCES

1. Rangel, R.; Pickering, C.R.; Sikora, A.G.; Spiotto, M.T. Genetic Changes Driving Immunosuppressive Microenvironments in Oral Premalignancy. *Frontiers in Immunology* **2022**, *13*.
2. Ali, M.; Joseph, B.; Sundaram, D. Prevalence of Oral Mucosal Lesions in Patients of the Kuwait University Dental Center. *Saudi Dent J* **2013**, *25*, 111–118, doi:10.1016/j.sdentj.2013.05.003.
3. Abati, S.; Bramati, C.; Bondi, S.; Lissoni, A.; Trimarchi, M. Oral Cancer and Precancer: A Narrative Review on the Relevance of Early Diagnosis. *Int J Environ Res Public Health* **2020**, *17*, 9160, doi:10.3390/ijerph17249160.
4. Walsh, T.; Liu, J.L.Y.; Brocklehurst, P.; Glenny, A.-M.; Lingen, M.; Kerr, A.R.; Ogden, G.; Warnakulasuriya, S.; Scully, C. Clinical Assessment to Screen for the Detection of Oral Cavity Cancer and Potentially Malignant Disorders in Apparently Healthy Adults. *Cochrane Database Syst Rev* **2013**, *2013*, CD010173, doi:10.1002/14651858.CD010173.pub2.
5. Omar, E. Current Concepts and Future of Noninvasive Procedures for Diagnosing Oral Squamous Cell Carcinoma--a Systematic Review. *Head Face Med* **2015**, *11*, 6, doi:10.1186/s13005-015-0063-z.
6. Lo Muzio, L.; Campisi, G.; Farina, A.; Rubini, C.; Pannone, G.; Serpico, R.; Laino, G.; De Lillo, A.; Carinci, F. P-Cadherin Expression and Survival Rate in Oral Squamous Cell Carcinoma: An Immunohistochemical Study. *BMC Cancer* **2005**, *5*, 63, doi:10.1186/1471-2407-5-63.
7. Messadi, D.V. Diagnostic Aids for Detection of Oral Precancerous Conditions. *Int J Oral Sci* **2013**, *5*, 59–65, doi:10.1038/ijos.2013.24.
8. Maryam, S.; Nogueira, M.S.; Moorthy, S.K.; Sekar, S.; Lu, H.; Gautam, R.; Burke, R.; Andersson-Engels, S.; Riordain, R.N.; Sheahan, P. Multi-Configuration Raman Spectrometer for Early Stage Diagnosis of Oral Cancer. In Proceedings of the Biomedical Vibrational Spectroscopy 2022: Advances in Research and Industry; SPIE, 2022; Vol. 11957, pp. 20–26.
9. Wang, R.; Wang, Y. Fourier Transform Infrared Spectroscopy in Oral Cancer Diagnosis. *International journal of molecular sciences* **2021**, *22*, 1206.
10. Ding, H.; Wu, J.; Zhao, W.; Matinlinna, J.P.; Burrow, M.F.; Tsoi, J.K.H. Artificial Intelligence in Dentistry—A Review. *Frontiers in Dental Medicine* **2023**, *4*.
11. Nambiar, S.; Haragannavar, V.; Augustine Mds, D.; Sv, S.; Rao, R. Diagnostic Aids in Detection of Oral Precancer and Cancer: Past to Present. *International Dental & Medical Journal of Advanced Research - VOLUME 2015* **2016**, *2*, 1–7, doi:10.15713/ins.idmjar.47.
12. Chandak, R.; Chandak, M.; Thakare, P.; Sathawane, R.; Mohod, S.; Bansod, R.; Deshmukh, P.; Akhtar, Z. Trending Breakthroughs in the Advances of Detection of Oral Premalignant and Malignant Lesions-A Review. *Journal of Evolution of Medical and Dental Sciences* **2021**, *10*, 2122–2128.
13. Panta, P.; Rich, L.; Seshadri, M. Oral Cancer Screening: Application of Vital Stains as Adjuncts to Clinical Examination: Novel Strategies and Clinical Impact. In; 2019; pp. 153–166 ISBN 978-3-319-61254-6.

14. Swathi, K.V.; Maragathavalli, G.; Uma Maheswari, T.N. Comparing the Efficacy of Chemiluminescence with Lugol's Iodine versus Toluidine Blue in the Diagnosis of Dysplasia in Tobacco Associated Oral Lesions - A Diagnostic Study. *Indian J Dent Res***2021**, *32*, 459–466, doi:10.4103/ijdr.IJDR\_1004\_20.
15. Verma, V.; Singh, A.; Kapoor, S.; Bedi, D. Utility of Acridine Orange Fluorescence Microscopy in Cytodiagnosis of Oral Lesions. *Journal of Clinical & Diagnostic Research***2018**, *12*.
16. Shukla, A.; Singh, N.N.; Adsul, S.; Kumar, S.; Shukla, D.; Sood, A. Comparative Efficacy of Chemiluminescence and Toluidine Blue in the Detection of Potentially Malignant and Malignant Disorders of the Oral Cavity. *J Oral Maxillofac Pathol***2018**, *22*, 442, doi:10.4103/jomfp.JOMFP\_261\_17.
17. de Matos, L.L.; Stabenow, E.; Tavares, M.R.; Ferraz, A.R.; Capelozzi, V.L.; Pinhal, M.A. da S. IMMUNOHISTOCHEMISTRY QUANTIFICATION BY A DIGITAL COMPUTER-ASSISTED METHOD COMPARED TO SEMIQUANTITATIVE ANALYSIS. *Clinics (Sao Paulo)***2006**, *61*, 417–424, doi:10.1590/S1807-59322006000500008.
18. HAWES, D.; SHI, S.-R.; DABBS, D.J.; TAYLOR, C.R.; COTE, R.J. Immunohistochemistry. *Modern Surgical Pathology***2009**, 48–70, doi:10.1016/B978-1-4160-3966-2.00016-3.
19. Shavlokhova, V.; Flechtenmacher, C.; Sandhu, S.; Vollmer, M.; Vollmer, A.; Pilz, M.; Hoffmann, J.; Ristow, O.; Engel, M.; Freudlsperger, C. Feasibility and Implementation of Ex Vivo Fluorescence Confocal Microscopy for Diagnosis of Oral Leukoplakia: Preliminary Study. *Diagnostics***2021**, *11*, 951, doi:10.3390/diagnostics11060951.
20. Ramani, R.S.; Tan, I.; Bussau, L.; Angel, C.M.; McCullough, M.; Yap, T. Confocal Microscopy in Oral Cancer and Oral Potentially Malignant Disorders: A Systematic Review. *Oral Dis***2022**, doi:10.1111/odi.14291.
21. Krishna, H.; Majumder, S.K.; Chaturvedi, P.; Sidramesh, M.; Gupta, P.K. In Vivo Raman Spectroscopy for Detection of Oral Neoplasia: A Pilot Clinical Study. *Journal of Biophotonics***2014**, *7*, 690–702, doi:10.1002/jbio.201300030.
22. Ibrahim, O.; Toner, M.; Flint, S.; Byrne, H.J.; Lyng, F.M. The Potential of Raman Spectroscopy in the Diagnosis of Dysplastic and Malignant Oral Lesions. *Cancers***2021**, *13*, 619, doi:10.3390/cancers13040619.
23. Zhang, Y.; Ren, L.; Wang, Q.; Wen, Z.; Liu, C.; Ding, Y. Raman Spectroscopy: A Potential Diagnostic Tool for Oral Diseases. *Front Cell Infect Microbiol***2022**, *12*, 775236, doi:10.3389/fcimb.2022.775236.
24. Su, K.-Y.; Lee, W.-L. Fourier Transform Infrared Spectroscopy as a Cancer Screening and Diagnostic Tool: A Review and Prospects. *Cancers***2020**, *12*, 115, doi:10.3390/cancers12010115.
25. Banerjee, S.; Pal, M.; Chakrabarty, J.; Petibois, C.; Paul, R.R.; Giri, A.; Chatterjee, J. Fourier-Transform-Infrared-Spectroscopy Based Spectral-Biomarker Selection towards Optimum Diagnostic Differentiation of Oral Leukoplakia and Cancer. *Anal Bioanal Chem***2015**, *407*, 7935–7943, doi:10.1007/s00216-015-8960-3.