

Morphological and biochemical and quality control analysis of peas genotypes under cadmium toxicity at the seedling stage

ABSTRACT

Cadmium is the most dangerous heavy metal because of its carcinogenic nature and no function in the body of living beings.

Cadmium toxicity is an emerging problem as it is the most abundant heavy metal in Pakistani soils. Peas are being grown near cities where an elevated level of cadmium persists. So, this study was conducted for the evaluation of 10 peas genotypes (TRIPLET, MID No. 2, Oskar, LM (OSE), METEOR, P-1, Rohina, PEA-09, LM-POIS, and LAMPO) under cadmium stress at the seedling stage in the greenhouse of Department of Plant Breeding and Genetics, University of Agriculture, Faisalabad. There were 3 treatments of cadmium (0, 4, and 8 mg/kg soil) and each treatment was replicated five times in a completely randomized design. Different morphological traits i.e., fresh root weight, fresh seedling weight, root length, seedling length, dry root weight, dry seedling weight and biochemical traits i.e., Cd in the root, shoot, and soil were recorded by atomic absorption spectrometry. Generally, T1 reduced root and shoot parameters while 8 ppm Cd application comparatively enhanced the growth of peas genotypes. LM-POIS and LM (OSE) genotypes performed well under Cd stress and LM-POIS had also an elevated level of Cd in the shoot on the other hand TRIPLET and LAMPO were the least performing genotypes. It is suggested that Cd contamination significantly reduces pea growth; therefore, Cd-tolerant-peas genotypes should be developed or introduced. Further, Cd exposure in the soil should be minimized and already Cd-polluted soils should be remediated.

Keywords: Cd toxicity; Peas genotypes; Quality control analysis; Soil pollution, Pre-breeding evaluation

1. INTRODUCTION

Peas are an important crop in Pakistan, with a significant economic impact on farmers. It is the most prevalent crop and, owing to its nutritious value, has a high commercial demand. It is grown on the plains during the winter and in the highlands during the summer (1). The major pea-growing districts in Punjab include Gujranwala, Nankana Sahib, Multan, Sahiwal, Toba Tek Singh, Sialkot, Jhang, and Sheikupura (2). Peas are mostly grown in Punjab during the months of October and November (3). The lack of resistant and high-yielding cultivars, as well as the use of inappropriate peas-growing methods, has resulted in a reduction in the per-acre average yield in Pakistan (4).

Biotic and abiotic stresses have hindered pea production in Pakistan, resulting in decreased yields (5). Abiotic stresses are becoming more severe because of climate change due to global warming (6). Abiotic stress includes drought, salt stress, cold stress, heat stress, and heavy metals in the soil (7). Heavy metals are accumulated in the soil when crops are irrigated with untreated sewage or industrial water (8). Cd is the most dangerous heavy metal for all species due to its carcinogenic properties (9). Cd polluted the vegetables grown near cities and irrigated with untreated industrial sewage wastewater (10).

Despite having any metabolic function in higher plants, cadmium is a highly toxic and nonessential heavy metal (11). It has the potential to go into the food chain and harm the biota (12). Toxic concentrations of Cd in plants obstruct plant growth, and its production and disrupt seedling physiological systems that ultimately decrease crop output in peas (13). The impact of Cd on peas and cowpea seeds was related to a lack of water absorption, limiting the amount of water available for seed embryo development (14).

Cd stress significantly reduces the growth and development of bean seedlings (15). Different peas genotypes have been evaluated under Cd toxicity to evaluate Cd-resistant genotypes (16). Another effort was made to evaluate peas genotypes (Hurst Green and Kelvedon Wonder) against Cd toxicity. It was determined that Kelvedon Wonder was tolerant to Cd toxicity (17). Generally, it has been discovered that peas are significantly affected by Cd toxicity hence, their growth reduce remarkably after Cd exposure at germination and seedling stage (18).

Keeping in view above scenario the current study aimed at the evaluation of cultivars and breeding lines under cadmium toxicity, at seedling stage in screenhouse. Determination of cadmium concentration helped to identify genotypes with better performance in cadmium toxicity.

2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

2.1 Experimental Site and Treatments

This experiment was conducted in the greenhouse of the Department of Plant Breeding and Genetics, University of Agriculture, Faisalabad. The experiment was carried out in a completely randomized design with five replicates in winter 2020-21. 500 ml plastic jars were filled with 500 mg sand to sow pea genotypes. Ten peas genotypes (TRIPLET, MID No. 2, Oskar, LM (OSE), METEOR, P-1, Rohina, PEA-09, LM-POIS, and LAMPO) were treated with 3 levels of Cadmium Nitrate ($\text{Cd}(\text{NO}_3)_2$) in solution form after 15 days of sowing:

- 1) 0 mg Cd/kg Soil (T0) (Control)
- 2) 4 mg Cd/kg Soil (T1)
- 3) 8 mg Cd/kg Soil (T2)

2.2 Morphological Analysis

The experiment was carried out for 30 days. Five seedlings from each genotype per treatment per replication were selected and the mean was calculated. Data was recorded after two weeks of Cd application. Data were recorded for the following morphological traits: seed length (cm), root length (cm), fresh shoot weight (g), fresh root weight (g), dry shoot weight (g), and dry root weight (g).

Comment [2A1]: Reconsider the spelling.

2.3 Biochemical Analysis

For this purpose, 0.25g of oven-dried samples were added into 5ml HNO₃ and HClO₄ solution (3:1). Prepared material was kept in the chamber overnight and then heated on a hot plate (350°C) until the solution volume remained 1ml. Now, the solution turned from dark yellow color to colorless. The Solution was poured out into a new flask having 50ml volume by filtering the solution with filter paper (What's man No.1). 50ml double distilled water was added to the samples which were ready to be used for Cd analysis in Atomic Absorption Spectrometer.

2.4 Statistical Analysis

Collected data were subjected to analysis of variance and Tuckey HSD Test for evaluation of significant differences among treatments (19). Analysis of variance and Tukey HSD test were performed by using Statistix 8.1 software. Bio-absorption co-efficient (BAC), Bio-concentration Factor (BCF), and Translocation Factor (TF) were calculated by following formulas given by (20).

$$\text{BAC} = (\text{Cd in shoot})/(\text{Cd in soil})$$

$$\text{BCF} = (\text{Cd in root})/(\text{Cd in soil})$$

$$\text{TF} = (\text{Cd in shoot})/(\text{Cd in root})$$

3. RESULTS AND DISCUSSION

Statistical analysis revealed that there were substantial variations among varieties and treatments. An increased level of cadmium was observed in root and seedlings by increasing the level of Cd in the soil. All genotypes responded differently to Cd application for all measured morphological and biochemical attributes.

3.1 Morphological Analysis

Generally, the fresh root weight of all genotypes was decreased by the application of Cd at 4 ppm, except TRIPLET and MID No. 2. The maximum root weight was recorded in LM (OSE) (2.66 g) when grown under Cd stress at 8 ppm. While METEOR had the lowest root weight (0.40 g) under Cd stress at 4 ppm. When Cd stress was increased up to 8 ppm the fresh root weight of almost all genotypes was increased compared to 4 ppm Cd stress. The root length of LM-POIS (27.66 cm) was maximum in Cd stress at 8 ppm, among the root lengths of all genotypes in all treatments. Cd stress at 4 ppm significantly reduced the root length of peas genotypes while Cd stress at 8 ppm comparatively increased the root length. The lowest root length was recorded in LAMPO (7.5 cm) and METEOR (7.67 cm) under 4 ppm Cd stress. There was a significant reduction in the dry root weight of all genotypes by Cd application but a great improvement in the dry root weight of Oskar and LM (OSE) in Cd stress at 8 ppm. LM-POIS (0.93g) had maximum dry root weight and the lowest dry root weight was noted in METEOR (0.12 g). There was a significant difference in fresh seedling weight of all genotypes in all three treatments. The maximum fresh seedling weight was recorded in P-1

(3.13 g) in control, on the other hand, the least fresh seedling weight was measured in LAMPO (0.65 cm) in 4 ppm Cd stress. There was also variation in fresh seedling weight in response to Cd stress at 8 ppm for all genotypes. The seedling length of all genotypes was greatly reduced by Cd application as compared to the control treatment. The seedling length of Oskar (42.67 cm) was maximum in the control condition than all genotypes in all treatments while the lowest seedling length was measured in LAMPO (8.25 cm) in response to Cd stress at 4 ppm. Dry seedling weight was generally reduced under Cd stress, but 4 ppm application of Cd had more severe effects on dry seedling weight of all genotypes. Maximum dry seedling weight was measured in P-1 (0.91 g) in control conditions than dry seedling weights of all genotypes in all treatments and LAMPO (0.8 g) had the lowest dry seedling weight. Results are summarized in Table 1-6 subjected under supplementary material.

Table 1: LSD all pairwise comparisons for fresh root weight of different peas genotypes grown under different Cd levels

| Genotypes | Control | Cd @ 4 ppm | Cd @ 8 ppm | Mean |
|-----------|-----------|------------|------------|----------|
| TRIPLET | 0.750 kl | 0.966 g-j | 1.150 e-g | 0.955 C |
| MID No. 2 | 1.266 de | 1.866 c | 1.866 c | 1.666 A |
| Oskar | 1.166 ef | 0.900 i-k | 2.400 b | 1.488 B |
| LM (OSE) | 1.066 f-i | 0.966 g-j | 2.666 a | 1.566 AB |
| METEOR | 1.133 e-h | 0.400 m | 1.300 de | 0.944 CD |
| P-1 | 1.133 e-h | 0.700 l | 0.966 g-j | 0.933 CD |
| Rohina | 2.333 b | 1.066 f-i | 1.366 d | 1.588 AB |
| PEA-09 | 0.966 g-j | 0.633 l | 0.900 i-k | 0.833 D |
| LM-POIS | 1.800 c | 0.966 g-j | 1.900 c | 1.555 AB |
| LAMPO | 0.966 g-j | 0.800 j-l | 0.950 h-j | 0.905 CD |
| Mean | 1.258 B | 0.926 C | 1.546 | |

Comment [2A2]: The first row and column shall be bold for all tables.

Comment [2A3]: Kl, de, ef, e-h,... etc shall be defined either as legend or as proper sentences for all tables.

Table 2: LSD all pairwise comparisons for dry root weight of different peas genotypes grown under different Cd levels

| Genotypes | Control | Cd @ 4 ppm | Cd @ 8 ppm | Mean |
|-----------|----------|------------|------------|----------|
| TRIPLET | 0.230 lm | 0.133 n | 0.260 kl | 0.207 G |
| MID No. 2 | 0.476 g | 0.613 e | 0.713 d | 0.601 A |
| Oskar | 0.373 h | 0.283 jk | 0.816 c | 0.491 D |
| LM (OSE) | 0.260 kl | 0.300 jk | 0.883 ab | 0.481 D |
| METEOR | 0.280 jk | 0.123 n | 0.370 h | 0.257 F |
| P-1 | 0.316 ij | 0.206 m | 0.360 hi | 0.294 E |
| Rohina | 0.870 b | 0.256 kl | 0.446 g | 0.524 C |
| PEA-09 | 0.310 j | 0.230 lm | 0.230 lm | 0.256 F |
| LM-POIS | 0.926 a | 0.223 lm | 0.563 f | 0.571 B |
| LAMPO | 0.303 jk | 0.200 m | 0.193 m | 0.232 FG |
| Mean | 0.434 B | 0.257 C | 0.483 A | |

Table 3: LSD all pairwise comparisons for root length of different peas genotypes grown under different Cd levels

| Genotypes | Control | Cd @ 4 ppm | Cd @ 8 ppm | Mean |
|-----------|------------|------------|------------|-----------|
| TRIPLET | 16.500 hi | 16.333 hi | 14.000 jk | 15.611 C |
| MID No. 2 | 14.333 j | 12.667 lm | 18.333 ef | 15.111 CD |
| Oskar | 17.333 f-h | 12.333 m | 17.333 f-h | 15.667 C |
| LM (OSE) | 16.667 hi | 12.000 mn | 24.667 b | 17.778 B |
| METEOR | 13.667 j-l | 7.667 q | 13.000 k-m | 11.444 F |
| P-1 | 16.667 hi | 10.333 o | 16.667 hi | 14.556 D |
| Rohina | 18.000 e-g | 12.667 lm | 15.667 i | 15.444 C |
| PEA-09 | 17.000 gh | 9.000 p | 11.000 no | 12.333 E |
| LM-POIS | 19.667 cd | 19.000 de | 27.667 a | 22.111 A |
| LAMPO | 20.333 c | 7.500 q | 11.000 no | 12.944 E |
| Mean | 17.017 A | 11.950 B | 16.933 A | |

Table 4: LSD all pairwise comparisons for the fresh seedling weight of different peas genotypes grown under different Cd levels

| Genotypes | Control | Cd @ 4 ppm | Cd @ 8 ppm | Mean |
|-----------|-----------|------------|------------|----------|
| TRIPLET | 0.933 mn | 1.200 jk | 0.866 n | 1.000 F |
| MID No. 2 | 2.033 cd | 1.200 jk | 1.433 h | 1.555 D |
| Oskar | 1.366 h-j | 0.966 l-n | 1.133 kl | 1.155 E |
| LM (OSE) | 2.300 b | 1.166 k | 1.666 fg | 1.711 C |
| METEOR | 2.333 b | 1.166 k | 1.433 h | 1.644 CD |
| P-1 | 3.133 a | 1.433 h | 1.533 gh | 2.033 A |
| Rohina | 2.200 bc | 1.766 f | 1.766 f | 1.911 B |
| PEA-09 | 2.166 bc | 1.233 i-k | 1.400 hi | 1.600 D |
| LM-POIS | 1.833 ef | 1.966 de | 0.900 n | 1.566 D |
| LAMPO | 1.166 k | 0.650 o | 1.100 k-m | 0.972 F |
| Mean | 1.946 A | 1.323 B | 1.275 B | |

Table 5: LSD all pairwise comparisons for dry seedling weight of different peas genotypes grown under different Cd levels

| Genotypes | Control | Cd @ 4 ppm | Cd @ 8 ppm | Mean |
|-----------|----------|------------|------------|----------|
| TRIPLET | 0.450 gh | 0.310 j-m | 0.240 m-o | 0.333 E |
| MID No. 2 | 0.676 cd | 0.276 k-n | 0.420 hi | 0.457 D |
| Oskar | 0.640 d | 0.276 k-n | 0.460 gh | 0.458 D |
| LM (OSE) | 0.686 cd | 0.230 mno | 0.506 fg | 0.474 CD |
| METEOR | 0.543 ef | 0.266 lmn | 0.470 fgh | 0.426 D |

| | | | | |
|---------|-----------|-----------|-----------|----------|
| P-1 | 0.913 a | 0.283 k-n | 0.366 ij | 0.521 BC |
| Rohina | 0.770 b | 0.640 d | 0.500 f-h | 0.636 A |
| PEA-09 | 0.726 bc | 0.350 i-k | 0.323 j-l | 0.466 D |
| LM-POIS | 0.623 de | 0.733 bc | 0.216 no | 0.524 B |
| LAMPO | 0.290 j-n | 0.180 o | 0.265 lmn | 0.245 F |
| Mean | 0.632 A | 0.354 B | 0.376 B | |

Table 6: LSD all pairwise comparisons for seedling length of different peas genotypes grown under different Cd levels

Comment [2A4]: Depict the tables in the form of graphs too. It will attract more views from audience.

| Genotypes | Control | Cd @ 4 ppm | Cd @ 8 ppm | Mean |
|-----------|-----------|------------|------------|-----------|
| TRIPLET | 29.500 c | 14.333 mn | 16.500 i-k | 20.111 E |
| MID No. 2 | 38.667 b | 17.333 h-j | 15.333 k-m | 23.778 B |
| Oskar | 42.667 a | 21.333 fg | 23.333 e | 29.111 A |
| LM (OSE) | 40.000 b | 13.667 no | 15.000 k-n | 22.889 BC |
| METEOR | 27.000 d | 17.667 hi | 15.333 k-m | 20.000 EF |
| P-1 | 28.333 cd | 20.667 g | 17.333 h-j | 22.111 CD |
| Rohina | 23.333 e | 18.667 h | 15.333 k-m | 19.111 F |
| PEA-09 | 23.000 e | 16.000 j-l | 14.667 l-n | 17.889 G |
| LM-POIS | 28.667 c | 12.667 op | 22.667 ef | 21.333 D |
| LAMPO | 11.667 p | 8.250 q | 12.500 op | 10.806 H |
| Mean | 29.283 A | 16.058 C | 16.800 B | |

3.2 Biochemical Analysis

For the mean comparison of genotypes, Cd concentration (5.90 ppm) in the root of Mid No. 2 was the highest while the lowest Cd concentration (4.25 ppm) was observed in the root of LM (OSE). By 4 ppm Cd application the lowest mean value (2.00 ppm) of Cd concentration was in the root of LM (OSE) whereas the highest Cd concentration was observed in the roots of Mid No. 2 (3.96) and LAMPO (3.83). TRIPLET (7.86 ppm) and Mid No. 2 (7.83) accumulated the highest Cd concentration under Cd stress at 8 ppm while the least Cd was observed in the roots of LM-POIS (5.23 ppm). Under Cd stress at 4 ppm Cd concentration in the shoot of LM (OSE) was maximum (4.00 ppm) while the least mean value was observed in Mid No. 2 (2.33 ppm). Under Cd stress at 8 ppm Cd concentration in shoot was maximum in LM-POIS (7.83 ppm). The lowest concentration was observed in Oskar (4.17 ppm) and P-1 (4.17 ppm). When Cd was applied at 4 ppm the average Cd concentration in soil samples was 3.77 ppm while those samples collected from soil treated with Cd at 8 ppm had 7.55 ppm. Results are summarized in Table 7-9 subjected under supplementary material.

Table 7: LSD all pairwise comparisons for Cd concentration in root of different peas genotypes grown under different levels of Cd stress

| Genotypes | Cd @ 4 ppm | Cd @ 8 ppm | Mean |
|-----------|------------|------------|---------|
| TRIPLET | 3.67 fg | 7.87 a | 5.76 AB |

| | | | |
|-----------|---------|----------|---------|
| MID No. 2 | 3.97 f | 7.83 a | 5.90 A |
| Oskar | 2.67 h | 7.70 a | 5.18 CD |
| LM (OSE) | 2.00 i | 6.50 cd | 4.25 E |
| METEOR | 3.47 g | 6.96 bc | 5.21 CD |
| P-1 | 3.53 fg | 7.66 a | 5.60 AB |
| Rohina | 3.56 fg | 5.37 e | 4.46 E |
| PEA-09 | 3.73 fg | 6.16 d | 4.95 D |
| LM-POIS | 3.36 g | 5.23 e | 4.30 E |
| LAMPO | 3.83 fg | 7.16 b | 5.50 BC |
| Mean | 3.38 B | 6.8467 A | |

Table 8: LSD all pairwise comparisons for Cd concentration in shoot of different peas genotypes grown under different levels of Cd toxicity

| Genotypes | Cd @ 4 ppm | Cd @ 8 ppm | Mean |
|-----------|------------|------------|---------|
| TRIPLET | 3.67 g | 4.23 ef | 3.95 DE |
| MID No. 2 | 2.33 h | 6.17 bc | 4.25 D |
| Oskar | 3.67 g | 4.16 ef | 3.91 E |
| LM (OSE) | 4.03 fg | 4.30 ef | 4.16 DE |
| METEOR | 3.67 g | 6.46 b | 5.06 B |
| P-1 | 3.63 g | 4.17 ef | 3.90 E |
| Rohina | 3.93 fg | 6.00 c | 4.97 B |
| PEA-09 | 3.93 fg | 4.53 e | 4.23 D |
| LM-POIS | 3.67 g | 7.83 a | 5.75 A |
| LAMPO | 3.86 fg | 5.43 d | 4.65 C |
| Mean | 3.64 B | 5.33 A | |

Table 9: LSD all-pairwise comparisons test of Cd concentration in soil

| Trt | Mean | Homogeneous Groups |
|-----|--------|--------------------|
| T1 | 3.7778 | B |
| T2 | 7.5556 | A |

3.3 Quality Control Analysis

The quality control analysis of peas genotypes was observed by calculating BAC, BCF, and TF. The BAC in LM (OSE) (1.07), Rohina (1.04), PEA-9 (1.04), and LAMPO (1.02) was increased than 1 under Cd stress at 4 ppm, and BAC of Cd in LM-POIS increased than 1 under Cd stress at 8 ppm. BCF for Cd in MID-No 2 (1.05) and LAMPO (1.01) was also elevated than 1 under 4 ppm Cd stress while BCF in TRIPLET (1.04), MID-No 2 (1.04), Oskar (1.02) P-1 (1.01) was the indicator of Cd uptake from soil to root in Cd stress at 8 ppm. TF under Cd stress at 4 ppm increased than 1 in almost genotypes except for MID-No.2. Surprisingly TF for Cd in LM(OSE) was 2.02, which was the highest value of TF for Cd in all genotypes under both treatments. Results are summarized in Table 10 subjected under supplementary material.

Table 10: Bio-absorption Coefficient, Bio-concentration factor and translocation factor for Cd in different peas genotypes grown under different levels of Cd stress

| Genotypes | Bio-absorption Coefficient | | Bio-concentration factor | | Translocation factor | |
|-----------|----------------------------|------|--------------------------|------|----------------------|------|
| | T1 | T2 | T1 | T2 | T1 | T2 |
| TRIPLET | 0.97 | 0.56 | 0.97 | 1.04 | 1.00 | 0.53 |
| MID No.2 | 0.61 | 0.81 | 1.05 | 1.03 | 0.58 | 0.78 |
| Oskar | 0.97 | 0.55 | 0.70 | 1.01 | 1.37 | 0.54 |
| LM(OSE) | 1.06 | 0.56 | 0.52 | 0.86 | 2.01 | 0.66 |
| METEOR | 0.97 | 0.85 | 0.91 | 0.92 | 1.05 | 0.92 |
| P-1 | 0.96 | 0.55 | 0.93 | 1.01 | 1.02 | 0.54 |
| Rohina | 1.04 | 0.79 | 0.94 | 0.71 | 1.1 | 1.11 |
| PEA-09 | 1.04 | 0.60 | 0.98 | 0.81 | 1.05 | 0.73 |
| LM-POIS | 0.97 | 1.03 | 0.89 | 0.69 | 1.08 | 1.49 |
| LAMPO | 1.02 | 0.71 | 1.01 | 0.94 | 1.00 | 0.75 |

Discussion

The root weight of peas genotypes greatly de-creases due to Cd stress (21-23) because Cd hinders the availability of other nutrients which were important for root growth (24). Surprisingly, it was observed in our study that the root growth of peas genotypes increased by increasing the Cd level by more than 4 ppm in the soil. Hence, Cd was applied in combination with nitrate, this could also be the reason for enhanced root growth due to a sufficient supply of nitrogen (25). The other reason could be the struggle of peas plants for survival under elevated Cd stress. As a result of high level of cadmium stress, roots

triggered itself under mechanism of stress resistance to get nutrients from soil that were hindered due to high application of Cd. Opposite results to our study were observed from other researchers as recently reduced peas root length was reported under Cd stress (26). A similar phenomenon was observed in the root length of almost all genotypes that Cd stress up to 4 ppm greatly reduced root length, but it was comparatively increased at 8 ppm Cd stress. (27) found reduced seedling weight when they subjected peas elevated Cd levels. The current study also supported the previous ones for decreased seedling weight with increased Cd level. Peas genotypes whose seedling weight was less affected by Cd stress can be selected for further studies to include in the next breeding programs to make Cd tolerant genotypes.

When the soil was polluted with Cd (28, 29) then grown vegetables had a higher accumulation of Cd in their parts (30). (31-34) reported elevated level of Cd in shoot of those plants grown under Cd rich soil. In the case of peas, a considerable reduction in the growth was also observed associated with increased Cd accumulation in the root and shoot (35). Peas mutants and *Brassica juncea* accumulated higher Cd in their shoot when grown under elevated levels (36). Those peas genotypes that enhanced their growth by stimulating more response to nitrogen availability could also be used for phytoremediation if they could be harvested at the seedling stage. The increased values of quality control indicators of more than 1 could suggest the higher uptake of Cd in respective parts of the plant. BAC indicated the transfer of Cd from the soil to the shoot of peas genotypes while BCF determined the transference of Cd from soil to root. TF explained the transfer of Cd from root to the shoot of peas genotypes. These are the quality control indicators and determine if the plant is storing metal in root or shoot. Those genotypes which have higher TF than 1, are considered to accumulate higher Cd in their shoot while those that have TF less than 1, have stored Cd in their roots.

Previously it has been determined that Cd adversely affects plant health (37) so, this study is the extension to suggest the negative effects of Cd on peas growth. Variation in response to Cd stress by all genotypes is also an important aspect of the study to suggest the genotypes for further studies. So, those genotypes which accumulated higher Cd and transferred to their shoot can be considered very important in the next studies to develop a breeding program in consideration of phytoremediation of Cd polluted soils.

4. CONCLUSION

10 peas genotypes were evaluated under two Cd stressed treatments (4 ppm and 8 ppm) and one control condition (0 ppm). There was a great variation in the data for all observed traits as every genotype differently responded to Cd application. Biochemical analysis of root, shoot and soil samples revealed that there was an elevated level of Cd in all samples when Cd applied in the soil but there was also variation in the data of Cd concentration in plant parts of each genotype because some genotypes might have tolerance at seedling stage against elevated Cd level. All morphological and biochemical analysis revealed the negative impact of Cd on both soil and peas genotypes. So, it is suggested to remediate Cd polluted soils and also develop Cd tolerant peas genotypes in aspect of increasing heavy metal stress in soils.

CONSENT

All authors declare that written informed consent was obtained from the patient (or other approved parties) for publication of this case report and accompanying images. A copy of

the written consent is available for review by the Editorial office/Chief Editor/Editorial Board members of this journal

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki

REFERENCES

1. Habib N, Zamin M. Off-season pea cultivation in Dir Kohistan valley. Asian Journal of Plant Sciences. 2003;2(3):283-5.
2. Arshad MN. Agro-Ecological Zoning in Punjab-Pakistan. 2019.
3. Dhall RK. Pea cultivation. Bulletin No PAU/2017/Elec/FB/E. 2017;29.
4. Chisti S, Iqbal M, Parveen N, Nadeem K, Iqbal M, Shahzad U, et al. Sarsabz: A new high yielding, well adapted pea (*Pisum sativum* L.) variety. Sarhad Journal of Agriculture. 2018;34(4):904-9.
5. Aznar Fernández T. Identification and characterization of resistance to biotic and abiotic stresses in pea. 2018.
6. Bisbis MB, Gruda N, Blanke M. Potential impacts of climate change on vegetable production and product quality—A review. Journal of Cleaner Production. 2018;170:1602-20.
7. Yadav S, Modi P, Dave A, Vijapura A, Patel D, Patel M. Effect of abiotic stress on crops. Sustainable Crop Production. 2020.
8. Naz S, Anjum MA, Ejaz S, Ali S, Saddiq B, Sardar H, et al. Sewage wastewater reclamation with sand column filter and reduction of heavy metal accumulation in tomato and okra. Environmental Science and Pollution Research. 2021:1-9.
9. Bosancic B, Zabic M, Mihajlovic D, Samardzic J, Mirjanic G. Comparative study of toxic heavy metal residues and other properties of honey from different environmental production systems. Environmental Science and Pollution Research. 2020;27(30):38200-11.
10. Mani PK, Mandal A, Mandal D, Irfan M, Hazra G, Saha S. Assessment of Non-Carcinogenic and Carcinogenic Risks Due to Ingestion of Vegetables Grown Under Sewage Water Irrigated Soils Near a 33 Years Old Landfill Site in Kolkata, India. Exposure and Health. 2021:1-22.
11. de Araújo RP, de Almeida A-AF, Pereira LS, Mangabeira PA, Souza JO, Pirovani CP, et al. Photosynthetic, antioxidative, molecular and ultrastructural responses of young cacao plants to Cd toxicity in the soil. Ecotoxicology and Environmental Safety. 2017;144:148-57.

12. Sonone SS, Jadhav S, Sankhla MS, Kumar R. Water contamination by heavy metals and their toxic effect on aquaculture and human health through food Chain. *Letters in applied NanoBioScience*. 2020;10(2):2148-66.
13. Haider FU, Liqun C, Coulter JA, Cheema SA, Wu J, Zhang R, et al. Cadmium toxicity in plants: Impacts and remediation strategies. *Ecotoxicology and Environmental Safety*. 2021;211:111887.
14. Huybrechts M, Cuypers A, Deckers J, Iven V, Vandionant S, Jozefczak M, et al. Cadmium and plant development: an agony from seed to seed. *International journal of molecular sciences*. 2019;20(16):3971.
15. Bahmani R, Bihamta M, Habibi D, Forozesh P, Ahmadvand S. Effect of cadmium chloride on growth parameters of different bean genotypes (*Phaseolus vulgaris* L.). *ARPN J Agric Biol Sci*. 2012;7:35-40.
16. Bahmani R, Habibi D, Bihamta M. Evaluation of common bean genotypes tolerance in response to cadmium stress at germination stage. 2015.
17. ur Rehman MZ, Rizwan M, Hussain A, Saqib M, Ali S, Sohail MI, et al. Alleviation of cadmium (Cd) toxicity and minimizing its uptake in wheat (*Triticum aestivum*) by using organic carbon sources in Cd-spiked soil. *Environmental Pollution*. 2018;241:557-65.
18. Jaouani K, Karmous I, Ostrowski M, El Ferjani E, Jakubowska A, Chaoui A. Cadmium effects on embryo growth of pea seeds during germination: Investigation of the mechanisms of interference of the heavy metal with protein mobilization-related factors. *Journal of plant physiology*. 2018;226:64-76.
19. Steel RG, Torrie JH. Principles and procedures of statistics, a biometrical approach. McGraw-Hill Kogakusha, Ltd.; 1980 Dec.
20. Hassan Z. Functional analysis of zinc hyperaccumulation related genes of *Noccaea (Thlaspi) caerulescens* for phytoremediation purposes. Wageningen University and Research; 2013.
21. Singh S, Zacharias M, Kalpana S, Mishra S. Heavy metals accumulation and distribution pattern in different vegetable crops. *Journal of Environmental Chemistry and Ecotoxicology*. 2012;4(4):75-81.
22. Garg N, Singla P, Bhandari P. Metal uptake, oxidative metabolism, and mycorrhization in pigeonpea and pea under arsenic and cadmium stress. *Turkish Journal of Agriculture and Forestry*. 2015;39(2):234-50.
23. Slima DF, Ahmed DAE-A. Trace metals accumulated in pea plant (*Pisum sativum* L.) as a result of irrigation with wastewater. *Journal of Soil Science and Plant Nutrition*. 2020;20(4):2749-60.
24. Nazar R, Iqbal N, Masood A, Khan MIR, Syeed S, Khan NA. Cadmium toxicity in plants and role of mineral nutrients in its alleviation. 2012.
25. Zhang R-R, Liu Y, Xue W-L, Chen R-X, Du S-T, Jin C-W. Slow-release nitrogen fertilizers can improve yield and reduce Cd concentration in pakchoi (*Brassica chinensis* L.)

grown in Cd-contaminated soil. *Environmental Science and Pollution Research*. 2016;23(24):25074-83.

26. Abbas T, Fan R, Hussain S, Sattar A, Khalid S, Butt M, et al. Protective effect of jasmonic acid and potassium against cadmium stress in peas (*Pisum sativum* L.). *Saudi Journal of Biological Sciences*. 2022;29(4):2626-33.

27. Shamshad S, Shahid M, Rafiq M, Khalid S, Dumat C, Sabir M, et al. Effect of organic amendments on cadmium stress to pea: a multivariate comparison of germinating vs young seedlings and younger vs older leaves. *Ecotoxicology and environmental safety*. 2018;151:91-7.

28. Grobelak A, Placek A, Grosser A, Singh BR, Almás AR, Napora A, et al. Effects of single sewage sludge application on soil phytoremediation. *Journal of Cleaner Production*. 2017;155:189-97.

29. Lwin CS, Seo B-H, Kim H-U, Owens G, Kim K-R. Application of soil amendments to contaminated soils for heavy metal immobilization and improved soil quality—A critical review. *Soil science and plant nutrition*. 2018;64(2):156-67.

30. Yang J, Guo H, Ma Y, Wang L, Wei D, Hua L. Genotypic variations in the accumulation of Cd exhibited by different vegetables. *Journal of Environmental Sciences*. 2010;22(8):1246-52.

31. Januškaitienė I. The effect of cadmium on several photosynthetic parameters of pea (*Pisum sativum* L.) at two growth stages. *Žemdirbystė= Agriculture*. 2012;99(1):73-85.

32. Tran TA, Popova LP. Functions and toxicity of cadmium in plants: recent advances and future prospects. *Turkish journal of Botany*. 2013;37(1):1-13.

33. García de la Torre VS, Coba de la Peña T, Pueyo JJ, Lucas MM. Cadmium-Tolerant and-Sensitive Cultivars Identified by Screening of *Medicago truncatula* Germplasm Display Contrasting Responses to Cadmium Stress. *Frontiers in plant science*. 2021;12:293.

34. Kolahi M, Kazemi EM, Yazdi M, Goldson-Barnaby A. Study of Oxidative Stress Indices, Morphological Response, Mineral Absorption in Chickpea (*Cicer Arietinum* L.) Under Cadmium Stress and Bioinformatics of HMA Poteins. 2021.

35. Metwally A, Safronova VI, Belimov AA, Dietz K-J. Genotypic variation of the response to cadmium toxicity in *Pisum sativum* L. *Journal of Experimental Botany*. 2005;56(409):167-78.

36. Belimov AA, Shaposhnikov AI, Azarova TS, Makarova NM, Safronova VI, Litvinskiy VA, et al. Microbial consortium of PGPR, rhizobia and arbuscular mycorrhizal fungus makes pea mutant SGECDt comparable with Indian mustard in cadmium tolerance and accumulation. *Plants*. 2020;9(8):975.

37. Cui J, Wang W, Peng Y, Zhou F, He D, Wang J, et al. Effects of simulated Cd deposition on soil Cd availability, microbial response, and crop Cd uptake in the passivation-remediation process of Cd-contaminated purple soil. *Science of the Total Environment*. 2019;683:782-92.