

# ASSESSMENT OF HEAVY METALS ON AFRICAN CATFISH (*CLARIAS GARIEPINUS*) FROM SOME FISH PONDS IN EMENE, ENUGU STATE, SOUTH-EAST NIGERIA

## Abstract

The research was conducted to determine the presence and bioaccumulation pattern of some heavy metals in African catfish (*Clarias gariepinus*) from three fish ponds in Emene, Enugu State, South-East Nigeria. Fish samples used for the analysis were collected during the dry season from Tollex, Sopulu and Nduka pond farms. The analysis of heavy metals: lead (Pb), cadmium (Cd), chromium (Cr), copper (Cu), zinc (Zn), and iron (Fe), in *C. gariepinus* gills and muscles was carried out using atomic absorption spectrophotometer (AAS) after digestion with concentrated HCl. Results of the analysis show that the investigated heavy metal in the fish samples from the ponds on the basis of their average concentration followed the order: Sopulu pond > Nduka pond > Tollex pond with highest concentration (7.427 ppm of Fe) found in the gill of fish from Sopulu fish pond and lowest concentration (0.013 ppm of Cd) in the muscle of fish from Tollex fish farm. The study has shown that there was a considerable amount of heavy metals in the muscle and gills of catfish from Tollex, Nduka and Sopulu fish farms.

Keywords: Toxic metal, *Clarias gariepinus*, Fish pond, Atomic absorption spectrometry, Emene-Enugu

## INTRODUCTION

Pollution of the environmental media could be as a result of pollutants such as polycyclic aromatic hydrocarbons (PAHs) and toxic metals which are ubiquitous. Pollutants' contamination of air, soil, water, food, beverages etc. has been reported by several scholars. For example, the reports of PAHs and heavy metals in air [1-3], soil [4-7], water [8-12], food [13-16], beverages [17-20] are well documented.

The contamination of aquatic environment especially by heavy metals affects not only the water body but also the aquatic organisms inhabiting the environment. According to Alam *et al.* [21], metallic elements are easily dissolved in and transported by water as well as quickly ingested by aquatic creatures. They also appear to be everywhere or in many locations simultaneously. The

density of heavy metals is at least five times that of water. Heavy metals typically do not degrade further into less dangerous components; instead, they build up where they are released and are hence stable and bio-accumulative [22]. They are sometimes passed up the food chain to humans [23]. At a certain amount of exposure, heavy metals are non-biodegradable and extremely damaging to aquatic life, vegetation, and people's health [24]. Some heavy metals, like copper or cobalt, are necessary for enzymatic function at low concentrations, but in greater quantities, they act as enzyme inhibitors.

At low levels, some heavy metals, such as copper or cobalt, are essential for enzymatic activity, but act as enzyme inhibitors at higher concentrations. Other metals, such as cadmium and lead, have no known essential role in living organisms and are toxic even at low concentrations [25]. In recent years, the need for a better understanding of heavy metal concentration and dispersion patterns in aquatic environments has been highlighted following the discovery of high levels of toxic heavy metals (particularly cadmium and lead) in fish and other living organisms [26, 27] and even in water resources [12], beverages [28, 29, 18, 19]. The most significant problem associated with heavy metals in the environment apart from accumulation through the food chain and persistence in nature is their toxicity [30].

Fishes, which are often at the top of the aquatic food chain absorb large amounts of these metals from both food chain and water [31]. Fishes have been considered good indicators of heavy metal contamination in aquatic systems because they occupy different trophic levels with different sizes and ages [32, 33]. Fish is widely consumed in many parts of the world by humans, and polluted fish may endanger human health. Studies on heavy metals in fish [34 - 37] have been a major environmental focus, especially during the last decade because fishes are notorious for their ability to accumulate heavy metals in their gills and muscles as they play important role in human nutrition. Consequently, heavy metals need to be carefully screened to ensure that unnecessarily high levels of some toxic metals are not being transferred to man through fish consumption [38]. *C. gariepinus* among the most commonly harvested and consumed fresh and pond water fish species and therefore a need for regular monitoring of the quality to prevent heavy metal contamination and bioaccumulation along the food chain.

In Enugu State, Emene is probably known to be an industrial area, thereby generating pollution by industrial activities by releasing pollutants especially heavy metals to the ecosystem each blessed day. Fish ponds in Emene have significant variations in the concentration of heavy metals

due to untreated industrial and sewage wastes that are added to the pond water through one channel or the other. For the majority of the people living in Enugu and Emene in particular, their only source of fish for protein and other beneficial uses is the pond fish. And so, it becomes necessary to examine the heavy metals concentration in fish harvested in some fish ponds in Emene, Enugu State, South-East Nigeria.

Studies have shown that all over the world, ponds are contaminated with heavy metals [39 – 41] and as a result, fishes harvested from them are seriously affected. Fish ponds in Emene, Enugu State, Nigeria may not be an exception as the ponds are natural habitats of fishes that are used for human consumption in the area. Hence, the health risks associated with heavy metal poisoning in humans via consumption of fish harvested from ponds cannot be overemphasized and it was against this backdrop that this research was conducted since there remains little or no literature on the level of heavy metals in Tollex, Sopulu, and Nduka pond farms in Emene, Enugu State. The current study therefore examines six heavy metals (Pb, Cd, Cr, Zn, Cu and Fe) concentrations in the gills and muscles of the African catfish, *C. gariepinus*.

Results obtained from this study would provide information on levels of heavy metals in the assessed fish species, thus contributing to the effective monitoring of the fish farms.

## MATERIALS AND METHODS

### Materials Procurement

The African catfish samples were collected from Tollex, Sopulu and Nduka fish ponds from Emene in Enugu state. All reagents and chemicals used during analysis were of analytical grade and pre-tested for possible heavy metal contamination. All the glassware and plastics were soaked overnight in 10% nitric acid and rinsed twice with distilled water before use. The instruments used were all calibrated and standardized with standard solutions prepared from commercially procured chemicals.

### Sample Preparation and Analysis

The samples were thawed at room temperature and then dissected for analysis using stainless steel scalpels. The gills and muscles on the dorsal surface of the fish were dried in an oven at 80°C for two hours. The samples were then removed from the oven and allowed to cool. Each

dried sample was pulverized using a porcelain mortar and pestle. For each pulverized fish sample, 0.3g was weighed with a Mettler H10 sensitive weighing balance and poured into a beaker containing 20mL of hydrochloric acid for digestion. Each digested sample was filtered using Whatman filter paper (0.45µm) and then transferred into a 50mL Erlenmeyer flask and diluted to mark with distilled water. Heavy metals (Pb, Cd, Cu, Cr, Zn and Fe) were determined in the solution using Atomic Absorption Spectrophotometry as adopted from a study by Orakwue *et al.* [42] with little modifications. Procedural blanks were run alongside samples as the analysis was processed in triplicate to ensure the quality of the results obtained. Commercial standards (Buck scientific) containing known concentration were further diluted to provide working standards of the following concentrations in ppm (0, 0.2, 0.4, 0.8 and 1.6) for each heavy metal and their corresponding absorbance recorded was used to plot the standard curve. From the standard curve, unknown concentrations of heavy metals in the samples were extrapolated. For each heavy metal, there is a specific hollow cathode lamp and an operational wavelength. The following: 283nm, 228.9nm, 357.9nm, 213.9nm, 325nm and 385nm are the wavelengths for lead, cadmium, chromium, zinc, copper and iron respectively.

## RESULTS AND DISCUSSION

There is the presence of all the heavy metals investigated in both the gills and the muscles of catfish species in all three ponds as shown in Table 1. Lead content in the muscle of the fish samples from the three ponds is Sopulu (0.137 ppm) > Nduka (0.047 ppm) > Tollex (0.033 ppm) while the order (Tollex (0.020 ppm) < Nduka and Sopulu (0.067 ppm) in the gills was observed. Pb concentrations in the muscle of catfish from Tollex and Sopulu ponds are higher than the metal concentration in the gills whereas the Pb concentration is higher in the gills than in the muscle of fish samples from Nduka pond. Lead poisoning in adults can affect the peripheral and central nervous systems, the kidneys, and blood pressure [43]. Hypertension has been associated with acute lead poisoning, along with renal failure. At lesser exposures, both experimental and epidemiological evidence of interference with renal function and elevations in blood pressure have been reported [44]. In children, exposure to Pb can cause acute symptomatic poisoning. Characteristics of this life-threatening syndrome are abdominal colic, constipation, fatigue, anemia, peripheral neuropathy, and in most cases, alteration of central nervous system function [45]. In severe cases, a full-blown acute encephalopathy with coma, convulsions, and

papilledema may occur. In milder cases, only headache or personality changes may be evident. In many instances, persons who have suffered from acute lead encephalopathy are left with permanent neurologic and behavioral sequelae [46].

Cadmium content in the muscle of the fish samples followed the order: Sopulu > Tollex > Nduka while that of the gills was of the order Tollex < Nduka < Sopulu. Cadmium induces tissue injury by creating oxidative stress [47 – 49], epigenetic changes in DNA [50 – 52], inhibition or upregulation of transport pathways [53 – 55] particularly in the proximal S1 segment of the kidney tubule [56].

Cu content in the muscle of the fish samples ranged between 1.197 ppm and 2.297 ppm with Tollex having the highest concentration while that of gills ranged from 1.180 ppm to 1.667 ppm with Nduka having the lowest concentration. Acute copper poisoning is rare. However, serious health problems from long-term exposure to copper can occur. Severe poisoning can cause liver failure and death [57]. In poisonings from a long-term buildup of copper in the body, the outcome depends on how much damage there is to the body's organs. Ingestion may lead to stricture formation throughout the gastrointestinal tract. Acute liver failure can occur due to direct copper toxicity-induced tissue necrosis [58].

Cr concentration in the muscles of the fish samples from the ponds indicated that Nduka > Sopulu > Tollex while that of the gills was of the pattern Sopulu > Nduka > Tollex. Chromium has been found to alter the epigenetic profile of cells at both the level of DNA methylation and histone modification [59 -61].

Zn content in the muscle of the fishes showed a range of 2.050 ppm to 3.580 ppm in the order Sopulu > Tollex > Nduka while the concentration in the gills indicated a range between 3.587 ppm and 4.923 ppm in the order Sopulu > Nduka > Tollex. There is a very low probability of zinc toxicity being fatal; however, the prognosis largely depends on how quickly the patient receives treatment. Large intentional ingestions are most likely to cause significant toxicity [62]. Severe toxicity may also mimic many other pathologies and rarely present with acute kidney injury, pancreatic injury, liver failure, and hemodynamic instability [63].

Table 1: Concentration of the investigated heavy metals (ppm) in muscles and gills of catfish in the selected fish ponds

	Tollex Pond		Nduka Pond		Sopulu Pond	
	Muscle	Gill	Muscle	Gill	Muscle	Gill
Lead	0.04	0.01	0.04	0.06	0.15	0.07
	0.03	0.03	0.05	0.06	0.1	0.08
	0.03	0.02	0.05	0.08	0.16	0.05
<b>AVERAGE</b>	<b>0.033</b>	<b>0.020</b>	<b>0.047</b>	<b>0.067</b>	<b>0.137</b>	<b>0.067</b>
<b>STD</b>	<b>0.006</b>	<b>0.010</b>	<b>0.006</b>	<b>0.012</b>	<b>0.032</b>	<b>0.015</b>
Cadmium	0.01	0.04	0.03	0.05	0.06	0.07
	0.02	0.06	0.03	0.06	0.08	0.09
	0.01	0.05	0.03	0.05	0.1	0.08
<b>AVERAGE</b>	<b>0.013</b>	<b>0.050</b>	<b>0.030</b>	<b>0.053</b>	<b>0.080</b>	<b>0.080</b>
<b>STD</b>	<b>0.006</b>	<b>0.010</b>	<b>0.000</b>	<b>0.006</b>	<b>0.020</b>	<b>0.010</b>
Copper	2.13	1.32	1.25	1.65	1.16	1.19
	2.11	1.44	1.22	1.66	1.19	1.18
	2.65	1.26	1.2	1.69	1.24	1.17
<b>AVERAGE</b>	<b>2.297</b>	<b>1.340</b>	<b>1.223</b>	<b>1.667</b>	<b>1.197</b>	<b>1.180</b>
<b>STD</b>	<b>0.306</b>	<b>0.092</b>	<b>0.025</b>	<b>0.021</b>	<b>0.040</b>	<b>0.010</b>
Chromium	0.07	0.05	0.08	0.07	0.06	0.07
	0.04	0.08	0.09	0.07	0.06	0.08
	0.05	0.06	0.09	0.06	0.05	0.07
<b>AVERAGE</b>	<b>0.053</b>	<b>0.063</b>	<b>0.087</b>	<b>0.067</b>	<b>0.057</b>	<b>0.073</b>
<b>STD</b>	<b>0.015</b>	<b>0.015</b>	<b>0.006</b>	<b>0.006</b>	<b>0.006</b>	<b>0.006</b>
Zinc	3.54	3.76	2.01	4.16	3.54	5.01
	2.97	3.35	2	4.22	3.44	4.98
	3.89	3.65	2.14	4.17	3.76	4.78
<b>AVERAGE</b>	<b>3.467</b>	<b>3.587</b>	<b>2.050</b>	<b>4.183</b>	<b>3.580</b>	<b>4.923</b>
<b>STD</b>	<b>0.464</b>	<b>0.212</b>	<b>0.078</b>	<b>0.032</b>	<b>0.164</b>	<b>0.125</b>
Iron	5.15	4.69	5.21	6.12	7.18	7.29
	5.22	5.02	5.29	6.22	7.19	7.43
	5.44	5.11	5.22	6.17	7.23	7.56
<b>AVERAGE</b>	<b>5.270</b>	<b>4.940</b>	<b>5.240</b>	<b>6.170</b>	<b>7.200</b>	<b>7.427</b>
<b>STD</b>	<b>0.151</b>	<b>0.221</b>	<b>0.044</b>	<b>0.050</b>	<b>0.026</b>	<b>0.135</b>

Ingested iron can cause direct caustic injury to the gastrointestinal mucosa, resulting in nausea, vomiting, abdominal pain, and diarrhea. Significant fluid and blood loss can lead to hypovolemia. Hemorrhagic necrosis of gastrointestinal mucosa can lead to hematemesis, perforation, and peritonitis [64]. At the cellular level, iron impairs cellular metabolism in

the heart, liver, and central nervous system. Free iron enters cells and concentrates in the mitochondria. This disrupts oxidative phosphorylation, catalyzes lipid peroxidation, forms free radicals, and ultimately leads to cell death [65]. Fe concentration was of the order Sopulu > Tollex > Nduka in the muscle of the fish samples while, in the gills, it was Sopulu > Nduka > Tollex. The concentration ranged between 4.940 ppm in the gill of fish from Tollex to 7.427 in the gill of the fish from Sopulu.

## CONCLUSION

Present research reveals significant variations in the concentration of the investigated heavy metals in the fish samples from the ponds which may be as a result of untreated industrial and sewage wastes that leach into the pond water. The study has shown that there was a considerable amount of heavy metals in the fish tissues from Tollex, Nduka and Sopulu fish farms. In most cases, Sopulu pond showed higher levels of heavy metals than the other sources. Nduka pond likewise showed higher levels of heavy metals than Tollex pond which showed the lowest levels of heavy metals in the entire sources. In general, catfish tissue samples showed higher levels of heavy metal in the gills than in the muscles from the three fish pond sources when compared to catfish tissue samples. Heavy metal concentrations in the fish samples from the ponds on the basis of their average followed the order: Sopulu pond > Nduka pond > Tollex pond. Further study is recommended for levels of toxic metals including silver (Ag), Arsenic (As), mercury (Hg), nickel (Ni) and their exposure risks from the fish harvested from the ponds. Again, the levels of heavy metals in the ponds should be continuously monitored to check on their levels in order to maintain standard recommended by appropriate authorities.

## REFERENCES

1. Abayalath N, Malshani I, Ariyaratne R, Zhao S, Zhong G, Zhang G, Manipura A, Siribaddana A, Karunaratne P, Kodithuwakku SP. Characterization of airborne PAHs and metals associated with PM10 fractions collected from an urban area of Sri Lanka and the impact on airway epithelial cells. *Chemosphere*. 2020; <https://doi.org/10.1016/j.chemosphere.2021.131741>.

2. Baldantoni D, Nicola FD, Alfani A. Air biomonitoring of heavy metals and polycyclic aromatic hydrocarbons near a cement plant. *Atmospheric Pollution Research*. 2014; 5(2): 262-269. <https://doi.org/10.5094/APR.2014.032>.
3. Samdudin M, Azahar H, Abas A, Zakaria Z. Determination of heavy metals and polycyclic aromatic hydrocarbons (PAHs) contents using the Lichen *Dirinaria picta* in Universiti Kebangsaan Malaysia. *Journal of Environmental Protection*. 2013; 4(8): 760-765. <https://doi.org/10.4236/jep.2013.48088>.
4. Ali M, Song X, Ding D, Wang Q, Zhang Z, Tang Z. Bioremediation of PAHs and heavy metals co-contaminated soils: Challenges and enhancement strategies. *Environmental Pollution*. 2022. <https://doi.org/10.1016/j.envpol.2021.118686>.
5. Gran-Scheuch A, Ramos-Zuñiga J, Fuentes E, Bravo D, Pérez-Donoso JM. Effect of co-contamination by PAHs and heavy metals on bacterial communities of diesel contaminated soils of South Shetland Islands, Antarctica. *Microorganisms*. 2020; 7;8(11):1749. <https://doi.org/10.3390/microorganisms8111749>.
6. Bandowe BAM, Shukurov N, Leimer S. *et al.* Polycyclic aromatic hydrocarbons (PAHs) in soils of an industrial area in semi-arid Uzbekistan: spatial distribution, relationship with trace metals and risk assessment. *Environ Geochem Health* 2021; 43:4847–4861. <https://doi.org/10.1007/s10653-021-00974-3>.
7. Akinsete SJ, Olatimehin AD. Priority polycyclic aromatic hydrocarbons and heavy metals in urban roadside soils of heavy-traffic density areas in Ibadan, Nigeria: Levels, sources and health risk assessment. *Environmental Forensics*. 2023; <https://doi.org/10.1080/15275922.2023.2172092>
8. Zhu Y, Yang Y, Liu M, Zhang M, Wang J. Concentration, distribution, source, and risk assessment of PAHs and heavy metals in surface water from the three Gorges reservoir, China. *Human and Ecological Risk Assessment: An International Journal*, 2015; 21(6):1593-1607. <https://doi.org/10.1080/10807039.2014.962315>
9. Yan M, Nie H, Wang W, Huang Y, Wang J. Occurrence and toxicological risk assessment of polycyclic aromatic hydrocarbons and heavy metals in drinking water

- resources of Southern China. *Int J Environ Res Public Health*. 2018;15(7):1422. <https://doi.org/10.3390/ijerph15071422>.
10. Nubi AO, Popoola SO, Dada OA, Oyatola OO, Unyimadu JP, Adekunbi OF, Salami AM. Spatial distributions and risk assessment of heavy metals and PAH in the southwestern Nigeria coastal water and estuaries, Gulf of Guinea. *Journal of African Earth Sciences*. 2022. <https://doi.org/10.1016/j.jafrearsci.2022.104472>.
11. Okafor VN, Omokpariola DO, Igbokwe EC, Theodore CM, Chukwu NG. Determination and human health risk assessment of polycyclic aromatic hydrocarbons (PAHs) in surface and ground waters from Ifite Ogwari, Anambra State, Nigeria. *Inter J Environ Anal Chem*. 2022b; <https://doi.org/10.1080/03067319.2022.2038587>
12. Okafor VN, Omokpariola DO, Obumselu OF, Eze GC. Exposure risk to heavy metals through surface and groundwater used for drinking and household activities in Ifite Ogwari, Southeastern Nigeria. *Appl Water Sci*. 2023; <https://doi.org/10.1007/s13201-023-01908-3>
13. Amos-Tautua BMW, Inengite AK1, Abasi CY, Amirize GC. Evaluation of polycyclic aromatic hydrocarbons and some heavy metals in roasted food snacks in Amassoma, Niger Delta, Nigeria. *African Journal of Environmental Science and Technology*. 7(10): 961-966. 2013; <https://doi.org/10.5897/AJEST2013.1545>
14. Nikolina J, Novakov NJ, eljko A, Mihaljev ŽA, Brankica D, Kartalović BD, Bojan J, Blagojević BJ, Jelana M, Petrović JM, Miroslav A, Ćirković MA, Dragan R, Rogan DR. Heavy metals and PAHs in canned fish supplies on the Serbian market. *Food Additives and Contaminants: Part B*. 2017;10(3): 208-215. <https://doi.org/10.1080/19393210.2017.1322150>
15. Rai PK, Lee SS, Zhang M, Tsang YF, Kim K. Heavy metals in food crops: Health risks, fate, mechanisms, and management. *Environment International*. 2019; 125:365-385. <https://doi.org/10.1016/j.envint.2019.01.067>.
16. Bansal V, Kim K. Review of PAH contamination in food products and their health hazards. *Environment International*. 2015. 84: 26-38. <https://doi.org/10.1016/j.envint.2015.06.016>.

17. Okafor VN, Uche UB, Abailim RC. Levels of polycyclic aromatic hydrocarbons (PAHs) in beers: Consumption and public health concerns: *Chemical Sciences International Journal*, 29 (1), 47-59, February, 2020; <https://doi.org/10.9734/CSJI/2020/v29i130157>.
18. Okafor VN, Umennadi PU, Odidika CC, Vinna DC. Metals and polycyclic aromatic hydrocarbons (PAHs) in beer: a review. *J Chem Soc Nig.* 2021; <https://doi.org/10.46602/jcsn.v46i4.646>
19. Okafor VN, Omokpariola DO, Okabekwa CV, Umezina EC. Heavy metals in alcoholic beverages consumed in Awka, South-East Nigeria: Carcinogenic and non-carcinogenic health risk assessments. *Chemistry Africa.* 2022a; <https://doi.org/10.1007/s42250-022-00477-3>
20. Okafor VN, Omokpariola DO, Agu MO, Odidika CC, Okabekwa CV, Ogbuo LC, Obiatuegwu MC. Assessment of polycyclic aromatic hydrocarbons (PAHs) in alcoholic beverages consumed in Awka, Southeast Nigeria, *Bull. Chem. Soc. Ethiop.* 2023; 37(4), 805-815.
21. Alam MGM, Tanaka A, Allison G, Laurenson LJB, Stagnitti F, Snow, E. A comparison of trace element concentrations in cultured and wild carp (*Cyprinus carpio*) of lake Kasumigaura, Japan. *Ecotoxicology and environmental safety.* 2002; 53:348-354.
22. Kennicutt MC, Wade TL, Presley BJ. Assessment of sediment contamination in Casco Bay. *Casco Bay Estuary Project.* Texas A&M University. 1992; 113 pp.
23. Fergusson JE. (1990) *The heavy elements: Chemistry, environmental impact and health effects.* Pergamon Press, Oxford. 1990; 85-547
24. Zaynab M, Al-Yahyai R, Ameen A, Sharif Y, Ali L, Fatima M, Khan KA, Li S. Health and environmental effects of heavy metals. *Journal of King Saud University – Science.* 34:(1). <https://doi.org/10.1016/j.jksus.2021.101653>.
25. Bryan GW. *Proceeding of the Royal Society London. Series 177.* 389-410. 1976. Some effects of heavy metal tolerance in aquatic organisms. In: *Effects of pollutants on aquatic organisms.* A.P.M. Lockwood. Ed. Cambridge, England: Cambridge University. 1971; 7–34.
26. McConchie DM., Mann AW, Lintern MJ, Longman D, Talbot V, Gabelish AJ, Needleman H. *Annu. Rev. Med.* 2003; 200(55):209–2022. <https://doi.org/10.1146/annurev.med.55.091902.103653>.

27. Storelli MM, Storelli A, D'ddabbo R, Marano C, Bruno R, Marcotrigiano GO. Trace elements in loggerhead turtles (*Caretta caretta*) from the eastern Mediterranean Sea: Overview and evaluation. *Environmental Pollution*. 2005; 135:163–170.
28. Okafor VN. Hops and potential Nigerian substitutes: Comparative studies in beer brewing, LAP, Lambert Academic Publishing- GmbH & Co. KG, Germany. 2016; ISBN-10 (97833330014817).
29. Okafor VN. Spectroscopic methods for evaluation of hop extracts and extract from *Gongronema latifolium* as substitute in the Nigerian beer industry: *International Journal of Advanced Research*. 2016; 4 (12):2499-2504. <https://doi.org/10.21474/IJAR01/2675>.
30. Censi P, Spoto SE, Saiano F, Sprovieri M, Mazzola S, Nardone G, Di Geronimo SI, Punturo R, Ottonello D. Heavy metals in coastal water systems. A case study from the northwestern Gulf of Thailand. *Chemosphere*. 2006; 64: 1167–1176.
31. Gibson RN. Impact of habitat quality and quantity on the recruitment of juvenile flatfishes. *Netherlands journal of Sea Research* 1994: 32: 191.
32. Burger J, Gaines KF, Shane BC, Stephens WL, Snodgrass J, Dixon C, McMahon M, Shukla S, Shukla T, Gochfeld M. Metal levels in fish from the Savannah River: potential hazards to fish and other receptors. *Environmental Research*. 2002; 89: 85-91.
33. Rashed MN. Monitoring of environmental heavy metals in fish from Nasser lake. *Environment International*. 2001; 27: 27–33.
34. Özmen H, Külahç F, Çukuroval A, Doğru M. Concentrations of heavy metal and radioactivity in surface water and sediment of Hazar Lake (Elazığ, Turkey). *Chemosphere*. 2004; 55: 401–408.
35. Nyarko E, Boateng CM, Asamoah O, Edusei MO, Mahu E. Potential human health risks associated with ingestion of heavy metals through fish consumption in the Gulf of Guinea. *Toxicol Rep*. 2023; 10:117-123. <https://doi.org/10.1016/j.toxrep.2023.01.005>.
36. Łuczyńska J, Pietrzak-Fiećko R, Purkiewicz A, Łuczyński MJ. Assessment of fish quality based on the content of heavy metals. *Int J Environ Res Public Health*. 2022; 19(4):2307. <https://doi.org/10.3390/ijerph19042307>.

37. Hashim R, Song TH, Muslim NZ, Yen TP. Determination of heavy metal levels in fishes from the lower reach of the Kelantan river, Kelantan, Malaysia. *Trop Life Sci Res.* 2014; 25(2):21-39.
38. Adeniyi AA, Yusuf KA. Determination of heavy metals in fish tissues, water and bottom sediments from Epe and Badagry Lagoons, Lagos, Nigeria. *Environmental Monitoring Assessment.* 2007; 37: 451-458.
39. Saah SA, Adu-Poku D, Boadi N. Heavy metal contamination and water quality of selected fish ponds at Sunyani, Ghana: A comparison with WHO standards. *Chemistry International.* 2021; 7(3):181-187., Available at SSRN: <https://ssrn.com/abstract=3865906>.
40. Onuoha S. Assessment of metal contamination in aquaculture fish ponds South Eastern, Nigeria. *World Applied Sciences Journal.* 2017; 35(1):124-127 <https://doi.org/10.5829/idosi.wasj.2017.124.127>
41. Leonard LS, Mahenge A, Mudara NC. Assessment of heavy metals contamination in fish cultured in selected private fishponds and associated public health risk concerns, Dar es Salaam, Tanzania. *Marine Science and Technology Bulletin,* 2022; 11(2), 246- 258. <https://doi.org/10.33714/masteb.1108314>
42. Orakwue FC, Okafor VN, Obumselu, FO, Okoli JO, Nnamdi KC. Phytochemical, antimicrobial screening and human health risk assessment of heavy metals of stem-bark and root extracts of *Newbouldia laevis* (boundary tree). *J Chem Soc Nig.* 2021; <https://doi.org/10.46602/jcsn.v46i5.673>
43. Vupputuri S, Muntner P, Bazzano LA, *et al.* Blood lead level is associated with elevated blood pressure in blacks. *Hypertension.* 2003; 41:463–68.
44. Evans M, Elinder C. Chronic renal failure from lead: myth or evidence-based fact?, *Kidney International.* 2021; 79(3):272-279. <https://doi.org/10.1038/ki.2010.394>.
45. Cullen MR, Robins JM, Eskenazi B. Adult inorganic lead intoxication: Presentation of 31 new cases and a review of recent advances in the literature. *Medicine (Baltimore)* 1983; 62:221-247.

46. Byers RK, Lord EE. Late effects of lead poisoning on mental development. *Am J Dis Child* 1943; 66:471-494.
47. Matovic V, A. Buha, Z. Bulat, and D. Dukić. Cadmium toxicity revisited: focus on oxidative stress induction and interactions with zinc and magnesium. *Arhiv za Higijenu Rada i Toksikologiju*. 2011; 62(1):65–76.
48. Patra RC, A. K. Rautray, D. Swarup. Oxidative stress in lead and cadmium toxicity and its amelioration. *Veterinary Medicine International*. 2011; Article ID 457327, 2011.
49. Cuypers AM, Plusquin T, Remans *et al.* Cadmium stress: an oxidative challenge. *BioMetals*, 2010; 23(5): 927–940.
50. Wang B, C. Shao C, Y. Li Y, Y. Tan Y, L. Cai L. Cadmium and its epigenetic effects. *Current Medicinal Chemistry*. 2012; 19(16):2611–2620.
51. Martinez-Zamudio R, Ha HC. Environmental epigenetics in metal exposure. *Epigenetics*. 2011; 6(7):820–827.
52. Luparello C, Sirchia R, Longo A, Cadmium as a transcriptional modulator in human cells. *Critical Reviews in Toxicology*. 2011; 41(1):75–82.
53. Thevenod F. Catch me if you can! Novel aspects of cadmium transport in mammalian cells. *BioMetals*. 2010; 23 (5):857– 875.
54. Wan L, H. Zhang. Cadmium toxicity: effects on cytoskeleton, vesicular trafficking and cell wall reconstruction. *Plant Signaling & Behavior*. 2012; 7(3):345–348.
55. Kerkhove VE, Pennemans V, Swennen Q, Cadmium and transport of ions and substances across cell membranes and epithelia. *BioMetals*. 2010; 23(5):823–855.
56. Vesey DA. Transport pathways for cadmium in the intestine and kidney proximal tubule: focus on the interaction with essential metals. *Toxicology Letters*. 2010; 198(1):13–19.

57. Royer A, Sharman T. Copper Toxicity. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023; <https://www.ncbi.nlm.nih.gov/books/NBK557456/> [Updated 2023 Mar 27].
58. Gamakaranage CS, Rodrigo C, Weerasinghe S, Gnanathanan A, Puvanaraj V, Fernando H. Complications and management of acute copper sulphate poisoning; a case discussion. *J Occup Med Toxicol*. 2011; 19(6):1-34.
59. Rager JE, Suh M, Chappell GA, Thompson CM, Proctor DM. Review of transcriptomic responses to hexavalent chromium exposure in lung cells supports a role of epigenetic mediators in carcinogenesis. *Toxicol Lett*. 2019; 305:40-50. <https://doi.org/10.1016/j.toxlet.2019.01.011>.
60. Arita A, Shamy MY, Chervona Y, Clancy HA, Sun H, Hall MN, Qu Q, Gamble MV, Costa M. The effect of exposure to carcinogenic metals on histone tail modifications and gene expression in human subjects. *J Trace Elem Med Biol*. 2012; 26(2-3):174-8. <https://doi.org/10.1016/j.jtemb.2012.03.012>.
61. Chervona Y, Costa M. The control of histone methylation and gene expression by oxidative stress, hypoxia, and metals. *Free Radic Biol Med*. 2012. 53(5): 1041–1047.
62. Plum LM, Rink L, Haase H. The essential toxin: impact of zinc on human health. *Int J Environ Res Public Health*. 2010; 7(4):1342-65
63. Agnew UM, Slesinger TL. Zinc Toxicity. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK554548/> [Updated 2022 Dec 11].
64. Yuen HW, Becker W. Iron Toxicity. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing. 2023; Available from: <https://www.ncbi.nlm.nih.gov/books/NBK459224/> [Updated 2023 Jun 26].
65. Baranwal AK, Singhi SC. Acute iron poisoning: management guidelines. *Indian Pediatr*. 2003; 40(6):534-40.