

# Comparative Analysis of the Essential Oil Contents of *Azadirachta indica* and *Humulus lupulus*

## ABSTRACT

Hops give beer its brewing qualities (iso-alpha acid and essential oil). Hop acid is responsible for bitterness while hop essential oil is responsible for aroma and palate fullness. The aim of the study is to investigate the potentiality of substituting hop oil with neem oil in beer brewing. The oils were analyzed using Fourier-Transform Infrared (FTIR) spectroscopic technique after extracting the oil by solvent extraction. Hydrocarbon and Sulphur contents were also investigated using their standard methods. FTIR results show that both oils contain similar functional groups. It was observed that some hydrocarbons (myrcene, ocimene and  $\alpha$  and  $\beta$ -pinene) present in essential oil of hop were also present in neem leaf oil. Volatile sulphur compounds such as dimethyl sulphide, dimethyl disulphide and dimethyltrisulphide present in hop oil were also found in neem oil. This study revealed that the brewing neem oil can substitute hop oil in beer brewing.

Keywords: *Azadirachta indica*, *Humulus lupulus*, Essential oil, spectroscopic method, beer brewing

## INTRODUCTION

A specific raw material is used to make every industrial product. Plants are a significant supplier of raw materials for many industrial products, including apparel, footwear, and a wide range of other items (Jegannathan and Nielsen, 2013). Additionally, plants serve as a source of raw materials for the production of biofuels (Hood, 2016), dyes (Yadav et al., 2023), fragrances (Ameh et al., 2021) insecticides ((Bos *et al.*, 1997; Blank *et al.*, 2011),), medicines (Sofowora et al., 2013), and beverages like beer (Okafor et al., 2016a;b; Eboatu et al., 2016).

Beer is consumed worldwide and in Nigeria, beer consumption has increased tremendously due to large population of youth with disposable income (Okafor et al., 2017a;b). According to a new analysis by the Worldwide Brewing Alliance (WBA), the beer industry had a \$2.2 billion impact on the Nigerian economy in 2019. The WBA stated during a webinar that the report, which was created by Oxford Economics, revealed the sector's impact on various economies. According to the paper, Nigeria's economy benefits from the domestic, international, and global supply chains of the global beer industry. The worldwide beer industry supported a \$2.2 billion gross value-added contribution to Nigeria's GDP in 2019. This was the same as 0.6% of the country's GDP, or 73% of Zaria's economy (<https://bizwatchnigeria.ng/wba-reports-beer-sector-contributed-2-2bn-to-nigerian-economy>, 2022). As a result, a sizable share of imports into the nation have continued to be hops, which are needed to satisfy the demands of the already flourishing brewing industry. The main ingredients used to make beer are malt, hops, water, and yeast. The brewing industry depends on hop plants because some of its special compounds may find use in the nutraceutical sector (Okafor, 2016).

Hops are used for their ability to bitter taste, flavour, and enhance aroma in beer. Hops are made up of essential oils and acids ( $\alpha$  and  $\beta$  acids). In contrast to the acids, which are used for bittering and improving foam stability as well as acting as an antibiotic against microorganisms, hop's essential oils are used to enhance flavours and aromas (Hough *et al.*, 1982; Okafor et al., 2020a;b).

The quality of hops and beer is currently only partially studied in the literature even though, the substitution of hops with Nigerian bitter vegetables abounds in literature (Rodolfiet *al.*, 2023; Okafor et al., 2020a;b;c;d). There hasn't been any research that used the identical experimental setup and also took into account both of these factors and the substitution of hop essential oil with *A. indica* oil. In the current study, the impact of fragrance improvement on beers infused with *A. indica* oil using a multidisciplinary approach for characterization, i.e., FTIR, GC/MS, and GC/FID was examined for the first time.

The aim of this research is to study comparatively the essential oils of hop and *A. indica* in order to establish the potentiality of substituting the hop oil with that of *A. indica* in tropical beer brewing. The objectives are (a) to extract essential oil from *A. indicaleaves* (b)to analyze the oil extracts for functional group analysis using FTIR, (c) to determine the hydrocarbon content of

the oil extracts using GC/MS, (d) to determine the sulphur content of the oil extracts GCFID. The substitution of hops with *A. indicain* beer brewing in Nigeria would not only save Nigeria's hard-earned foreign exchange since Nigeria's brewing industry won't need to import hops but would give our local farmers jobs in the Agricultural sector.

## MATERIALS AND METHODS

### **Extraction of Essential oil from *A. indica***

The leaves of *A.indica* used for the research was obtained in in the herbarium of Nnamdi Azikiwe University, Awka, Anambra State and was identified by a taxonomist. The leaves were dried after which they were pulverized into powder using manual grinder. A 352g of the pulverized powder was placed into the thimble and placed in the Soxhlet chamber containing 500mL N-hexane. After completing the extraction process, the solvent and extractor were placed on water bath to evaporate the solvent and the oil was collected.

### **Sample Preparation for FTIR Analysis**

We employed a protocol used by Ezea *et al.* (2019) in the preparation of the oil for FTIR analysis. To get spectra heights, 0.5 mL of the oil was combined with 0.5 g of Kbr powder, and 1 mL of Nujol was pipetted into the mixture to create a paste. This paste was then placed into the instrument sample mould and allowed to scan at wavelengths between 600 cm<sup>-1</sup> and 4000 cm<sup>-1</sup>.

### **GC-MS and GC-FID analysis**

Method adopted by some researchers in the determination of hydrocarbon and volatilesulphur compounds (Bakkali et al., 2008; Figueredo et al., 2009; Jaimand et al., 2006) was used in this work. Briefly, gas chromatography-flame ionization detection (GC-FID) and gas chromatography-mass spectrometry (GC-MS) were carried out using an Agilent 6890N gas chromatograph and an Agilent 5973N mass selective detector (70 eV). Helium was used as the carrier gas, and its flow rate was 1000 L/min. A sample volume of 1 L (1% solution) was injected. The range of the scanning frequency was 50 to 550 m/z. A capillary column made of HP-5MS (30 m x 0.25 mm x 0.25 m) was used. The column temperature was first set at 50°C for 2 minutes, then increased to 150°C at 2 degrees per minute for 2 minutes, held for 2 minutes, and

then increased to 250°C at 10 degrees per minute for 5 minutes. In order to compute the retention index (RI), a homologous sequence of n-alkanes (C<sub>8</sub>-C<sub>40</sub>) were used.

## RESULTS AND DISCUSSION

Table 1 presents all characteristic spectra observed in the sample for essential oils of *A. indica* as well as specific functional groups assigned to particular vibration spectra. *A. indica* essential oil displays very intense and distinct bands, which can be attributed to specific vibrations of appropriate functional groups corresponding in turn to the components contained therein. Essential oils are mixtures of various compounds, for example, carvone, limonene, pinene, cymol, terpenic alcohols, menthol and menthone with various isomers, piperitone, methyl acetate, and germacrene. In many studies, the authors have made appropriate adjustments to specific bands in the spectra of essential oils of plant origin of various species (Guillen and Cabo, 1997; 2002; Rohman et al., 2009; Tapp et al., 2003) regarding specific vibrations in molecules or their moieties. However, many bands are difficult to properly assign to a specific functional group which may correspond to the content of a given amount of different substances (the reason is the fact that the substances present in particular oils often have similar chemical compositions). Table 1 presents in detail the frequencies of the characteristic spectra, along with the major extensions of the respective bands in the spectra of the essential oil sample, their assignment to the appropriate functional group. The bottom indices also show the intensity of observed bands in typical spectra in the IR region.

The results of hydrocarbon content in both hop and neem oils indicated that some hydrocarbons (myrcene, ocimene and  $\alpha$  &  $\beta$ -pinene) present in essential oil of hop were also present in *A. indica* essential oil while humulene and caryophyllene were only present in the essential oil of hop but absent in *A. indica*. As secondary metabolites, essential oil components are influenced by environmental (climatic and soil) factors and plant organs (Koutsaviti et al., 2018; Manion and Widder, 2017; Sarkhail, 2014; Zahang et al., 2019; Yadav et al., 2019; Calo et al., 2015) and this may explain the reason for the discrepancies in the composition of essential oil in *A. indica* and hops. The result however was in partial agreement with the findings of Dieckmann and Palamand (1974).

Table 1: Result of FTIR spectra of *Azadiratcha indica* oil

S/N	Wavelength(cm <sup>-1</sup> )	Functional group	Compounds
1	752.3616	C-Br	Bromo compound C-Br stretch
2	842.7057	C-Cl	Aliphatic Chloro compound C-Cl stretch
3	1025.068	C-O-C	Ether CO symmetric stretch
4	1378.737	H <sub>2</sub> C-CH <sub>3</sub>	Ethane CH symmetric stretch
5	1477.960	H-CW	Ethene C=C anti-symmetric stretch
6	1620.787	R-COONH <sub>2</sub>	Amide CN stretch
7	1838.131	R-COOR	Cyclic ester COO stretch
8	2058.256	RCOO	Carboxylic acid CO stretch
9	2226.013	R <sub>2</sub> C=O	Carbonyl C=O anti- symmetric stretch
10	2294.906	R <sub>2</sub> C-O	Carbonyl C=O anti-symmetric stretch'
11	2448.348	R-C=N	Nitrile CN anti-symmetric stretch
12	2524.535	CH <sub>2</sub> SH	Thiol SH stretch
13	2766.888	CH <sub>2</sub>	Methylene CH symmetric stretch
14	2949.748	R-S-ON	Thiocyanate SCN antisymmetric stretch
15	3043.655	RCHOH	1° alcohol OH stretch
16	3197.924	RCHOH	1° alcohol OH stretch
17	3444.58	R-NH <sub>2</sub>	2 amine N-H bend
18	3695.667	R <sub>3</sub> N	3 amine N-H bend
19	381 1.956	R <sub>3</sub> CHOH	Phenol OH symmetric stretch

Volatile sulphur compounds such as dimethyl sulphide, dimethyl disulphide and dimethyltrisulphide present in hop oil were also found in *A. indica* oil while 3-(methylthio)propanal, 3-(methylthio)propanol, 3-(methylthio)propyl acetate, ethyl 3-(methylthio)-propanoate, dihydro-2-methyl-3(2H)-thiophenone, 2-thiophene-carboxyaldehyde, 5-methyl-2-thiophenecarboxyaldehyde, benzothiophene and benzothiazole were present in hops

only (Houghs et al., 1982). Therefore, in terms of volatile sulphur compounds, *A. indica* oil could not be a good alternative for hops in beer brewing.

## CONCLUSION

FTIR results show that both oils contain similar functional groups. Myrcene, ocimene, and  $\alpha$  and  $\beta$ -pinene were present in both essential oil of hop and *A. indica* oil. Sulphur compounds such as dimethyl sulphide, dimethyl disulphide and dimethyltrisulphide present in hop oil were also found in *A. indica* oil. This study revealed that the essential oil of *A. indica* possessed promising potential in aroma enhancement in tropical beer brewing. Consequently, it is recommended that beer brewing industries may consider the use of *A. indica* as possible hop substitute.

## REFERENCES

Ameh OE, Achika JI, Bello NM, Owolaja, AJ (2021) Extraction and Formulation of Perfume from *Cymbopogon citratus* (Lemongrass) J. Appl. Sci. Environ. Manage. Vol. 25 (8) 1461-1464 August 2021

Bakkali F, S. Averbeck, D. Averbeck and M. Waomar (2008). Biological effects of essential oils – A review, Food Chem. Toxicol. 46, 446-475.

Blank A. F., T.C.P. Sant'ana, P.S. Santos, M.F. Arrigoni-Blank, A.P.D. Prata, H.C.R. Jesus and P.B. Alves (2011). Chemical characterization of the essential oil from patchouli accessions harvested over four seasons, Ind. Crops Prod. 34, 831-837

Bos R., H.J. Woerdenbag, F.M. van Putten, H. Hendriks and J.J. Scheffer (1997) Seasonal variation of the essential oil, valerenic acid and derivatives and valepotriates in *Valeriana officinalis* roots and rhizomes and the selection of plants suitable for phytomedicines, Planta Med. 64, 143-147.

BWA (2022) <https://bizwatchnigeria.ng/wba-reports-beer-sector-contributed-2-2bn-to-nigerian-economy>,

Calo J. R., P.G. Crandall, C.A. O'Bryan and S.C. Ricke (2015). Essential oils as antimicrobials in food systems - A review, Food Control 54, 111-119.

Dieckmann, R. H., &Palamand, S. R. (1974). Autoxidation of some constituents of hops. I. Monoterpene hydrocarbon, myrcene. *Journal of Agricultural and Food Chemistry*, 22(3), 498-503.

Ezea VN, Umerie SC and Ubaoji KI 2019. Chemical modifications of cellulose from palm kernel de-oiled cake to microcrystalline cellulose and its evaluation as a pharmaceutical excipient. *A J Pure Appl Chem*, 13(4), 49-57.

Figueredo G., J.C. Chalchat, S. Petrovic, Z. Maksimovic, M. Gorunovic, P. Boza and J. Radic (2009). Composition of essential oils of flowers, leaves, stems and rhizome of *Peucedanum officinale* L.(Apiaceae), *J. Essent. Oil Res.* 21, 123-126.

Guillen, M.D.; Cabo, N. Infrared spectroscopy in the study of edible oils and fats. *J. Sci. Food Agric.* 1997,75, 1–11.34.

Guillén, M.D.; Cabo, N. Fourier transform infrared spectra data versus peroxide and anisidine values to determine oxidative stability of edible oils. *Food Chem.* 2002, 77, 503–510.35.

6115

Hood EE. Plant-based biofuels. *F1000Res.* 2016 Feb 17;5:F1000 Faculty Rev-185. doi: 10.12688/f1000research.7418.1. PMID: 26949525; PMCID: PMC4758376.

Jaimand K, E. Sharifi and M. Dini (2006). Chemical constituents of the leaf and seed oils of *Peucedanum officinale* L. cultivated in Iran, *J. Essent. Oil Res.* 18, 670-671.

Jegannathan K.R, P. H. Nielsen (2013) Environmental assessment of enzyme use in industrial production – a literature review, *J Cleaner Production*, 42:228-240. <https://doi.org/10.1016/j.jclepro.2012.11.005>

Koutsaviti A, V. Antonopoulou, A. Vlasi, S. Antonatos, A. Michaelakis, D.P. Papachristos and O.Tzakou (2018). Chemical composition and fumigant activity of essential oils from six plant families against *Sitophilus oryzae* (Col: Curculionidae), *J. Pest Sci.* 91, 873-886.

Manion C. R. and R.M. Widder (2017). Essentials of essential oils, *Am. J. Health-Syst. Pharm.* 74, E153-E162.

Okafor V. N., Ifeyinwa Blessing Tabugbo, Regina Igwe Anyalebechi, Ugochukwu Wilson Okafor, Joy Ngozika Obiefuna (2020a) A Review of Nigerian Potential Hop Substitutes in Beer Brewing: 1983 – 2020. *International Research Journal of Pure and Applied Chemistry*, 21(15), 50 – 73. <https://doi.org/10.9734/IRJPAC/2020/v21i1530252>.

Okafor V. N., Regina Igwe Anyalebechi, Ugochukwu Wilson Okafor, Chinwe Priscilla Okonkwo, Joy Ngozika Obiefuna, Matthew Chiemezie Obiadi (2020b) Phytochemical Constituents of Extracts of Hops and Some Potential Nigerian Hop Substitutes: A Comparative Study in Beer Brewing. *International Journal of Biological and Chemical Research*, 1 (1), 1-7, July, 2020, (Science Assist International)

Okafor, V. N.; Okafor, U. W.; Anyalebechi, R. I. (2020c) Gas Chromatography-Mass Spectrometry (GC-MS) Profiling of Compounds of Hops and Some Nigerian Potential Hop Substitutes: Comparative Studies in Beer Brewing. *Journal of Chromatography and Separation Techniques*, 11 (2), No. 429, 1-9. <https://doi.org/10.35248/2157-7064.20.11.429>

Okafor V. N., U. B. Uche, R. C. Abailim (2020d) Levels of Polycyclic Aromatic Hydrocarbons (PAHs) in Beers: Consumption and Public Health Concerns: *Chemical Sciences International Journal*, 29 (1), 47-59, February, 2020, <https://doi.org/10.9734/CSJI/2020/v29i130157>.

Okafor V. N., Ugochukwu Wilson Okafor, Stella Nwakaego Ezem (2017a). Metabolites Composition of *Garcinia kola* Extract as Potential Substitutes for Isomerised Hop Extract in Beer Brewing: *Chemistry Research Journal*, 2(5), 153-162.

Okafor V. N and Obodoeze J. J (2017b) Comparative Evaluation of Phytochemical Constituents, Bitterness Characters and Essential Oil Contents of Extracts from Four Nigerian Plants as Potential Substitutes for Isomerized Hop Extract in Beer Brewing: *Chemistry Research Journal*, 2 (4),16-21.

Okafor V. N (2016a). Hops and Potential Nigerian Substitutes: Comparative Studies in Beer Brewing, LAP, Lambert Academic Publishing- GmbH & Co. KG, Germany. ISBN-10 (97833330014817)

Okafor V. N (2016b) Spectroscopic Methods for Evaluation of Hop Extracts and Extract from *Gongronemalatifolium* as Substitute in the Nigerian Beer Industry: *International Journal of Advanced Research* 4 (12), 2499-2504. <http://dx.doi.org/10.21474/IJAR01/2675>

Eboatu A. N., Emmanuel Nnamdi Ejike, Vincent Nwalieji Okafor. Evaluation of Anti-Nutritional Factors in Hop Extracts and Extract from *Azadirachta indica* as Potential Substitute in the Nigerian Beer Industry: EPH-International Journal of Applied Sciences 2 (11), 1-14, October, 2016. <https://doi.org/10.53555/ephas.v2i11.46>

Okafor V. N., Augustine NnalueEboatu, Regina Igwe Anyalebechi, Ugochukwu Wilson Okafor (2016c) Comparative Studies of the Physicochemical Properties of Beers Brewed with Hop Extracts and Extracts from Four Selected Tropical plants: Journal of Advanced Chemical Sciences 2 (4), 282-286.

Okafor V. N., Regina Igwe Anyalebechi, Ugochukwu Wilson Okafor (2016d). Characterization of Hop Extracts and Extracts from some Selected Nigerian Plants by GC/MS and Brewing Qualities Analyses: IJRDO-Journal of Applied Science 2 (9), 39-64. <https://doi.org/10.53555/as.v2i9.1404>

Rodolfi M, Valentoni A, Pretti L, Sanna M, Guidotti S, Marchioni I, Ganino T. From Hop to Beer: Influence of Different Organic Foliar Fertilisation Treatments on Hop Oil Profile and Derived Beers' Flavour. *Plants*. 2023; 12(9):1861. <https://doi.org/10.3390/plants12091861>

Rohman, A.; Che Man, Y.B. Monitoring of virgin coconut oil (VCO) adulteration with palm oil using Fourier transform infrared spectroscopy. *J. Food Lipids* 2009, 16, 618–628

Sarkhail P (2014). Traditional uses, phytochemistry and pharmacological properties of the genus *Peucedanum*: A review, *J. Ethnopharmacol.* 156, 235-270

Sofowora A, Ogunbodede E, Onayade A. The role and place of medicinal plants in the strategies for disease prevention. *Afr J Tradit Complement Altern Med.* 2013 Aug 12;10(5):210-29. doi: 10.4314/ajtcam.v10i5.2. PMID: 24311829; PMCID: PMC3847409.

Tapp, H.S.; Defernez, M.; Kemsley, E.K. FTIR spectroscopy and multivariate analysis can distinguish the geographic origin of extra virgin olive oils. *J. Agric. Food Chem.* 2003, 51, 6110–

Yadav S, Kanha Singh Tiwari, Chitrasen Gupta, Mahendra Kumar Tiwari, Arbaj Khan, Sankatha P. Sonkar (2023) A brief review on natural dyes, pigments: Recent advances and future perspectives, *Results in Chemistry*, 5: 100733, <https://doi.org/10.1016/j.rechem.2022.100733>.

Zhang H. M., S.S. Guo, B. Fan, S.S. Du, Y.Y. Wang and Z.W. Deng (2019). Evaluation of efficacy of the essential oil from *Ostericumviridiflorum* (Turcz.) Kitagawa in control of stored product insects, *Environ.Sci. Pollut. Res.* 26, 1406-1413

UNDER PEER REVIEW