

Original Research Article

Prevalence of Intestinal Parasites in Children Accessing Health Care in Federal Medical Centre Keffi, Nasarawa State, Nigeria.

- Comment [WU1]: Among
- Comment [WU2]: Medical Healthcare
- Comment [WU3]: At the Federal Medical Centre Keffi

ABSTRACT

Aim: This work was conducted between April 2022 and March 2023 to investigate the prevalence and predisposing factors of enteric parasites among children accessing health care in Federal Medical Center keffi, Nasarawa State.

- Comment [WU4]: Between the period of April 2022 and March 2023
- Comment [WU5]: Medical healthcare
- Comment [WU6]: At the

Study Design: The study was a cross-sectional study.

Place and Duration of Study: This research was carried out in Federal Medical Center Keffi between April 2022 and March 2023.

- Comment [WU7]: Delete

Methodology: Fresh stool samples were collected into sterile bottles from children aged 1-15 years (246 males and 194 females) who accessed health care in Federal Medical Center keffi, Nasarawa State from April 2022 and March 2023. Useful information such as age, gender, and sources of drinking water was obtained from the subjects through the administration of a structured questionnaire. The stool specimen was observed visually for colour, consistency, and presence of blood, pus, and adult worms while the concentration of the eggs, cysts, and larvae of the intestinal parasites was carried out using the formol ether method and viewed with X10 and X40 magnifications of the binocular microscope. The data obtained from this study were analyzed using the chi-square test by the use of Statistical Package for Social Sciences (SPSS) version (21.0). Values obtained were considered significant at 95% probability.

- Comment [WU8]: Medical healthcare
- Comment [WU9]: At the
- Comment [WU10]: Keffi
- Comment [WU11]: To
- Comment [WU12]: Biographic
- Comment [WU13]: Were
- Comment [WU14]: Specimens
- Comment [WU15]: Were
- Comment [WU16]: were
- Comment [WU17]: Data obtained
- Comment [WU18]: V
- Comment [WU19]: Review this values

Results: An overall prevalence of 27.26% was obtained in the study participants with a total of five species of intestinal parasites identified. The parasites include; *Entamoeba coli*, *Entamoeba histolytica*, *Giardia lamblia*, *Strongyloides stercoralis*, and *Trichomonas hominis*. *Entamoeba histolytica* was the most prevalent (15.45%) while *S. stercoralis* was the least prevalent (1.3.6%). In this study, males recorded a higher prevalence of *Entamoeba. histolytica* (16.49%) than females (14.63%) likewise for *E. coli* (5.14%) and (4.07%), *Strongyloides stercoralis* (1.55%) and (1.22%) respectively. In contrast, females recorded a higher prevalence of *Giardia lamblia* (2.85%) than males (2.58%) and *Trichomonas hominis* (3.25%) and (1.03%) respectively. The prevalence of intestinal parasites varied significantly among the age groups, with the age group 1-5 years being the most susceptible to *E. histolytica* (45.71%). Children who drank river water had the highest prevalence of parasitic infection (57.14%) while those who drank borehole water had the lowest majority (14.29%).

- Comment [WU20]: Your statistical analysis is not showing in your results

Conclusion: The high prevalence of intestinal parasites observed in this study in relation to the sources of drinking water of the subjects suggests a major predisposing factor to intestinal

parasitic infections. Therefore, the provision of potable drinking water and public enlightenment on proper hygiene practices in the study area is of great importance.

Keywords: Intestinal parasites, children, prevalence, sources of drinking water.

1. INTRODUCTION

Enteric protozoan infections are a leading cause of morbidity and mortality in children, especially in developing countries such as Nigeria [1, 2]. Having knowledge of the distribution of enteric parasitic diseases, as well as the areas of overlap, is critical for identifying hotspots where there is a need for consolidated prevention and control interventions [3-5]. In underdeveloped areas, these actions are particularly important as social determinants such as deficiencies in sanitation, poor personal hygiene and human cohabitation with domestic animals, favour the maintenance of infections, reinfections and coinfections [6]. These resultant infections can lead to impairments in physical and cognitive development, and eventually death [7].

Infections are mainly established when infective stage(s) (cysts or eggs/larvae) of the protozoa are transmitted via ingestion of water, soil or food contaminated by faeces [8]. Enteric parasitic infection is one of the neglected tropical diseases (NTD) that thrive where there is poverty. Populations most affected are the poor, living in isolated rural areas, urban slums or crisis-prone areas [9,10]. Soil-transmitted helminth infections especially hookworm infection which causes childhood and maternal anaemia, result in the greatest disability, and the highest burden of neglected tropical diseases [11]. The resultant physiologic changes as a result of these infections include iron-deficiency anaemia, growth retardation in children, intestinal obstruction and some other physical and mental health problems [12,13].

The helminths *Trichuris trichiura*, *Ascaris lumbricoides* and the hookworms as well as the protozoa such as *Entamoeba histolytica* cause infection in 800, 1,400, 1,200 and 48 million people respectively worldwide [14,15]. Studies have shown that more than three billion people are infected with intestinal parasites with the prevalence being higher among children below six years old by varying degrees in countries around the globe [16], with children more susceptible and amounting to the most significant risk population and can contribute to malnutrition, especially in children in daycare centres and orphanages. This is due to ignorance, low levels of safety, direct contact and sharing toys with other children, absence of functional sanitary facilities and overcrowding [17]. An exemplary case is the Infections of *Giardia lamblia* which damages the intestinal mucosa and results in malabsorption of nutrients, particularly fat [18,19]. The high prevalence of intestinal parasites among children globally and especially in developing countries which Nigeria is one has prompted the need to investigate the prevalence in the study area.

2. Materials and Methods

Comment [WU21]: Delete

Comment [WU22]: Helminthes such as

Comment [WU23]: Please kindly state the prevalence of intestinal parasitic infections in Africa, Nigeria, Nasarawa State as well as Keffi.

2.1 Study Area

The study was carried out at Federal Medical Centre Keffi, Nasarawa Nigeria. Keffi is located approximately 68km from Abuja, the Federal Capital Territory and 128km from Lafia, the capital of Nasarawa State. It is located geographically between latitude 8°3'N of the equator and longitude 7°50'E and situated at an altitude of 850m above sea level [20].




Figure 1: Map of Nasarawa state with the study area highlighted in yellow.

2.2 Sample Size Determination

In this study, the total number of samples was estimated using the standard formula described by [21]

$$N = \frac{Z^2 pq}{D^2}$$

Where; N=sample size,

Z= standard normal distribution at 95% confidence interval=1.96,

p= prevalence rate of 54.55% recorded by [22].

q=1-p

D= the allowable error, which is taken as 5% =0.05

Substituting the values in the formula;

$$q=1-p (1- 0.55) =0.45$$

$$N = \frac{1.96^2 \times 0.55 \times 0.45}{0.0025} = \frac{0.950769}{0.0025} = 380 \quad N= 380$$

This represents the minimum number of samples to be collected

2.3 Inclusion Criteria

Only children between the ages of 1 and 15 years who have not taken any anti-parasitic drugs in recent past 3 weeks were enrolled for this study.

2.4 Exclusion Criteria

Children more than 15 years of age, those who are on anti-parasitic drugs, and those who did not consent were excluded from the study.

2.5 Samples and Data Collection

A total of 440 fresh stool samples were collected from participants who consented to sterile bottles from children aged 1-15 years (246 males and 194 females) who accessed health care in Federal Medical Center keffi, Nasarawa State between April 2022 and March 2023. Useful information such as age, gender, and sources of drinking water was obtained from the subjects through the administration of a structured questionnaire.

2.6 Examination of Stool Samples

Samples were collected in sterile universal containers, labeled properly, and taken to the Parasitology Unit, Medical Laboratory Services Federal Medical Centre Keffi. The direct wet mount and formol ether concentration technique described by [23] was used for the preparation of the stool samples. Thereafter, the stool preparation was transferred onto a clean slide, and a drop of iodine was added before covering it with a cover glass and observed microscopically using the X10 and X40 magnifications.

2.7 Data Analysis

The prevalence of intestinal parasitic infection among the examined children was calculated using the number of infected students divided by the number of children examined. The significance of the infection among the children was tested by Chi-square (X^2). Values obtained were considered significant at 95% probability using the statistical package for social sciences (SPSS) version 21.0

Comment [WU24]: Who are assessing medical healthcare

Comment [WU25]: At the

Comment [WU26]: Biographic

Comment [WU27]: Delete

Comment [WU28]: Direct

Comment [WU29]: Delete

Comment [WU30]: Prevalence

Comment [WU31]: Are you working with students or children?

3. RESULTS

3.1 Identified Intestinal Parasites and their Prevalence

Of the 440 children examined in this study, 122 (27.26%) were infected with intestinal parasites. The parasites detected include; *Entamoeba histolytica* (15.45%), *Entamoeba coli* (4.55%), *Giardia lamblia* (2.27%), *Trichomonas hominis* (2.27%) while *Strongyloides stercoralis* and cases of co-infection were (1.36%) respectively as shown in Table 1.

Comment [WU32]: Of the total of 440

Comment [WU33]: Delete

Table 1: Overall Prevalence of Intestinal Parasites among Children attending Federal Medical Centre Keffi

Intestinal Parasites	No. Positive	Prevalence (%)	P. value
<i>Entamoeba coli</i>	20	4.55%	
<i>Entamoeba histolytica</i>	68	15.45%	
<i>Giardia lamblia</i>	12	2.27%	0.000
<i>Strongyloides stercoralis</i>	6	1.36%	
<i>Trichomonas hominis</i>	10	2.27%	
Co-infection	6	1.36%	

Comment [WU34]: Include another row for the totals

3.2 Overall Prevalence of Intestinal Parasites with Respect to Some Socio-demographic Variables

The result in Table 2 shows the prevalence of intestinal parasites among the children examined in relation to age and gender and sources of drinking water. Children aged between 1-5 had a higher prevalence (65.71%) followed by those aged 6-10 (19.59%) while those aged 11-15 had the least prevalence (18.18%). Females had a higher prevalence of 31.96% than males 21.95%. the prevalence in relation to sources of drinking water was higher among children who drank river water (57.14%) followed by those who drank rain water (35.00%) while the least prevalence was seen among children who drank borehole water (14.29%).

Comment [WU35]: Delete

Comment [WU36]: Table showed

Table 2. Prevalence of intestinal parasites among children attending Federal Medical Centre Keffi with regards to Sociodemographic variables

Age Groups (Years)	Number Examined	Number Negative	Number Positive	Prevalence	P-value
1-5	70	24	46	65.71%	
6-10	194	156	38	19.59%	0.000
11-15	176	144	32	18.18%	
Gender					
Males	246	192	54	21.95%	0.018
Females	194	132	62	31.96%	
Sources of Drinking Water					
Well	160	117	43	26.88%	
Tap	122	99	23	18.85%	
Borehole	63	54	9	14.29%	0.001

River	35	15	15	57.14%
Rain	60	39	21	35.00%

Comment [WU37]: Include another row for totals

4.0 DISCUSSION

In this study which was based on the prevalence of intestinal parasites in children accessing health care in Federal Medical Centre (FMC) Keffi, Nasarawa State Nigeria, the overall prevalence of intestinal parasites was 27.62%. Enteric parasites detected in this study include *Entamoeba coli* (4.55%), *Entamoeba histolytica* (15.45%), *Giardia lamblia* (2.27%), *Strongyloides stercoralis* (1.36%), *Trichomonas hominis* (2.27%) with $P=0.05$. Also, co-infection was observed in this study with a prevalence of 1.36%. *E. histolytica* had the highest prevalence of 15.45% while *S. stercoralis* had the least prevalence of 1.36%. These intestinal parasites have been reported in several parts of Nigeria including [10] in Kaduna metropolis, [24] in Gwagwalada, Abuja, [25] in Benue, [26] in Delta State and in Ilorin by [27]. It appears that the detection of these intestinal parasites among school-age children is primarily due to low standards of personal hygiene, poor sanitary conditions, poverty, lack of toilet facilities, and lack of portable drinking water [28].

Comment [WU38]: This study was conducted to determine the

Comment [WU39]: Among

Comment [WU40]: .

Comment [WU41]: The

The overall prevalence of 27.26% obtained in this study is in agreement with a prevalence of 27.2% reported in some communities in Akwanga local Government area of Nasarawa State by [29]. Similarly, [1] reported an overall prevalence of 27.66% in Rivers State Nigeria. Also, [30] reported a prevalence of 27.3% in Katakao area, North-Central Nigeria. The prevalence obtained in this study is higher than the 23.32% and 23.6% reported by [27,31] respectively, both in Ilorin, Kwara State Nigeria, and a prevalence of 16.2% reported in Okpokwu, Benue State Nigeria by [32], also [30] reported a prevalence of 21.2% among preschool children in Kushe, North-Central Nigeria. In a study carried out in Southern Nigeria, [33] reported a prevalence of 23.95% among primary school-aged children. Elsewhere, a prevalence of 22.43% was reported by [34] in Egypt, 24.8% reported by [35] in Sudan, and a prevalence of 22.2% was reported by [17] in India. On the contrary, a very low prevalence of 5.9% was reported in Europe over a period of five years by [36]. A slightly higher prevalence of 29.0% was reported by [37] among selected age groups within Okada South-South Nigeria while a prevalence of 30% was reported by [38] in Abia State Nigeria. Similarly, a prevalence of 34.2% was reported among primary school children in three geo-political zones of Imo State Nigeria by [12] and 38.6% among public primary school children in Ibadan Oyo Nigeria by [39]. Generally, the result obtained in this study is low compared to what was obtained in other local government areas of Nasarawa State. The result obtained by [22] in two local Government areas of Nasarawa state gives a prevalence of 54.55%. Similarly, a prevalence of 47.5% was reported by [40] (2019) in Lafia, Nasarawa State, also a prevalence of 33.5% among school-age children in Lafia was reported by [41]. In another study carried out in Karu, Nasarawa State by [42], 45.1% was reported among school children in two communities of Auta balefi. Studies from other parts of the country with higher prevalence have also been reported. The difference in prevalence could be as a result of seasonal variation in sample collection, processing and the overall differences in living standards across study areas.

Comment [WU42]: Local

Comment [WU43]: Similarly

With respect to gender difference, this study shows a prevalence of 32.99% among females while 21.95% was recorded for males ($P=0.05$). This is in agreement with reports by [43] in Minna Niger State, Nigeria, who reported that females had a higher rate of infection at 94% while males were 51.4%. Also, [17] in India reported that females had a prevalence of 56% while males were 43.9%. On the contrary, the prevalence of intestinal parasites was 29% among males while it was 20.9% among females reported by [44], [38] in Abia State Nigeria, reported 42.86% for males and 23.08% for females. [30] reported a prevalence of 57.60%, in males while 42.40% was reported for females in North-Central Nigeria. [45] in Egypt reported a prevalence of 58% for males and 42% for females while [46] reported a prevalence of 64.8% among males and 35.1% in females in Pakistan.

This study showed the highest prevalence of enteric parasites among children aged 1-5 (65.71%) followed by those aged 6-10 (19.59%) and those aged 11-15 (18.18%) respectively with $P=0.05$. A similar case was reported by [26] with children aged 5-7 having a prevalence of 54.22% while those aged between 14-16 years had a prevalence of 42.86%. Similarly, rate of infection was high among children of 6-8 (77.63%) and 9-11 (85.00%) years of age compared to those aged 12-14 (53.13%). In Abia State Nigeria, [38] reported a highest prevalence rate in age group of 5-8 years 24 (32.43%) whereas 9-12 years age group had the lowest prevalence rate of 24 (27.91%). [47] in North Eastern Nigeria reported a prevalence of 85.7% among children aged 6-8, while the least prevalence of 77.7% was reported in the 13-16 years age bracket. Furthermore, [24] in Gwagwalada, FCT, Abuja Nigeria reported a prevalence of 28(43.1%) among children aged 4 to 5 years. Also, [35] reported the highest prevalence rate (35%) among the 6-8 years age groups, while the lowest prevalence rate (10.3%) was reported among 12-14 years age groups. A high prevalence rate of 54.22% among the age group 5-7 years while the age group 14-16 years recorded the lowest prevalence of 42.86% as reported by [26] among school children in Deta State Nigeria. Conversely, [46] reported a higher prevalence among children aged 10-12 (94.2%) compared to those aged 4-6 (72%), while [45], reported an overall prevalence of 32% among children aged 11-20 years. Studies have shown that children in these age groups often spend more of their leisure time outdoors, playing both in school and at home, and are fond of picking food from the floor and touching unclean surfaces that end up in their mouths. They are also more often in contact with soil and eat indiscriminately with unwashed hands [34]. Also, the lack of sanitation facilities or non-functional sanitation facilities may be a factor contributing to the high prevalence of intestinal parasites among these age groups [48].

The relationship between the overall detection of enteric parasites and the source of drinking water was exploited in this study with $P=0.05$. For well water it was 26.88%, tap water: 18.85% borehole: 14.29%, river: 57.14%, and rain water was 35.00% respectively. The highest prevalence was recorded among children who used river water (57.14%) while the least was recorded among those who used borehole water (14.29%). Reports within and outside Nigeria show varying prevalence with regard to sources of drinking water. [27] reported a prevalence of 45% among children who drank river water in Kwara, Nigeria, similarly, [49] in Katsina-ala, Benue State reported a prevalence of 29.6% among children who obtained their drinking water from streams, 45.5% from wells, 37.8% from rivers, 26.1% from taps and 0.0% from water vendors. In other parts of Africa, [50] from Kenya reported that children who used rainwater for drinking were more likely to have intestinal protozoan parasite infections than those who used tap water, bought water from vendors, or used water from other sources as not being significant. However, [51] reported in Ethiopia that drinking water from river sources was rated as the highest associated risk with 36.8% of parasitic infections. Also in Ethiopia, [52] reported a

significant relationship between sources of drinking water and the detection of parasites. Rainfall and agricultural residues that transfer parasites to drinking water sources contribute to the spread of the parasites and in turn the high prevalence [53].

5.0 CONCLUSION

A prevalence of 27.26 % obtained in this study shows a high presence of intestinal parasites in the study area, the prevalence was found to be higher in females than in males. Children between ages 1-5 years had a higher prevalence of infection compared to others. This could be a result of the propensity of children within this age range to put their dirty hands in their mouths and also due to picking dirt from the ground. Intestinal parasitic infections in the children were found to be significantly associated with fecal oral route transmission via unhygienic water.

CONSENT

Written informed consent was taken from each participant.

ETHICAL APPROVAL

The ethical approval for this research was obtained from the Health Research Ethics Committee, Federal Medical Centre Keffi (NHREC/2012/2012).

REFERENCES

1. Abah, A. E. and Arene, F. O. I. Status of Intestinal Parasitic Infections among Primary School Children in Rivers State, Nigeria. Hindawi Publishing Corporation Journal of Parasitology Research Volume 2015, Article ID937096, <https://doi.org/10.1155/2015/937096>
2. Ugboko, H. U., Nwinyi, O. C., Oranusi, S. U., & Oyewale, J. O. Childhood diarrhoeal diseases in developing countries. *Heliyon*, 2020;6(4), e03690. <https://doi.org/10.1016/j.heliyon.2020.e03690>
3. Akinbo, F. O., Omoregie, R., Eromwon, R., Igbenimah, I. O., & Airueghiomon, U. E. Prevalence of intestinal parasites among patients of a tertiary hospital in Benin city, Nigeria. *North American journal of medical sciences*, (2011). 3(10), 462–464. <https://doi.org/10.4297/najms.2011.3462>
4. Bergquist R., Yang G.J., Knopp S., Utzinger J., Tanner M. Surveillance and response: Tools and approaches for the elimination stage of neglected tropical diseases. *Act. Trop.* 2015;141:229–234. doi: 10.1016/j.actatropica.2014.09.017.

5. Tambo E., Ai L., Zhou X., Chen J.H., Hu W., Bergquist R., Guo J.G., Utzinger J., Tanner M., Zhou X.N. Surveillance-response systems: The key to elimination of tropical diseases. *Infect. Dis. Poverty*. 2014;5:49:1–49:12. doi: 10.1186/2049-9957-3-17
6. Dantas-Torres, F., & Otranto, D. Best Practices for Preventing Vector-Borne Diseases in Dogs and Humans. *Trends in parasitology*, (2016). 32(1), 43–55. <https://doi.org/10.1016/j.pt.2015.09.004>
7. Rodríguez, L., Cervantes, E., & Ortiz, R. Malnutrition and gastrointestinal and respiratory infections in children: a public health problem. *International journal of environmental research and public health*, (2011). 8(4), 1174–1205. <https://doi.org/10.3390/ijerph8041174>
8. Mekonnen et al. Assessment of seasonality in soil-transmitted helminth infections across 14 schools in Jimma Town, Ethiopia. *Pan African Medical Journal*. (2019).;32:6. doi: [10.11604/pamj.2019.32.6.16085](https://doi.org/10.11604/pamj.2019.32.6.16085)
9. Bangert, M., Molyneux, D.H., Lindsay, S.W. *et al.* The cross-cutting contribution of the end of neglected tropical diseases to the sustainable development goals. *Infect Dis Poverty* (2017). 6, 73. <https://doi.org/10.1186/s40249-017-0288-0>
10. Hadiza M.K., Maikaje D.B. and Ijah U.J.J. Prevalence of Intestinal Parasites Among Children Attending Daycare and Orphanage Centers in Kaduna Metropolis, Kaduna. *Science World Journal* (2019) Vol. 14(No 3).
11. Hotez PJ, Alvarado M, Basáñez M-G, Bolliger I, Bourne R, Boussinesq M, et al. The Global Burden of Disease Study 2010: Interpretation and Implications for the Neglected Tropical Diseases. *PLoS Negl Trop Dis* (2014) 8(7): e2865. <https://doi.org/10.1371/journal.pntd.0002865>
12. Udensi. U.J., Ifenyinwa, M. C., Ijeoma, E. N., Godson, U. M., & Nwaku, A. I. Prevalence of intestinal parasites among primary school children in three geopolitical zones of imo state, Nigeria. *Science Journal of public health*, (2015). 3(5-1), 25-28.
13. Ihejirika, O. C., Nwaorgu, O. C., Ebirim, C. I., & Nwokeji, C. M. Effects of intestinal parasitic infections on nutritional status of primary children in Imo State Nigeria. *The Pan African medical journal*, (2019).33,34. <https://doi.org/10.11604/pamj.2019.33.34.17099>
14. Houmsou, R, E Amuta, T Olusi. Prevalence of intestinal parasites among primary school children in Makurdi, Benue State Nigeria. *The Internet Journal of Infectious Diseases*. 2009 Volume 8 Number 1.
15. Arora, D. R. and Arora, B. B. *Medical Parasitology* 3rd Edition. CBS Publishers and Distributors PVT Limited, New Delhi, India. (2011). PP 3-195.
16. Anvari Tafti M H, Mirjalili M M, Aghabagheri M. The Prevalence of Intestinal Parasites in Children Attending Day-Care Centers in Yazd City, Iran. *JCHR*; (2014). 3 (2) :96-102 URL: <http://jhr.ssu.ac.ir/article-1-189-en.html>
17. Kumar Manochitra, Shashiraja Padukone, Selvaratthinam Ajay Philips and Subhash Chandra Parija. Prevalence of Intestinal Parasites among Patients attending a Tertiary Care Centre in South India. *International Journal of Current Microbiology and Applied Sciences* ISSN: 2319-7706 Volume 5 Number 9 (2016).
18. Bartelt, L. A., & Sartor, R. B. Advances in understanding Giardia: determinants and mechanisms of chronic sequelae. *F1000prime reports*, (2015). 7, 62. <https://doi.org/10.12703/P7-62>

19. Solaymani-Mohammadi S. Mucosal Defense Against *Giardia* at the Intestinal Epithelial Cell Interface. *Frontiers in Immunology*, (2022). 13, 817468. <https://doi.org/10.3389/fimmu.2022.817468>
20. Akwa VL. Binbol NL. Samaila KL. Marcus ND. Geographical perspective of Nasarawa State, Onaive Printing and Publishing Company Ltd, Keffi, 2007; p 503
21. Naing L, Winn T. Rusli BN. Practical issues in calculating the sample size for prevalence studies. *Archives of Orofacial Sciences*. 2006;1:9-14
22. Ombugadu, R.J., Makpo, J.K., BanyingI, H.A. and Eke, S.S. Intestinal parasitic infection with special reference to Entamoeba histolytica in two local government areas of Nasarawa State, Nigeria. (2010) VL - 8DO - 10.4314/tzool.v8i1.66696JO - Zoologist
23. Cheesbrough M. District laboratory practice in tropical countries Part 1. Second edition. Cambridge University press, USA, 2009; p 191-204
24. Gimba, U.N. and Dawam, N.N. Epidemiological Status of Intestinal Parasitic Infection Rates in Children Attending Gwagwalada Township Clinic, Fct-Abuja, Nigeria. *American Journal of Research Communication*, (2015) 3(2): 97-110
25. Tyoalumun Kpurkpur, Sani Abubakar, and Nongu Christopher. Prevalence of Intestinal Parasitic Infections and their Association with Nutritional Status of Rural and Urban Pre-School Children in Benue State, Nigeria. *International Journal of MCH and AIDS* (2016), Volume 5, Issue 2, 146-152.
26. Obioma Jacinta Eboh, Christopher E. Okaka & Thankgod Onuoha. Prevalence of Gastrointestinal Parasites among School Children in Delta State, Nigeria. *London Journal of Research in Science: Natural and Formal* (2022) Volume 22 | Issue 8 | Compilation 1.0
27. Blessing Olawunmi Amisu, Olalekan John Okesanya, Noah Olabode Olaleke, Celina Omolade Ologun, Don Eliseo Lucero-Prisno III, Victoria Oluwafunmilayo Ogunwale, Racheal Ayeni Ahuoyiza, Emery Manirambona, Bijaya Kumar Padhi, Abhishek Mewara (2023) Socio-Environmental Determinants of Parasitic Intestinal Infections Among Children: A Cross-Sectional Study in Nigeria. *Journal of Global Health Science*. Jun;5(1): e6
28. Nxasana N, Baba K, Bhat V, Vasaikar S. Prevalence of intestinal parasites in primary school children of mthatha, eastern cape province, South Africa. *Ann Med Health Sci Res*. 2013 Oct;3(4):511-6. doi: 10.4103/2141-9248.122064. PMID: 24380000; PMCID: PMC3868115.
29. Ali AA, Pam VA, Uzoigwe NR, Ombugadu A and Maikenti JI. Prevalence of Gastrointestinal Infections Among Human Population in Some Communities in Akwanga Local Government Area, Nasarawa State, Nigeria. *Trends Tech Sci Res*. 2023; 5(5): 555674. DOI: 10.19080/TTSR.2023.05.555674
30. Magaji, P. J., & Magaju, J. The prevalence of gastrointestinal parasites among primary school children in Kagarko local government area, Kaduna State, Nigeria. *American Journal of Health, Medicine and Nursing Practice*, (2021). 6(1), 1-17. doi:10.47672/ajhmn.646
31. Shuaibu, I., Umar, Y. A., & Chanding, A. Y. Prevalence and Associated Risk Factors of Gastro-Intestinal Parasites among Primary School Pupils in Ilorin East Local

- Government Area, Kwara State, Nigeria. *Researchers World*, (2018). 9(4), 137-144. DOI : 10.18843/rwjasc/v9i4/16
32. Ogbeyi, G., Jenewari, J., Abah, J., Akor, J., & Ikpom, E. Current Status of Intestinal Parasites among Pre School Children in Eke, Okpokwu Local Government Area, Benue State, Nigeria. *Western Journal of Medical and Biomedical Sciences*, (2020). 1(2), 150–157. <https://doi.org/10.46912/wjmbs.28>
 33. Gbonhinbor J, Abah AE, Awi-Waadu G. Prevalence of Intestinal Parasitic Infection and Associated Risk Factors Among Primary School-Aged Children (5-15 years) in Southern Nigeria. *Int Journal of Infection* 2022; 9(3)e123721. doi:10.5812/iji-123721
 34. Ahmed, Wafaa F.A. Intestinal Parasites Among Primary School Children in Urban and Rural Tanta, Gharbia, Governorate, Egypt. *Egypt. J. Exp. Biol. (Zool.)*, (2013) 9(2): 257 – 262.
 35. Abuobieda Sirelkhatem Mohammed Elameen¹, Mosab Nouralde in Mohammed Hamad, Mohammed Baha Eldin Ahmed. Prevalence of Intestinal Parasitic Infections among Primary Schools aged Children in Ombda Locality. *Saudi Journal of Biomedical Research*, Dec., 2019; 4(12): 412-415
 36. Maria, K., Karalexi, M. A., Vrioni, G., & Tsakris, A. Prevalence of Intestinal Parasitic Infections among Children in Europe over the Last Five Years. *Tropical medicine and infectious disease*, (2021). 6(3), 160. <https://doi.org/10.3390/tropicalmed6030160>.
 37. Okafor-Elenwo, E. J., Izevbuwa, E.O., & Akpoka, O. A. Prevalence of Intestinal Parasites amongst selected age groups within Okada, South –South Nigeria. *International Journal of Applied Biology*, (2020). 4(1), <https://doi.org/10.20956/ijab.v4i1.9549>.
 38. Ithemanma C.A, Oladele O.T. Prevalence of Intestinal Parasites and Associated Risk Factors among School Children (5 - 16 Years) in Ihite-Ude, Ofeme Community in Umuahia North L.G.A, Abia State. *International Journal of Research & Review* (www.gkpublication.in) Vol.2; Issue: 12; December 2015, pp732-738.
 39. Nnolim, C., Adekeye, T. A., & Awobode, H. O. Intestinal helminth infection and malnutrition among public primary school children in Ibadan, Nigeria. *Nigerian Journal of Parasitology*, 2020;41(1).
 40. Tongjura, J.D.C. & Ombugadu, J.R. & Abdullahi, M.M. & Blessing, M.A. & Amuga, G.A. & Mafuyai, Hayward. Intestinal parasites amongst primary school children attending Ta'al Model Primary School in Lafia Local Government Area of Nasarawa State, Nigeria. *Nigerian Journal of Parasitology*.2019; 40. 92. 10.4314/njpar.v40i1.15.
 41. Eniola, M. A., Echeta, O. C., Ombugadu, A., Ajah, L., Aimankhu, P. O., & Oluwole, A. S. Helminthiasis among School-Age Children and Hygiene Conditions of Selected Schools in Lafia, Nasarawa State, Nigeria. *Tropical medicine and infectious disease*, 2019;4(3), 112. <https://doi.org/10.3390/tropicalmed4030112>
 42. Ishaku, M. J., Onyeacho, C. P., & Koggie, A. Z. Prevalence of Gastrointestinal Helminth Parasites among School Children Attending Two Community Schools in Auta Balefi, Karu, Nasarawa State. *Annual Research & Review in Biology*, 2020; 35(2), 96-106. <https://doi.org/10.9734/arrb/2020/v35i230193>

43. Eke, S. S., **Omalu**, I. C. J., Ochayi, Q., Pam, V. A., Otuu, C. A., Ibeh, E. O., Hassan, S. C. and Garba, Y. Comparative Study on The Prevalence of Intestinal Parasites Among Primary School Children in Tudun-Fulani Model School and Hasha International School Minna, Niger State. *NSUK Journal of Science & Technology*, Vol. 6: No. 2. 2016. pp. 80 - 85 ISSN: 1597- 5527.
44. Ismail, K.A. Prevalence of Intestinal Parasitic Infection among School Children in Taif. *Insights Biomed.* (2018) Vol.3 No.2:10
45. Marwa Omar and Heba O. Abdelal. Current status of intestinal parasitosis among patients attending teaching hospitals in Zagazig district, Northeastern Egypt. *Parasitology Research* (2022) 121:1651–1662
46. Ulhaqa, Z, W. Khanb, M. F. Khana, M. Kabirc, A. A. Ujjand, W. Ullaha, Z. Masoode, S. Khana and P. De los Ríos Escalantef.g. Prevalence of intestinal parasitic diseases in school children of rural areas of district Lower Dir, Pakistan. *Brazilian Journal of Biology*, 2022, vol. 82, e243150
47. Damen, J.G., Luka, J., Biwan, E.I., Lugos, M. Prevalence of Intestinal Parasites among Pupils in Rural North Eastern, Nigeria. *Nigerian Medical Journal* Vol. 52 Issue1. (2011) Doi:10-4314/NMJ.V52i1.66872
48. Yoseph, A., & Beyene, H. The high prevalence of intestinal parasitic infections is associated with stunting among children aged 6-59 months in Boricha Woreda, Southern Ethiopia: a cross-sectional study. *BMC public health*, (2020); 20(1), 1270. <https://doi.org/10.1186/s12889-020-09377-y>
49. Agbo, Joseph & Hemen, Agere & Dawuda, Manah & Agatsa, David & Alfred, A. Prevalence of Intestinal Parasites Among Children in Selected Primary Schools in Katsina-Ala Local Government Area of Benue. (2019);4. 478-482.
50. Chege, N. M, Bartholomew N. Ondigo, Frank G. Onyambu, Alex Maiyo Kattam, Nancy Lagat, Tabitha Irungu, Elizabeth Jemaiyo Matey. The prevalence of intestinal parasites and associated risk factors in school-going children from informal settlements in Nakuru town, Kenya. *Malawi Medical Journal*. (2020) 32 (2): 80-86. <https://doi.org/10.4314/mmj.v32i2.5>
51. Zemene, T., Shiferaw, M.B. Prevalence of intestinal parasitic infections in children under the age of 5 years attending the Debre Birhan referral hospital, North Shoa, Ethiopia. *BMC Res Notes* (2018). 11, 58. <https://doi.org/10.1186/s13104-018-3166-3>
52. Dagne, N and Aleign, A. Prevalence of Intestinal Protozoan Parasites and Associated Risk Factors among School Children in Merhabete District, Central Ethiopia. *Journal of Parasitology Research* Volume, (2021). Article ID 9916456, 7 pages <https://doi.org/10.1155/2021/9916456>
53. Baqer, N. N., Hammood, A. H., Hassan, K. F., & Hassan, E. S. A. Detection of Water-Borne Parasites in Drinking Water of Baghdad, Iraq. *African journal of infectious diseases*, (2018). 12(2), 1–6. <https://doi.org/10.21010/ajid.v12i2.1>

Comment [WU44]: Omalu