

A Comprehensive Investigation of the Interactions between Proteins and Ligands in the Crystal Structures of Mycobacterium Tuberculosis

ABSTRACT

Mycobacterium tuberculosis (MTB), an acid-fast aerobic bacterium that may grow on gram stain as either a gram-positive or gram-negative bacterium, is the disease-causing agent of tuberculosis (TB). Rifampin, isoniazid, pyrazinamide, and ethambutol, the first-line anti-tubercular drugs, can all have hepatotoxic side effects. The new medicine needs to work through a novel mode of action or to a novel target, be more active than presently available treatments, and shorten the course of treatment for the MDR-TB and XDR-TB. And active against both active and latent bacteria, and not interact with antiretroviral medications because many TB patients also have HIV. Additionally, it must work well with other anti-TB medications in order to form at least an effective three-drug regimen. This article discusses the analysis of few FDA approved anti-tubercular medications and their binding locations with respective targeted proteins. This mainly focuses on the amino acids of the proteins which are responsible for the formation of interactions with a drug molecule. So researchers can modify the existing drugs or its derivatives or can construct a new molecule according to the binding sites of enzymes corresponding to mycobacterium tuberculosis.

Keywords: *Anti-tubercular drugs; mycobacterium tuberculosis; MDR-TB; XDR-TB.*

1. INTRODUCTION

TB is still among the deadliest infectious diseases in the world. Nearly 30,000 people become ill with this preventable and treatable disease every day, and close to 4400 people lose their lives to TB. Access to TB diagnosis and treatment as well as the burden of TB disease continue to be negatively impacted by the COVID-19 epidemic. Global TB objectives are off track and advancement made in the years leading up to 2019 has slowed, stopped, or reversed. The number of people who infected TB was expected to be 10.6 million (95% UI: 9.9-11 million) in 2021, an increase of 4.5% from 10.1 million (95% UI: 9.5-10.7 million) in 2020. The number of new cases of tuberculosis (TB) per 100,000 people increased by 3.6% between 2020 and 2021, reversing annual decreases of roughly 2% for the most of the preceding two decades. The net decrease from 2015 to 2021 was 10%, which is barely halfway to the first End TB Strategy milestone [1]. Reductions in number of new TB cases in 2020 and 2021 indicate that there are more people with the disease who are undiagnosed and untreated, which first leads to more TB deaths and community transmission of the infection before, after a lag, increasing the number of new cases of

TB. We anticipate that the impact of COVID-19 will vary depending on the context. For instance, nations with high rates of tuberculosis (TB), like India and Viet Nam, have seen substantially disparate COVID-19 occurrence [2, 3]. The Mycobacterium tuberculosis infections medications approved by the FDA include Rifampin, Isoniazid, Pyrazinamide, and Ethambutol[4]. Approximately 13 new compounds have been recently discovered, including Bedaquiline, Delamanid, Pretomanid, Contezolid, Delpazolid, Sutezolid, GSK-3036656, Macozinone, OPC-167832, Q203, SQ109, TBA-7371, and TBI-166. This occurs following almost 40 years of active research and development work without any novel approved TB drugs. Based on their phase II clinical trials, two of these drugs bedaquiline and delamanid have been given accelerated or conditional regulatory approval[5-7]. Linezolid, Levofloxacin, Moxifloxacin, and Nitrazoxanide are the additional substances. For better TB illness therapy, more research is required. Although the TB typically assaults the lungs, they can also affect the kidney, spine, and brain[8]. Droplets from speaking, coughing, and sneezing propagate the illness. Microbacterial cultures, chest X-rays, and tuberculin skin tests are used to make the diagnosis[9]. The appearance of tuberculosis varies, and based on clinical manifestation, it is categorized into pulmonary TB (PTB) and extrapulmonary TB (EPTB)[10]. Because extrapulmonary TB (EPTB) accounts for more than 50% of all TB diagnoses in HIV-positive individuals, the situation is made more difficult by the HIV pandemic[11]. EPTB frequently presents a significant challenge for early diagnosis due to its heterogeneity of presentation. Constitutional symptoms including fever, anorexia, weight loss, malaise, and exhaustion may be present[12]. Drug resistance is a recurring issue in the treatment of TB. For instance, in 2016, it was reported that there were 490,000 instances of multidrug-resistant tuberculosis (MDR-TB) and an additional 110,000 cases of TB that was susceptible to isoniazid but resistant to rifampicin (RR-TB), the best first-line anti-TB medication[13]. MDR-TB is resistant to two most potent anti-TB medications, rifampicin and isoniazid. Extensively drug-resistant tuberculosis (XDR-TB), also known as MDR-TB and is resistant to at least one fluoroquinolone plus a second-line injectable medication (kanamycin, capreomycin, or amikacin), is a different form of resistant TB. In the intensive phase, which lasts for 6 to 12 months, all XDR-TB cases are treated with injections of Capreomycin, Moxifloxacin (MIFX), PAS high dose Isoniazid, clofazimine, Linezolid, and Co-amoxiclav. The rest of the drugs, with the exception of the injections, are continued for 18 months in the continuation phase. The following obstacles must be overcome by researchers working in the field of TB treatment, among others: the newer drug must be more effective than currently available drugs in order to shorten the length of treatment; it must also be active against both active and latent bacteria; it must act by a novel mechanism of action or to a novel target, particularly for the treatment of MDR-TB and XDR-TB; and it must not interact with antiretroviral drugs because many TB patients also take these drugs[14,15]. In this review, 14-FDA approved Anti-tubercular drugs obtained from different literatures and their protein-ligand interactions are discussed.

2. FDA approved drugs used in TB

2.1 Bedaquiline

For the treatment of tuberculosis, bedaquiline, a diarylquinoline, is offered under the trade name Sirturo. Both the FDA and the EMA authorized bedaquiline for the treatment of MDR-TB in December 2012 and March 2014, respectively[16-20].

Mechanism of Action: Bedaquiline is extremely effective and works through a novel mechanism that involves inhibition of the proton pump of mycobacterial ATP synthase [21].

Binding Pockets: In their Nucleic Acid Research publication, Kailu Yang and Junjie Zhang reported that bedaquiline has hydrophobic connections with PHE-691, LEU-721, PRO-172, VAL-176, GLU-654, ILE-705, ALA-669, and TYR-681[22].

2.2 Delamanid

Delamanid, also marketed as Deltyba, is a derivative of nitrodihydroimidazooxazole (OPC-67683). On April 28, 2014, the EMA granted delamanid marketing authorization for use.

Mechanism of Action: Delamanid works by preventing the mycolic acid synthesis. A Keto-mycolic and methoxy-mycolic acid, two crucial elements of M. tuberculosis cell wall, is inhibited, which results in bacterial cell death [23].

Binding pockets: Cellitti S.E. came to the conclusion in their study that the Delamanid exhibits hydrophobic and other interactions with LYS-55A, ARG-60A, PRO-63A, LEU-64A, and MET-87A, VAL-46A, LYS-55A, THR-56A, ASN-62A, PRO-63A, TYR-65A, ALA-76A, LYS-79A, SER-78A, MET-87A, TRP-88A, and A5N-91A are other molecules with which the delamanid formed H-bonds. Delamanid also creates the TYR-65A that stacks. Delamanid joins forces with GLY-53A, TYR-133A, LYS-104A, ARG-54A, and PRO-86A to form hydrophobic and other connections [24]. Liglot and 2D complexes of bedaquiline and delamanid shown in figure 1.




Figure 1. A) Ligplot and B) 2D Complex of Bedaquiline with 7JGCC) Ligplot and D) 2D Complex of Delamanid with 3R5R

2.3 Pretomanid

Pretomanid, also known as PA-824, is a nitroimidazole medication that is presently undergoing clinical studies to see whether it has any effect on both drug-susceptible and drug-resistant TB. Pretomanid,

shorten the duration of TB treatment, surpassed rifampicin and pyrazinamide in terms of its ability to kill non-replicating *Mycobacterium tuberculosis* [25, 26].

Mechanism of Action: The suppression of lipid and protein synthesis is its mode of action. It prevents the formation of the mycolates found in mycobacterial cell walls [27]. A recent investigation on *M. smegmatis*, a species that is genetically related to MTB, however, revealed that pretomanid may operate instead by interfering with bacterial sugar phosphate metabolism and causing the bacteria to produce the deadly substance methylglyoxal [28].

Binding pockets: Barry C. Finzel and Courtney C. Aldrich in the *Journal of Medicinal Chemistry* came to the conclusion that pretomanid exhibits hydrophobic interactions with PRO-24A and TYR-25A. Pretomanid had the ability to stack with TRP-64A at various distances and TYR-157A. Additionally, GLY-156A forms halogen bonds with it. Together with ILE-256A, ALA-226A, GLY-227A, PRO-24A, MET-91A, THR-318B and PH-92B, it also forms the other bonds [29].

2.4 Contezolid:


The first oxazolidinone antibiotic to show potential for use in the management of infections driven on by Gram-positive pathogens was contezolid or MRX-1 [30].

Mechanism of Action: In general, oxazolidinones work by preventing the production of new proteins. It was believed that they accomplish this by obstructing translational integrity [31, 32]. Figure 2 depicts the Liglot and 2D complexes of pretomanid and contezolid.

2.5 Sutezolid

An oxazolidinone derivative of linezolid called sutezolid (PNU-100480 or U-100480) is now being tested in clinical studies to treat tuberculosis. In a murine model, sutezolid was found to be more effective than linezolid, and its combination with I and II-line anti-TB medications demonstrated excellent bactericidal action with the ability to accelerate therapy.

Mechanism of Action: Inhibits protein synthesis [33-35].






Figure 2. A) Ligplot and B) 2D Complex of Pretomanid with 4XJOC) Ligplot and D) 2D Complex of Contezolid with 6WQN


Binding Pockets: According to the research paper by Mendes, V. Sutezolid formed hydrophobic and other interactions with ASN-179A, TRP-103, PHE-110-GLU-180, LEU-183, PHE-114, MET-142, TRP-145, TRY-148, THR-149, TRP-138, ILE107, GLY-106, and LEU-87[36].

2.6 Macozinone

Macozinone did not interact negatively or favourably with I-line or II-line drugs in a trial conducted by Lupine and colleagues. Macozinone had improved bactericidal action in the lungs and spleen that was comparable to isoniazid when tested in a chronic TB murine model.

Mechanism of Action: By binding covalently to a cysteine residue in the active site of decaprenylphosphoryl-b-D-ribose 2'-epimerase (DprE1), inhibits the formation of cell wall[37].

Binding pockets: According to the research paper by Vadim Makaro and Stewart T. Cole, Macozinone exhibits hydrophobic interactions with VAL-365A. Macozinone joins LYS-134A and LYS-418A to produce H-bonds. It also displays ASN-385A halogen bonding. Additionally, it displays interactions with SER-228A, GLY-133A, HIS-132A, PHE-369A, GLY-117A, and CYS-387A as well as additional interactions [38]. Figure 3 depicts the Liglot and 2D complexes of sutezolid and macozinone.






Figure 3. A) Ligplot and B) 2D Complex of Sutezolid with 5ZN1C) Ligplot and D) 2D Complex of Macozinone with 4NCR

2.7 Linezolid

Linezolid exhibits significant action against *M. tuberculosis* that is rapidly developing, slowly growing, and incapable of reproducing itself, demonstrating its potential to be used in TB therapy regimens that are shorter in duration[39].

Mechanism of Action: An oxazolidinone antibiotic called linezolid works to suppress translation by attaching to the ribosome's 50S subunit and competing with the cell's natural substrates[40-41].

Binding pockets: According to the research paper by Mendis V., Linezolid exhibits hydrophobic and other interactions with ASN-179, ASN-176, VAL-152, THR-149, TYR-148, TRP-145, MET-142, LEU-183, MET-102, TRP-103, TRP-207, ILE-107 and PHE-110[38].

2.8 Levofloxacin

The L-isomer of the D, L-racemate ofloxacin, levofloxacin is an oral fluoroquinolone antibiotic that was created in Japan. Levofloxacin has a tolerability profile that is comparable to other oral fluoroquinolones, with gastrointestinal and central nervous system side effects being the most frequent. The pharmacokinetics of levofloxacin is not influenced by food, age, or gender.

Mechanism of Action: Modifications to the cell membrane porin channels or the DNA gyrase A or B subunits are two examples of chromosomal abnormalities that might cause bacterial resistance to levofloxacin[42-46].

Binding pockets: Levofloxacin is used in the research work by L. Mark Fisherb and Mark R. Sandersona, and they discover that it exhibits hydrophobic interactions with ILE-170A. Along with ARG-117A, GLY-173A, ALA-175A, ASN-326A, SER-436D, LYS-458C, ASN-461C, and ASN-473, it also forms H-bonds with them[47]. Figure 4 displays the linezolid and levofloxacin ligplot and 2D complexes.




Figure 4. A) Ligplot and B) 2D Complex of Linezolid with 5NZC) Ligplot and D) 2D Complex of Levofloxacin with 3RAE

2.9 Moxifloxacin

A fluoroquinolone antibiotic that exhibits a broad spectrum of activity, moxifloxacin is used to treat endocarditis, sinusitis, pneumonia, and TB [48, 49].

Mechanism of Action: Inhibits the synthesis of bacterial DNA[50-52].

Binding pockets: In their article, Pan F. Chan, Velupillai Srikannathasan, and Jianzhong Huang came to the conclusion that moxifloxacin exhibits hydrophobic interactions as well as others with ARG-122A, GLY-459B, SER-84A, and GLY-473B [53].

2.10 Rifampicin

Rifampicin, also known as rifampin, was developed in 1968 and is still one of the main medications used to treat tuberculosis (TB) because of its sterilizing properties and capacity to accelerate recovery times when used at high doses[54-58]. In order to treat patients with pulmonary TB who were chronically resistant to treatment, rifampicin was initially used in clinical settings in the late 1960s[59-61].

Mechanism of Action: The enzyme responsible for DNA transcription in bacteria, RNA polymerase, is inhibited by rifampicin[62].

Binding pockets: Gerard D. Wright and his research team demonstrated in their study, "Rifamycin Resistance Enzyme with an Unprecedented Mechanism of Action," that rifampicin interacts hydrophobically with LEU-176A, PHE-257A, and PRO-284A. The H-bonds with GLY-9A, THR-285A, ARG-213A, MET-205, ALA-204, VAL-215, ARG-201, MET-342, GLY-286, GLY-287, PHE-74, VAL-93, GLY-44, GLN-43, VAL-69 and ARG-196 are also shown [63].

2.11 Isoniazide

This synthetic drug was first designated by Meller and Melley in the year 1912. The antibiotic isoniazid, also referred to as isonicotinic acid hydrazide, is used to treat tuberculosis. It is often combined with rifampicin, pyrazinamide, and either streptomycin or ethambutol to treat active tuberculosis. It is frequently used alone to treat latent tuberculosis.

Mechanism of Action: Inhibits mycolic acid biosynthesis[64].

Binding pockets: According to Gegia M. study, the isoniazide interacts hydrophobically with PHE-78, PRO-79, ALA-80, LEU-88 and TYR-85A. Instead, it displays an H-bond with the amino acid THR-307A. It Displays other interactions with THR307, HIS-269, and HIS-84.[65] Figure 5 displays the moxifloxacin, rifampicin and isoniazideliglot and 2D complexes.

2.12 Pyrazinamide

Because of its unique method of action (interference with ATP synthesis), pyrazinamide will probably continue to play a significant role for treatment of drug-susceptible and multidrug-resistant TB (MDR-TB).




Figure 5. A) Ligplot and B) 2D Complex of Moxifloxacin with 5CDQC) Ligplot of Isoniazidecomplexed with 4PAE D & E) 2D Complex of Isoniazide with 4PAE&Rifampicin with 6BRD

Mechanism of Action:Inhibit the growth of bacteria[66-67].

Binding pockets:According to Petrella, S. study published in thePLoS One journal, pyrazinamide exhibits interactions with the following amino acids: ARG-255A, HIS-109, GLN-105A, ASP-108A, GLU-258 and PRA-598[68].

2.13 Ethambutol

The main purpose of the drug ethambutol is to cure tuberculosis. It is typically administered along with other anti-tuberculosis drugs such isoniazid, rifampicin, and pyrazinamide. Mycobacterium avium complex and Mycobacterium kansasii may also be treated with it.

Mechanism of Action:The bacteriostatic medication ethambutol prevents bacilli from reproducing by interfering with the formation of arabinogalactan in the cell wall [69].

Binding pockets:In their article, Zhang, L. demonstrated that ethambutol has hydrophobic and other interactions with VAL-1004A, ASP-279A, TRP-965, TRP-572, TYR-282, ILE-283, TYR-314 and ARG-383[70].

2.14 Para Amino Salicylic Acid


In the 1940s, it was discovered that para-amino salicylic acid (PAS), a second-line anti-TB medication (SLD), was useful for treating tuberculosis.

Mechanism of Action: Inhibition of follic acid synthesis[71].

Binding pockets:In their article, Schreuder, H.A. demonstrated that the para amino salicylic acid exhibits interactions with TRP-185, TYR-222, PRO293, ALA-296, LEU-210, TYR-201, LEU-199, VAL-47, ARG-214, SER-212, THR-296 and ARG-44 that are hydrophobic and others[72].Figure 6 displays the pyrazinamide, ethambutol and para-amino salicylic acid liglot and 2D complexes.

3. RESULTS AND DISCUSSION

The 3D crystal structures of all the enzymes were retrieved from the protein data bank-RCSB (<http://www.rcsb.org>). For interpretation and visualization of interactions of protein-ligand complex, Mastero and Biovia discovery studio visualizer and freely accessible server PDBsum (<http://www.ebi.ac.uk/pdbsum>) for Ligplot were used[73].






Figure 6. A) Ligplot and B) 2D Complex of Pyrazinamide with 3PL1 C) Ligplot and D) 2D Complex of Ethambutol with 7BVE E)Ligplot and D) 2D Complex of PAS with 1PBC

Bedaquiline forms non-bonded contacts with amino acids of 7JGC protein. The oxygen and carbons of amino acid proline-172A forms non bonded contacts with carbon and oxygen atoms of bedaquiline with varying distances (3.05-3.65Å). Isoleucine-173A forms two non bonded contacts with O

and C atoms of bedaquiline with distance 3.05-3.24Å. Valine-176A formed two interactions with carbons having distance 3.51 and 3.35Å. Five different number carbons of Glutamic acid-65 interacts with C and N atoms of bedaquiline and oxygen atoms of seven different positions of Glutamic acid-65 forms interactions with C and O atoms of the molecule BQ with varying distances. The Tyrosine-68 residue forms non bonded contacts with different C and N atoms of bedaquiline. Also C and O atoms of Alanine-66 and 67, Leucine-63 and 72 and Phenylalanine-69 residue creates interactions with C atoms of BQ (distances ranges from 3.10 to 3.90 Å).

The drug delamanid formed 18 (hydrogen bonds) bonded and 140 non-bonded interactions with 3R5R.O and N atoms of amino acids from A chain Arginine-54 and 60, Lysine-55 and 79, Thyronine-56, Proline-63, Tyrosine-65 and 133, Alanine-76, Methionine-87, Treptophan-88, Asparagine-91 and Valine-46 formed hydrogen bonds (having distance ranges from 2.56 to 3.26Å) with different number of O and N atoms of delamanid molecule. Pretomanid forms one hydrogen bond and 62 non-bonded interactions with protein 4XJO. Nitrogen atom of Glycine-93B formed hydrogen bond with oxygen-34 atom of pretomanid molecule with distance 2.92Å. In non-bonded interactions of Proline-24A and 317B, Tyrosine-25A, tryptopan-64A, Tyrosine-157A, Glycine-172A, 316B and 93, Methionine-91, Threonine-318B amino acids were involved. Sutezolid forms one hydrogen bond and 67 non-bonded interactions with protein 5NZ1. Nitrogen atom of Asparagine-179A formed hydrogen bond with oxygen-10 of sutezolid molecule with distance 2.19Å. Macozinone molecule forms 3-(hydrogen bonds) bonded and 33 non-bonded contacts with protein 4NCR. Nitrogen atom of Lysine-134A and 418A interacts with oxygen of molecule with distance 3.16Å and 3.03 respectively. S atom of Cystine-187A formed hydrogen bond with N atom with distance 1.66 Å.

Linezolid forms single hydrogen bond and 53 non-bonded interactions with protein 5NZ0. Nitrogen atom of Asparagine-179A formed hydrogen bond with C-18 atom of linezolid molecule with distance 3.13Å. MET-102A, TRP-103A, GLY -106A, ILE-107A, PHE-110A, TRP-145A, TYR-148A, THR-149A, ASN-176A and 179A and TRP-207A residues contributes in non-bonded contacts. Levofloxacin forms a hydrogen bond and 87 non-bonded interactions with protein 3RAE. OxygenG atom of Serine-79A formed hydrogen bond with O-18 atom of levofloxacin molecule with distance 2.81Å. Moxifloxacin molecule forms 3-(H-bond) bonded and 87 non-bonded contacts with protein 5CDQ. Nitrogen atom of Lysine-134A and 418A interacts with oxygen of molecule with distance 3.16 and 3.03Å respectively. Rifampicin formed 2 hydrogen bonds and 52 non-bonded interactions with protein 6BRD. Nitrogen from Arginine-196 and 216A binds with oxygen atom no.8 and 2 of rifampicin molecule with distance 2.67 and 3.18Å respectively. Isoniazide formed one hydrogen bond and 53 non-bonded contacts with protein 4PAE. Amino acid Thyronine-307A binds with N-3 of Isoniazide molecule with distance 2.61Å. Pyrazinamide formed 1 hydrogen bond and 5 non-bonded contacts with protein 3PL1. Amino acid Glanine-105A formed hydrogen bond with N of pyrazinamide molecule with distance 2.59Å. Ethambutol formed 1 hydrogen bond and 20 non-bonded interactions with the protein 7BVE. Aspartic acid-279A formed hydrogen bond with nitrogen of ethambutol with distance 5.14Å. Para-amino salicylic acid formed 10 hydrogen and 42 non-bonded contacts with the protein 1PBC. The residues involved in the hydrogen bonds are Tyrosine-201 and 222A, Serine-212A, Arginine-214 and Proline-293A. The information about crystal structures of the entire protein-ligand complexes were summarized in table 1.

Table 1. FDA Approved Antitubercular drugs MOA and their PDB details:

| S.N. | Drug | PDB Code | Classification | Organism | Mutat ion | Resolu tion | Mechanism of Action |
|------|-------------|----------|-----------------------|-----------------|-----------|-------------|--|
| 1 | Bedaquiline | 7JGC | Hydrolase | M. Smegmatis | No | 3.40A° | Inhibits the Proton pump ATP Synthase |
| 2 | Delamanid | 3R5R | Oxidoreductase | M. tuberculosis | No | 2-10A° | Inhibits mycolic acid biosynthesis. |
| 3 | Pretomanid | 4XJO | Transferase inhibitor | M. tuberculosis | No | 1.50 A° | Inhibition of mycolic acid synthesis |
| 4 | Contezolid | 6WQN | Ribosome | S. aureus | Yes | 2.90A° | Inhibits protein synthesis. |
| 5 | Sutezolid | 5NZ1 | Transcription | M. Tuberculosis | No | 2.33 Å | Inhibits protein synthesis. |
| 6 | Macozinone | 4NCR | Oxidoreductase | M. Tuberculosis | No | 1.88 A° | By forming a covalent bond with a cystine residue, it prevents the synthesis of cell wall. |
| 7 | Linezolid | 5NZ0 | Transcription | M. tuberculosis | No | 1.82 Å | Inhibit translocation by binding to the 50s subunits |

| | | | | | | | |
|----|--------------|------|------------------------------|-------------------------|-----|---------|--|
| 8 | Levofloxacin | 3RAE | Isomerase/ DNA/Antibiotic | S. pneumoniae | No | 2.90Å° | of ribosomes. Inhibit DNA gyrase subunit (Gyrase A & Gyrase B) topoisomerase iv subunit (Par C & Part E). |
| 9 | Moxifloxacin | 5CDQ | Hydrolase | S.aureussubaureus | No | 2.95Å° | Inhibit DNA gyrase subunit (Gyrase A & Gyrase B) topoisomerase iv subunit (Par C & Part E). |
| 10 | Rifampicin | 6BRD | Oxidoreductase | S.venezuelige | No | 3.32 Å° | Inhibit RNA polymerase. |
| 11 | Isoniazide | 4PAE | Oxidoreductase | Synechococcus elongatus | Yes | 3.21 Å | Inhibit synthesis of mycolic acid. |
| 12 | Pyrazinamide | 3PL1 | Oxidoreductase | A. flavus | Yes | 1.68 Å° | Inhibit the growth of bacteria. |
| 13 | Ethambutol | 7BVE | Transferase | M. Smegmatis | No | 2.81 Å | Inhibition of arabinosyl transferase. |
| 14 | PAS | 1PBC | Oxidoreductase | Pseudomonas fluorescens | No | 2.80 Å | Inhibition of follic acid synthesis. |

4. CONCLUSION

Collectively, our research offers a structural basis for understanding the significance of binding sites of amino acids of a particular protein. It serves as a useful tool for understanding the relationship between structure and function. On the basis of significant binding pockets, including donors and acceptors of amino acid residues, the researchers can investigate the newer molecular architectures for treating M. tuberculosis. The main finding is that future research using combined ligands and receptor-based design will generate a new class of anti-tubercular drugs that will benefit millions of patients.

REFERENCES

- <https://www.who.int/teams/global-tuberculosis-programme/tb-reports/global-tuberculosis-report-2022>.
- Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis*. 2020; 20(5): 533–534. doi: [https://doi.org/10.1016/S1473-3099\(20\)30120-1](https://doi.org/10.1016/S1473-3099(20)30120-1).
- Hale, T., Angrist, N., Goldszmidt, R. et al. A global panel database of pandemic policies (Oxford COVID-19 Government Response Tracker). *Nat Hum Behav*. 2021; 5, 529–538. <https://doi.org/10.1038/s41562-021-01079-8>
- Padda IS, Muralidhara Reddy K. Antitubercular Medications. 2022 Nov 30. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan. PMID: 32491598.
- World Health Organization . (2018). Global tuberculosis report 2018. World Health Organization. <https://apps.who.int/iris/handle/10665/274453>.
- Poce G, Cocozza M, Consalvi S, Biava M. SAR analysis of new anti-TB drugs currently in pre-clinical and clinical development. *Eur J Med Chem*. 2014; 86:335-351. doi: 10.1016/j.ejmech.2014.08.066.
- Cohen J. Infectious disease. Approval of novel TB drug celebrated with restraint. *Science*. 2013 Jan 11;339 (6116):130. doi: 10.1126/science.339.6116.130. PMID: 23307714.
- Iacobino A, Piccaro G, Giannoni F, Mustazzolu A, Fattorini L. Fighting tuberculosis by drugs targeting nonreplicating Mycobacterium tuberculosis bacilli. *Int J Mycobacteriol*. 2017;6(3):213-221. doi: 10.4103/ijmy.ijmy_85_17.
- Alzayer Z, Al Nasser Y. Primary Lung Tuberculosis. 2023 Jan 2. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan. PMID: 33620814.
- Lee JY. Diagnosis and treatment of extrapulmonary tuberculosis. *Tuberc Respir Dis (Seoul)*. 2015; 78(2):47-55. doi: 10.4046/trd.2015.78.2.47.

11. Purohit M, Mustafa T. Laboratory Diagnosis of Extra-pulmonary Tuberculosis (EPTB) in Resource-constrained Setting: State of the Art, Challenges and the Need. *J Clin Diagn Res.* 2015;9(4):EE01-6. doi: 10.7860/JCDR/2015/12422.5792.
12. Lange C, Mori T. Advances in the diagnosis of tuberculosis. *Respirology.* 2010;15(2):220-40. doi: 10.1111/j.1440-1843.2009.01692.x.
13. World Health Organization . (2017). Global tuberculosis report 2017. World Health Organization. <https://apps.who.int/iris/handle/10665/259366>.
14. Zumla A, Nahid P, Cole ST. Advances in the development of new tuberculosis drugs and treatment regimens. *Nat Rev Drug Discov.* 2013;12(5):388-404. <https://doi.org/10.1038/nrd4001>.
15. Marriner GA, Nayyar A, Uh E. et al. The medicinal chemistry of tuberculosis chemotherapy. In: Elliott R. ed. Third World Diseases. *Top Med Chem.* 2011; 7: 47–124. https://doi.org/10.1007/7355_2011_13.
16. Matteelli A, Carvalho AC, Dooley KE, Kritski A. TMC207: The first compound of a new class of potent anti-tuberculosis drugs. *Future Microbiol.* 2010;5(6):849e858. <https://doi.org/10.2217/fmb.10.50>.
17. Diacon AH, Pym A, Grobusch M, et al. The diarylquinoline TMC207 for multidrug-resistant tuberculosis. *N Engl J Med.* 2009;360:2397e2405. <https://doi.org/10.1056/NEJMoa0808427>.
18. Cohen J. Infectious disease. Approval of novel TB drug celebrated with restraint. *Science.* 2013;339(6116):130. doi: 10.1126/science.339.6116.130.
19. Guglielmetti L, Jaspard M, Le Dû D, et al. Long-term outcome and safety of prolonged bedaquiline treatment for multidrug-resistant tuberculosis. *Eur Respir J.* 2017; 49: 1601-799. <https://doi.org/10.1183/13993003.01799-2016>.
20. Andries K, Verhasselt P, Guillemont J, et al. A diarylquinoline drug active on the ATP synthase of Mycobacterium tuberculosis. *Science.* 2005;307(5707):223e227. <https://doi.org/10.1126/science.1106753>.
21. Ryan NJ, Lo JH. Delamanid: first global approval. *Drugs.* 2014;74(9):1041e1045. <https://doi.org/10.1007/s40265-014-0241-5>.
22. Kailu Yang, Jeng-Yih Chang, Zhicheng Cui, Xiaojun Li, Ran Meng, Lijun Duan, Jirapat Thongchol, Joanita Jakana, Christoph M. Huwe, James C. Sacchettini and Junjie Zhang. Structural insights into species-specific features of the ribosome from the human pathogen Mycobacterium tuberculosis. *Nucleic Acids Res.* 2017; 45 18. doi:10.1093/nar/gkx785.
23. Tiberi S, Du Plessis N, Walzl G, et al. Tuberculosis: progress and advances in development of new drugs, treatment regimens, and host-directed therapies. *Lancet Infect Dis.* 2018;18(7):e183ee198. [https://doi.org/10.1016/S1473-3099\(18\)30110-5](https://doi.org/10.1016/S1473-3099(18)30110-5).
24. Cellitti, S.E., Shaffer, J., Jones, D.H., Mukherjee, T., Gurumurthy, M., Bursulaya, B., Boshoff, H.I., Choi, I., Nayyar, A., Lee, Y.S., Cherian, J., Niyomrattanakit, P., Dick, T., Manjunatha, U.H., Barry, C.E., Spraggon, G., Geierstanger, B.H. Structure of Ddn, the deazaflavin-dependent nitroreductase from Mycobacterium tuberculosis involved in bioreductive activation of PA-824. *Structure.* 2012;20:101-112. <https://doi.org/10.1016/j.str.2011.11.001>
25. Tiberi S, du Plessis N, Walzl G, et al. Tuberculosis: progress and advances in development of new drugs, treatment regimens, and host-directed therapies. *Lancet Infect Dis.* 2018;18(7):e183ee198. [https://doi.org/10.1016/S1473-3099\(18\)30110-5](https://doi.org/10.1016/S1473-3099(18)30110-5).
26. Somasundaram S, Anand RS, Venkatesan P, Paramasivan CN. Bactericidal activity of PA-824 against Mycobacterium tuberculosis under anaerobic conditions and computational analysis of its novel analogues against mutant Ddn receptor. *BMC Microbiol.* 2013;13:218. <https://doi.org/10.1186/1471-2180-13-218>.
27. Stover CK, Warrener P, Van Devanter DR, et al. A small molecule nitroimidazopyran drug candidate for the treatment of tuberculosis. *Nature.* 2000;405(6789):962e966. <https://doi.org/10.1038/35016103>.
28. Baptista R, Fazakerley DM, Beckmann M, Baillie L, Mur LAJ. Untargeted metabolomics reveals a new mode of action of pretomanid (PA-824). *Sci Rep.* 2018;8(1):5084. <https://doi.org/10.1038/s41598-018-23110-1>.
29. Feng Liu, Surendra Dawadi, Kimberly M. Maize, Ran Dai, Sae Woong Park, Dirk Schnappinger, Barry C. Finzel and Courtney C. Aldrich. Structure-Based Optimization of Pyridoxal 5'-Phosphate-Dependent Transaminase Enzyme (BioA) Inhibitors that Target Biotin Biosynthesis in Mycobacterium tuberculosis; *J. Med. Chem.* 2017;60:5507–5520. doi: 10.1021/acs.jmedchem.7b00189.
30. Li C-R, Zhai Q-Q, Wang X-K, et al. In Vivo antibacterial activity of MRX-I, a new oxazolidinone. *Antimicrob Agents Chemother.* 2014;58(4):2418e2421. <https://doi.org/10.1128/AAC.01526-13>.

31. Jadhavar PS, Vaja MD, Dhameliya TM, Chakraborti AK. Oxazolidinones as anti-tubercular agents: discovery, development and future perspectives. *Curr Med Chem*. 2015;22(38):4379e4397. doi: 10.2174/0929867323666151106125759.
32. Thompson J, O'Connor M, Mills JA, Dahlberg AE. The protein synthesis inhibitors, oxazolidinones and chloramphenicol, cause extensive translational inaccuracy in vivo. *J Mol Biol*. 2002;322(2):273e279. doi: 10.1016/s0022-2836(02)00784-2.
33. Wallis RS, Dawson R, Friedrich SO, et al. Mycobactericidal activity of sutezolid (PNU-100480) in sputum (EBA) and blood (WBA) of patients with pulmonary tuberculosis. *PLoS One*. 2014;9(4):e94462. <https://doi.org/10.1371/journal.pone.0094462>.
34. Williams KN, Stover CK, Zhu T, et al. Promising antituberculosis activity of the oxazolidinone PNU-100480 relative to that of linezolid in a murine model. *Antimicrob Agents Chemother*. 2009;53(4):1314e1319. <https://doi.org/10.1128/AAC.01182-08>.
35. Williams KN, Brickner SJ, Stover CK, et al. Addition of PNU100480 to first-line drugs shortens the time needed to cure murine tuberculosis. *Am J Resp Crit Care*. 2009;180(4):371e376. <https://doi.org/10.1164/rccm.200904-0611OC>.
36. Nikiforov, P.O., Blaszczyk, M., Mendes, V., Boshoff, H.I., Barry, C.E., Blundell, T.L., Abell, C. The antibiotics linezolid and sutezolid are ligands for Mycobacterium tuberculosis EthR. <https://www.rcsb.org/structure/5nz0>.
37. Lupien A, Vocat A, Foo CS-Y, et al. Optimized background regimen for treatment of active tuberculosis with the next generation benzothiazinone macozinone (PBTZ169). *Antimicrob Agents Chemother*. 2018;62(11). <https://doi.org/10.1128/AAC.00840-18>. e00840-18.
38. Makarov, V., Lechartier, B., Zhang, M., Neres, J., van der Sar, A.M., Raadsen, S.A., Hartkoorn, R.C., Ryabova, O.B., Vocat, A., Decosterd, L.A., Widmer, N., Buclin, T., Bitter, W., Andries, K., Pojer, F., Dyson, P.J., Cole, S.T. Towards a new combination therapy for tuberculosis with next generation benzothiazinones. *EMBO Mol Med*. 2014; 6: 372-383. <https://doi.org/10.1002/emmm.201303575>
39. Drusano GL, Myrick J, Maynard M, et al. Linezolid kills acid phase and nonreplicative-persisters of Mycobacterium tuberculosis in a hollow-fiber infection model. *Antimicrob Agents Chemother*. 2018;62(8). <https://doi.org/10.1128/AAC.00221-18>.
40. Brown AN, Drusano GL, Adams JR, et al. Preclinical evaluations to identify optimal linezolid regimens for tuberculosis therapy. *mBio*. 2015;6(6). <https://doi.org/10.1128/mBio.01741-15>. e01741-15.
41. Ippolito JA, Kanyo ZF, Wang D, et al. Crystal structure of the oxazolidinone antibiotic linezolid bound to the 50S ribosomal subunit. *J Med Chem*. 2008;51(12):3353e3356. <https://doi.org/10.1021/jm800379d>.
42. Wimer SM, Schoonover L, Garrison MW. Levofloxacin: a therapeutic review. *Clin Ther*. 1998; 20(6):1049-70. doi: 10.1016/s0149-2918(98)80104-5.
43. Davis R, Bryson HM. Levofloxacin. A review of its antibacterial activity, pharmacokinetics and therapeutic efficacy. *Drugs*. 1994;47(4):677e700. <https://doi.org/10.2165/00003495-199447040-00008>.
44. Langtry HD, Lamb HM. Levofloxacin. Its use in infections of the respiratory tract, skin, soft tissues and urinary tract. *Drugs*. 1998;56(3):487e515. <https://doi.org/10.2165/00003495-199856030-00013>.
45. Fish DN, Chow AT. The clinical pharmacokinetics of levofloxacin. *Clin Pharmacokinet*. 1997;32(2):101e119. <https://doi.org/10.2165/00003088-199732020-00002>.
46. Chien SC, Wong FA, Fowler CL, et al. Double-blind evaluation of the safety and pharmacokinetics of multiple oral once-daily 750-milligram and 1-gram doses of levofloxacin in healthy volunteers. *Antimicrob Agents Chemother*. 1998; 42(4):885-8. doi: 10.1128/AAC.42.4.885.
47. Dennis A, Veselkov, Ivan Laponogov, Xiao-Su Pan, Jogitha Selvarajah, Galyna B. Skamrova, Arthur Branstrom, Jana Narasimhan, Josyula V. N. Vara Prasad, L. Mark Fisher and Mark R. Sanderson. Structure of quinolone-stabilized cleavage complex of topoisomerase IV from Klebsiella pneumoniae and comparison with a related Streptococcus pneumoniae complex. *Acta Cryst*. 2016; D72:488–496. <http://dx.doi.org/10.1107/S2059798316001212>.
48. Samilski JA, Lau TT, Elbe DH, Aulakh AK, Lun EM. Drug use evaluation of moxifloxacin (avelox) using a hand-held electronic device at a Canadian teaching hospital. *P T*. 2012; 37(5):291-9.
49. Gillespie SH. The role of moxifloxacin in tuberculosis therapy. *Eur Respir Rev*. 2016;25(139):19e28. <https://doi.org/10.1183/16000617.0085-2015>.
50. Wimer SM, Schoonover L, Garrison MW. Levofloxacin: a therapeutic review. *Clin Therapeut*. 1998;20(6):1049e1070. [https://doi.org/10.1016/S0149-2918\(98\)80104-5](https://doi.org/10.1016/S0149-2918(98)80104-5).
51. Langtry HD, Lamb HM. Levofloxacin. Its use in infections of the respiratory tract, skin, soft tissues and urinary tract. *Drugs*. 1998;56(3):487e515. <https://doi.org/10.2165/00003495-199856030-00013>.

52. Yamaba Y, Ito Y, Suzuki K, et al. Moxifloxacin resistance and genotyping of Mycobacterium avium and Mycobacterium intracellulare isolates in Japan. *J Infect Chemother.*2019;25:995e1000. <https://doi.org/10.1016/j.jiac.2019.05.028>.
53. Pan F, Chan, Velupillai Srikanthasan, Jianzhong Huang, Haifeng Cui, Andrew P. Fosberry, Minghua Gu, Michael M. Hann, Martin Hibbs, Paul Homes, Karen Ingraham, Jason Pizzollo, Carol Shen, Anthony J. Shillings, Claus E. Spitzfaden, Robert Tanner, Andrew J. Theobald, Robert A. Stavenger, Benjamin D. Bax & Michael N. Gwynn. Structural basis of DNA gyrase inhibition by antibacterial QPT-1, anticancer drug etoposide and moxifloxacin. *Nat. Comm.* 2015; 6:10048. doi:10.1038/ncomms10048.
54. Sensi P. History of the development of rifampin. *Rev Infect Dis.* 1983;5(suppl 3):S402eS406. https://doi.org/10.1093/clinids/5.supplement_3.s402.
55. Prideaux B, Via LE, Zimmerman MD, et al. The association between sterilizing activity and drug distribution into tuberculosis lesions. *Nat Med.* 2015;21(10):1223e1227. <https://doi.org/10.1038/nm.3937>.
56. Dickinson JM, Mitchison DA. Experimental models to explain the high sterilizing activity of rifampin in the chemotherapy of tuberculosis. *Am Rev Respir Dis.* 1981;123 (4Pt1):367e371. <https://doi.org/10.1164/arrd.1981.123.4.367>.
57. Svensson EM, Svensson RJ, te Brake LHM, et al. The potential for treatment shortening with higher rifampicin doses: relating drug exposure to treatment response in patients with pulmonary tuberculosis. *Clin Infect Dis.* 2018;67(1):34e41. <https://doi.org/10.1093/cid/ciy026>.
58. Dutta NK, Karakousis PC. Can the duration of tuberculosis treatment be shortened with higher dosages of rifampicin? *Front Microbiol.* 2015;6. <https://doi.org/10.3389/fmicb.2015.01117>.
59. Grobbelaar M, Louw GE, Sampson SL, van Helden PD, Donald PR, Warren RM. Evolution of rifampicin treatment for tuberculosis. *Infect Genet Evol.* 2019;74:103937. <https://doi.org/10.1016/j.meegid.2019.103937>.
60. Murray JF, Schraufnagel DE, Hopewell PC. Treatment of tuberculosis. A historical perspective. *Ann Am Thorac Soc.* 2015;12(12):1749e1759. <https://doi.org/10.1513/AnnalsATS.201509-632PS>.
61. Jawahar MS. Current trends in chemotherapy of tuberculosis. *Indian J Med Res.* 2004; 120(4):398-417.
62. Wehrli W. Rifampin: mechanisms of action and resistance. *Rev Infect Dis.* 1983;5(suppl3):S407eS411. https://doi.org/10.1093/clinids/5.supplement_3.s407
63. Kalinka Koteva, Georgina Cox, Jayne K. Kelso, Matthew D. Surette, Haley L. Zubyk, Linda Ejim, Peter Stogios, Alexei Savchenko, Dan Sørensen, and Gerard D. Wright. Rox, a Rifamycin Resistance Enzyme with an Unprecedented Mechanism of Action. *Cell Chemical Biology.* 2018; 25:1–10. <https://doi.org/10.1016/j.chembiol.2018.01.009>.
64. Unissa AN, Subbian S, Hanna LE, Selvakumar N. Overview on mechanisms of isoniazid action and resistance in Mycobacterium tuberculosis. *Infect Genet Evol.* 2016;45: 474-492. doi: 10.1016/j.meegid.2016.09.004
65. Gegia M, Winters N, Benedetti A, van Soolingen D, Menzies D. "Treatment of isoniazid-resistant tuberculosis with first-line drugs: a systematic review and meta-analysis". *The Lancet. Infectious Diseases.* 2017;17 (2):223–234. [doi:10.1016/S1473-3099\(16\)30407-8](https://doi.org/10.1016/S1473-3099(16)30407-8)
66. Shi W, Chen J, Feng J, Cui P, Zhang S, Weng X, Zhang W, Zhang Y. Aspartate decarboxylase (PanD) as a new target of pyrazinamide in Mycobacterium tuberculosis. *Emerg Microbes Infect.* 2014;3(8):e58. doi: 10.1038/emi.2014.61
67. Shi W, Zhang X, Jiang X, Yuan H, Lee JS, Barry CE III, et al. Pyrazinamide Inhibits Trans-Translation in Mycobacterium tuberculosis. *Science.* 2011;333(6049):1630–2. doi: 10.1126/science.1208813.
68. Petrella, S., Gelus-Ziental, N., Maudry, A., Laurans, C., Boudjelloul, R., Sougakoff, W. Crystal Structure of the Pyrazinamidase of Mycobacterium tuberculosis: Insights into Natural and Acquired Resistance to Pyrazinamide. *PLoS One.* 2011; 6: e15785-e15785. <https://doi.org/10.1371/journal.pone.0015785>
69. Palomino JC, Martin A. Drug Resistance Mechanisms in Mycobacterium tuberculosis. *Antibiotics (Basel).* 2014;3(3):317-40. doi: 10.3390/antibiotics3030317.
70. Zhang, L., Zhao, Y., Gao, Y., Wu, L., Gao, R., Zhang, Q., Wang, Y., Wu, C., Wu, F., Gurcha, S.S., Veerapen, N., Batt, S.M., Zhao, W., Qin, L., Yang, X., Wang, M., Zhu, Y., Zhang, B., Bi, L., Zhang, X., Yang, H., Guddat, L.W., Xu, W., Wang, Q., Li, J., Besra, G.S., Rao, Z. Structures of cell wall

arabinosyltransferases with the anti-tuberculosis drug ethambutol. *Science*. 2020; 368(6496), 1211-1219. doi.org/10.1126/science.aba9102

71. Minato Y, Thiede JM, Kordus SL, McKlveen EJ, Turman BJ, Baughn AD. Mycobacterium tuberculosis folate metabolism and the mechanistic basis for para-aminosalicylic acid susceptibility and resistance. *Antimicrob Agents Chemother*. 2015;59(9):5097–5106. doi: 10.1128/AAC.00647-15
72. Schreuder, H.A., Mattevi, A., Obmolova, G., Kalk, K.H., Hol, W.G., van der Bolt, F.J., van Berkel, W.J. Crystal structures of wild-type p-hydroxybenzoate hydroxylase complexed with 4-aminobenzoate, 2,4-dihydroxybenzoate, and 2-hydroxy-4-aminobenzoate and of the Tyr222Ala mutant complexed with 2-hydroxy-4-aminobenzoate. Evidence for a proton channel and a new binding mode of the flavin ring. *Biochemistry*. 1994;33: 10161-10170. doi: 10.1021/bi00199a044.
73. Laskowski RA, Hutchinson EG, Michie AD, Wallace AC, Jones ML, Thornton JM PDBsum: a Web- based database of summaries and analyses of all PDB structures. *Trends Biochem Sci*. 1997; 22(12):488-90. doi: 10.1016/s0968-0004(97)01140-7.

UNDER PEER REVIEW