

Original Research Article

EVALUATION OF THE MEDICINAL POTENTIALS OF SOURSOP (*ANNONA MURICATA* SAFFORD) LEAF EXTRACTS AGAINST *EIMERIA TENELLA* INFECTION IN BROILER CHICKENS

ABSTRACT

Aims: Poultry industry in Nigeria has recorded considerable expansion in recent times but not without poultry diseases as one of the major threats to boosting poultry production. Protozoa cause the majority of parasitic infections and cause high mortalities. Parasitic diseases such as coccidiosis are of particular importance because of their high incidence in poultry occasioned by the tropical environmental conditions under which the farmer operates. The resistance to common anticoccidial drugs has led to exploring other natural sources for prophylaxis and therapy. This research evaluated the therapeutic efficacies of *Annonamuricata* at different dosages of 100, 125, 165mg/kg body weight using 120 broilers at 4 weeks of age.

Place and Duration of Study: Animal Parasitology and Microbiology Research Unit, Department of Animal Production and Health, Federal University of Technology, Akure, Nigeria, between September and December, 2022.

Methodology: 120 broilers of 4 weeks old were used for this study using completely randomized study design. The birds were infected with sporulated *Eimeria tenella* oocysts at week 4. Treatment with ethanolic extract of *Annonamuricata* was administered orally at week 5 using different dosages of 100, 125, 165mg/kg body weight for five consecutive days. Treatment commenced after the isolation of *Eimeria* oocysts in the faeces and onset on coccidial clinical signs as parameters to confirm manifestation of the infection. Blood parameters, faecal oocysts count and biochemical assays were conducted.

Results: Results revealed that therapeutic administration of *Annonamuricata* leaf extracts at minimum dose 100mg/kg body weight had significant reduction ($P < 0.05$) on excreted *Eimeria* oocysts counts without any significant deviation from normal haematological and biochemical indices of the broilers.

Conclusion: This study suggests the use of *Annonamuricata* in the treatment of coccidiosis in broiler chickens. It is therefore recommended that its extracts be further explored for actual phytochemical constituents responsible for the displayed anticoccidial properties.

Keywords: *Annonamuricata*; anticoccidial activities; *Eimeria tenella*; faecal oocysts; blood parameter; biochemical assay.

1. INTRODUCTION

Nowadays, poultry production has high demand all over the world. Poultry industry in Nigeria has recorded considerable expansion in recent times but not without poultry diseases as one of the major threats to boosting poultry production. This increasing demand has led to the usage of numerous antibiotic-free products. There is an increased pressure to reduce the number of antibiotics that are used as bacteriostatic or bactericidal agents for poultry, so there is a crucial requisite for unconventional resolutions to sustain the productivity and efficiency of poultry [1]. Now, there is also the use of herbal plants as an alternate for the prevention of intestinal parasitosis [2], [3].

Protozoa and helminths cause the majority of parasitic infections and cause high mortalities [4]. Parasitic diseases such as Coccidiosis are of particular importance because of their high incidence in poultry occasioned by the tropical environmental conditions under which the farmer operates [5]. Of all nine *Eimeria* species identified in poultry [6], *Eimeriatenella* and *Eimeriamaxima* are the most common [7], causing tissue damage, lowered feed intake, poor absorption of nutrients from the feed, dehydration and blood loss [5].

Vaccines as well as chemotherapeutic agents have been employed to suppress the menace of coccidiosis in the past, but are expensive to produce, can lose their potencies over a long storage period, resident organisms can undergo genetic shift/drift making the use of vaccine unreliable. The reduction in the use of chemically manufactured drugs can be attributed to poverty, inaccessibility, and decaying infrastructure. Also, resistance to some drugs over a period of time is a risk factor [8]. The use of alternative medicine, as a result, has been advocated [4]. Several diseases can be cured using traditional medicines that utilize plant, herb, or mineral ingredients [9].

Hence, the need for this research to explore the abundance of plants to provide surplus remedies against the havoc posed by coccidiosis to broiler farming in Nigeria. *Annonamuricata* is identified candidates to be explored for possible remedy to coccidiosis menace in the poultry industry.

Soursop (*Annonamuricata*) is endemic to the warmest areas of the tropics of South and Central America and the Caribbean. It is now widely found in tropical and subtropical regions of Central and South America, Western Africa, and Southeast Asia [10]. The aqueous extract of *A. muricata* is used to control lepidopteran larvae, aphids and thrips, among others [11], [12],[13]. Also, the leaves, bark, and roots of *A. muricata* have been used for its anti-inflammatory, hypotensive, sedative, hypoglycemic, smooth muscle relaxant, antiplasmodic and antitrypanosomal effects [14], [15], [16] among other uses.

2. METHODOLOGY

2.1 Test Organisms

Typed *Eimeriatenella* was obtained from Nigeria Veterinary Research Institute, (NVRI) Vom, Jos, Ngeria.

2.2 Identification and Authentication of Plants

The leaves of *Annonamuricata* were collected from their natural habitats around The Federal University of Technology, Akure, Ondo State and proper plant identifications was done by the Department of Pharmacognosy, Faculty of pharmacy, ObafemiAwolowo University, Ile Iife with voucher number FPI 2402.

2.3 Crude Extraction of *Annonamuricata*

The plants leaves were dried at room temperature, then pulverized with electric hammer mill. 1000g of the fine powder was soaked in 5liters of ethanol for 76hours. The extract was filtered using muslin cloth and filtrate was concentrated to dryness using Rotary-evaporator. The resulting organic extracts was stored in sterile bottles at refrigeration temperature (4 - 8°C) before carrying out the in vivo activities.

2.4 Phytochemical Analysis of Plants' Crude Extract

The powdered crude extracts of the plants leaves were subjected to phytochemical analysis for the presence of plant secondary metabolites using standard qualitative procedures [17], [18], [19].

Determination of LD₅₀: Lorkes' method as described by Eneget *et al.*, [20] was used to determine the Lethal Dose₅₀ (LD₅₀) from which the graded doses were calculated. Determination of graded doses: No mortality was recorded in any of the birds of all the dose variations, so the random starting dose was picked as 100mg/kg with the other doses as 125mg/kg and 165mg/kg.

2.5 Experimental design

A hundred and twenty (150) broiler chickens were purchased from a reputable hatchery and reared as a single group from day old to fourteen (14) days with appropriate vaccination/medication schedules. Certified commercial feed but without anticoccidial agents of any kind was served *ad libitum*.

2.6 Infection and monitoring

Each bird was infected with 0.2 ml of a suspension of sporulated *Eimeriatenella* containing about 50,000 viable oocysts per millilitre as illustrated by Gotepet *et al.*, [21].

After confirmation of the disease manifestation, the broilers were distributed into 9 groups consisting 15 birds each. Extract was administered at 100mg/kg, 125mg/kg and 165mg/kg to group 1-3. Groups 4, 5 and 6 were the negative control, uninfected untreated and positive control groups respectively. Each treatment group were treated with the plant extracts, once a day for five (5) days per os.

Blood samples were collected for haematology and serum biochemistry. The intestine (caecum) and liver were harvested for histological examination. All collated data from the study were subjected to Analysis of Variance (ANOVA) using the SPSS version 22.

3. RESULTS AND DISCUSSION

3.1 Eimeria oocysts count

Table I: Therapeutic effects of *A. muricata* leaf extracts on the Macmaster oocysts count of broiler chickens infected with *Eimeria* oocysts.

	Pre-infection (wk 4)	Post-infection (wk 5)	Post-treated (wk 7)	% change from 5th week to 7th week
AnM100	175.00±0.00	718.75±44.19 ^b	200.00±17.68 ^a	-60.23±0.32 ^a
AnM125	200.00±70.71	931.25±397.75 ^{bc}	360.25±26.52 ^a	-52.58±14.84 ^a

AnM165	200.00±35.36	843.75±61.87 ^{bc}	300.25±8.84 ^a	-59.60±3.94 ^a
NiF	137.50±53.03	131.25±8.84 ^a	127.00±0.00 ^a	-3.10±0.00 ^a
INT	112.50±17.68	1368.75±150.26 ^c	1400.34±471.41 ^b	11.93±46.61 ^b
ITT	112.50±35.36	712.50±88.39 ^b	210.25±26.52 ^a	-71.60±7.38 ^a

Means on the same column with different superscripts are significantly different (P<0.05) Percentage change in oocysts count between period of infection (week 5) and period of treatment indicates reduction in oocysts due to the extracts administered. Though without significant differences along the column, the control drug had the highest percentage reduction (70%) while the two plants extracts posed same efficacies.

KEY

AnM100 *Annonamuricata* extract at 100mg/kg
AnM125 *Annonamuricata* extract at 125mg/kg
AnM165 *Annonamuricata* extract at 165mg/kg
NiF uninfected untreated
INT infected untreated
ITT infected and treated with toltrazuril at 10mg/kg

3.2 Haematological indices

Table II: Percentage change in haematological indices

	PCV(%)	HB CONC(g/l)	RBC COUNT(10 ¹² /l)	BASOPHILS (%)	HETEROPHILS (%)	MONOCYTES (%)	LYMPHOCYTES (%)	EOSINOPHIL (%)
AnM100	21.74±6.15 ^{abc}	16.05±9.00 ^a	92.39±81.47 ^a	-78.41±4.82 ^a	-21.15±0.00 ^a	-85.00±2.36 ^a	-38.10±6.74 ^a	-100.00±0.00 ^a
AnM125	6.55±2.52 ^{ab}	17.47±11.24 ^a	36.71±44.84 ^a	-70.84±5.89 ^a	-7.32±4.80 ^a	58.59±20.00 ^{abc}	-16.67±0.00 ^{ab}	-95.42±0.59 ^a
AnM165	12.70±11.81 ^{ab}	7.64±8.78 ^a	76.32±33.50 ^a	77.09±14.73 ^a	136.67±146.14 ^a	-56.82±9.64 ^{abc}	-52.57±19.95 ^a	-96.67±4.72 ^a
NiF	28.03±7.50 ^{bc}	3.75±3.01 ^a	63.90±26.09 ^a	-69.05±3.37 ^a	-18.73±9.51 ^a	-64.11±3.63 ^{abc}	-47.78±11.00 ^a	-97.62±3.37 ^a
INT	0.00±7.07 ^a	19.34±61.03 ^a	-16.56 ±16.98 ^a	0.00±0.00 ^b	5.60±13.71 ^a	-28.28±24.28 ^{bc}	-15.00±7.07 ^{ab}	-14.53±13.12 ^a
ITT	1.79±2.52 ^a	-0.53±3.54 ^a	1.35±5.35 ^a	0.00±0.00 ^b	-6.55±3.60 ^a	-20.00±28.28 ^c	29.17±5.90 ^b	150.00±212.13 ^b

Values on the same column with different superscripts are significantly different (P<0.05)

3.3 Biochemical indices

Table III: Percentage change in biochemical indices

	AST(IU/L)	ALT (IU/L)	ALP (IU/L)	TP(g/dl)
AnM100	-65.51±54.43 ^b	-14.53±56.22 ^a	-42.76±17.18 ^a	125.54±81.74 ^b
AnM125	-89.94±24.34 ^{ab}	-20.15±12.68 ^a	-44.57±7.69 ^a	151.31±5.22 ^b
AnM165	-92.71±33.88 ^{ab}	-21.24±16.84 ^a	-35.96±0.35 ^a	164.62±6.53 ^b

NIF	-114.80±93.55 ^a	-37.55±16.33 ^a	8.81±4.12 ^b	143.33±61.28 ^b
INT	50.33±8.95 ^c	10.18±32.87 ^b	3.84±27.22 ^b	-6.72±25.99 ^a
ITT	-30.66±5.79 ^b	-17.18±30.67 ^a	-20.64±21.02 ^a	5.92±20.62 ^{ab}

Values on the same column with different superscripts are significantly different (P<0.05)

3.4. Discussion

Coccidiosis is a major source of concern to most especially poultry farmers around the world due to its difficulty to control, high cost of treatment and high risk of mortality following carcass emaciation and immunosuppression. Infections with species of the genus *Eimeria*, the causative agent of coccidiosis in poultry has been shown to be due to *Eimeriatenella*, *Eimerianecatrix*, *Eimeriabrunette* and *Eimeriaacervulina* in Nigeria [22].

Chemoprophylaxis and anticoccidial feed additives have been used in managing coccidiosis but emergence of drug resistant strains due to indiscriminate anti-coccidial drugs use [22] has complicated the management of coccidiosis on farms.

Totravet drug proved to be most effective in the treatment of coccidiosis among all drugs used because it treated the largest percentage of infected birds [7].

In this research, the Lethal Dose 50 (LD₅₀), efficacy parameters, haematological effect, liver function and phytochemical tests of the leaves are extrapolated. Infected birds displayed various clinical signs such as nutrients blood in faeces, anorexia, impaired body weight gain, and in some cases, mortality.

Phytochemicals analysis reveals that *Annonamuricata* both possesses tannins, flavonoids, phenolics, saponins. The presence of these phytochemicals in *Annonamuricata* has explored and utilised in them as antistress, anticancer agents, antifungi[23], antibacterial [24]. These phytochemicals present are believed to reduce the oxidative stress posed by the *Eimeria* organism.

The Lethal Dose 50 carried out following guidelines as laid down by OECD showed that *Annonamuricata* has no toxic effects on the birds at 2500mg/kg body. Hence, it can be considered non-toxic and safe.

Compared with the uninfected group with decreased percentage oocysts count of (-51.00%), no significant difference was observed with *Annonamuricata* extract-100mg/kg, 125mg/kg, 165mg/kg and the control drug (toltrazuril) groups. Whereas, infected untreated birds had an increased percentage OPG count of 11.93%, compared with the initial OPG.

This implies that the oocysts population available for continuation of the *Eimeria* life cycle had been reduced by the plants extracts but unhindered in the infected untreated group, though, numerically, the toltrazuril group, with decreased %OPG of -70.60 was more effective in cutting down the *Eimeriaspp* population. Hence, the effectiveness of extract can be said to be dose dependent.

Haematological Parameters

The coccidia agent had a deteriorative effect on the PCV, HB, RBC, heterophil, the basophil, monocyte, lymphocyte and eosinophil. Post infection, there was a significant decrease in the PCV of all the treatment groups compared to the Uninfected (control) group associated with caecal haemorrhage following *Eimeria* attack as reported by Akhtar, [25].

The Red Blood Cell count, PCV, and Hb decreased post infection but improved post treatment with *Annonamuricata* extract. Post treatment, the RBC count reflected no significant difference between the treatment groups and the uninfected groups. This infers that the previous sources of blood loss, caecal haemorrhage has been amended by the plant extract.

Similar to the observations of Akhtar[25] and Davies *et al.*, [26], the post-infection heterophil population of every other group was significantly higher than that of the uninfected groups indicative of infestation's immunosuppressing activities.

Post infection and post treatment, a significantly high level of eosinophils was recorded in the infected untreated groups. This is indicative of reduction in the level of damage done by the *Eimeria* organism.

In this experiment, infested broilers with coccidian parasites showed highly significant increase in serum ALT and serum AST level as compared with control group (NiF) which is line with Patraet *al.*,[27] and Mondalet *al.*,[28] who reported that liver function test of the infected broiler chicken with *Eimeria* spp. showed a significant increase in the serum ALT, AST. This is suggestive that a significant assault was inflicted on the cell lining of the caecal wall and their inflammation is seen from the increased and severe blood loss causing tissue loss from the body may attribute to increased AST activity.

The highest reduction rates were noticed in AnM 165, -107% and -89% of each plants respective and this translates to the absence of as many damaged tissues (intestinal wall) as there was in the post-infection stage. It can also be said that the extracts helping in facilitating ulcer healing.

Haemogram of the pre-extract, post-extract administration and Post-infection stages reflected that all the haematological indices, except the eosinophils at the post-infection stage, had no significant differences when compared with the control, the NiF group.

The eosinophil levels, at the post-infection stage, are significantly highest in the AnM 100 and least in AnM 125. The lower the value of the eosinophil population, the more efficiency the extracts in ameliorating the detrimental effects of the coccidial organism. From this result, it suggests that the plants extracts, *Annonamuricata* 125mg/kg (AnM 125) had the highest anticoccidial potentials of the doses in use.

4. CONCLUSION

Treatment with *Annonamuricata* decreased percentage oocysts count, improved the Red Blood Cell count, PCV, and Hb while increasing serum ALT and serum AST levels in the extract treatment groups compared with the untreated. Based on this result and characteristics displayed by the *Annonamuricata* extract, the plant can be said to possess anticoccidial property. Further purification, isolation and identification of bio active component from the plant is therefore recommended.

CONSENT

Not applicable.

ETHICAL APPROVAL

All of the animals used in this study received humane care according to the criteria outlined in the Guide for the Care and the Use of Laboratory Animals prepared by the National Academy Science and published by the National Institute of Health (USA). The ethic regulations have been followed in accordance with national and institutional guidelines for the protection of animals' welfare during experiments.

REFERENCES

- [1] Raza QS, Saleemi MK, Gul S, Irshad H, Fayyaz A, Zaheer I., et al. Role of essential oils/volatile oils in poultry production—A review on present, past and future contemplations. *Agrobiol. Rec.* 2022;7:40–56. doi: 10.47278/journal.abr/2021.013.
- [2] Degla LH, Kuiseu J, Olounlade PA, Attindehou S, Hounzangbe-Adote MS, Edoor PA., et al. Use of medicinal plants as alternative for the control of intestinal parasitosis: Assessment and perspectives. *Agrobiol. Rec.* 2022;7:1–9. doi: 10.47278/journal.abr/2021.011.
- [3] Khater HF, Ziam H, Abbas A, Abbas RZ, Raza MA, Hussain K., et al. Avian coccidiosis: Recent advances in alternative control strategies and vaccine development. *Agrobiol. Rec.* 2020;1:11–25. doi: 10.47278/journal.abr/2020.004.
- [4] Ndjonka D, Rapado LN, Silber AM, Liebau E, Wrenger C. Natural products as a source for treating neglected parasitic diseases. *Int. J. Mol. Sci.* 2013;14:3395–3439. doi: 10.3390/ijms14023395
- [5] Seifert H. *Tropical Animal Health*. Kluwee academic Publishers, Boston.2006; p57.
- [6] Joyner L, Long P. The specific characters of the Eimeria, with special reference to the Coccidia of the fowl. *AvianPathology.* 2008; 3(3):145–157.
- [7] Adewole S. The Efficacy of Drugs in The Treatment of Coccidiosis In Chicken In Selected Poultryes. *Academic Research International.* 2012.
- [8] Barbour E, Ayyash D, Iyer A, Harakeh S, Kumosani T. A Review of Approaches Targeting the Replacement of Coccidiostat Application in Poultry Production. 2015.
- [9] Kasilo OM, Trapsida JM, MwikisaNgenda C, Lusamba-Dikassa PS. World Health Organization An overview of the traditional medicine situation in the African region. *Afr. Health Monit.* 2010;1:7–15
- [10] Coria-Téllez AV, Montalvo-González E, Yahia EM, Obledo-Vázquez EN. *Annonamuricata*: a comprehensive review on its traditional medicinal uses, phytochemicals, pharmacological activities, mechanisms of action and toxicity. *Arabian Journal on Chemistry.* 2016;50(2):10–18
- [11] Moghadamtousi SZ, Fadaeinasab M, Nikzad S, Mohan G, Ali HM, HA, Kadir. *Annonamuricata* (Annonaceae): A review of its traditional uses, isolated acetogenins and biological activities. *International Journal of Molecular Science.*2015; 16: 15625-15658.
- [12] Leatemia JA, Isman MB. Insecticidal activity of crude seed extracts of *Annona* spp., *Lansiumdomesticum* and *Sandoricumkoetjape* against lepidopteran larvae. *Phytoparasitica.* 2004; 32: 30–37
- [13] Isman MB, Akhtar Y. Plant natural products as a source for developing environmentally acceptable insecticides,” in *Insecticides Design Using Advanced Technologies*, eds I, Ishaaya, F. Nauen, A. R Horowitz (Berlin; Heidelberg: Springer-VerlagNeitherland). 2007; 235–248.
- [14] Adewole S, Ojewole J. Protective effects of *Annonamuricata* Linn. (Annonaceae) leaf aqueous extract on serum lipid profiles and oxidative stress in hepatocytes of streptozotocin-treated diabetic rats. *African Journal on Traditional Complementary Alternative Medicine.*2009; 6: 30–41.
- [15] Mishra S, Ahmad S, Kumar N, Sharma B. *Annonamuricata* (the cancer killer): a review. *Global Journal of Pharmaceutical Resources.* 2013;2: 1613–1618

- [16] Durojaye CO, Osho IB. Therapeutic antitrypanosomal activity of *Alchornealaxiflora* leaf extract in rats experimentally infected with *Trypanosomabruceibrucei*. *European Journal of Pharmaceutical and Medical Research*. 2023; 10(2):193-198.
- [17] Sati SC, Kumar P. Assessment of Himalayan juniper, *Juniperussquamata* Buch-Ham ex D. Don for phytochemical screening and antimicrobial potential against some infection causing pathogens. *World Journal of Pharmaceutical Research*. 2015; 23: 998–1011.
- [18] Loman AA, Ju LK. Enzyme-based processing of soybean carbohydrate: Recent developments and future prospects. *Enzyme and Microbial Technology*. 2017; 106:35-47.
- [19] Sorescu AA, Nuta A, Ion RM, Iancu L. Qualitative Analysis of Phytochemicals from Sea Buckthorn and Gooseberry. *InTech*. doi: 10.5772/intechopen.77365. 2018.
- [20] Enevide C, David A, Fidelis S. A new Method for determining Acute toxicity in animal models. 2013.
- [21] Gotep JG, Tanko JT, Forcados GE, Muraina IA, Ozele N, Dogonyaro BB., et al. Therapeutic and safety evaluation of combined aqueous extracts of *Azadirachta indica* and *Khayasenegalensis* in chickens experimentally infected with *Eimeria* oocysts. *J. Parasitol. Res*. 2016; 9. [10.1155/2016/4692424](https://doi.org/10.1155/2016/4692424)
- [22] Ruff, MD. Resistance of coccidian to medications. In: proc. World's poultry congr, 2006; 2: 427-430.
- [23] Temitope OO, Thonda OA. Comparative Study of the Antibacterial And Antifungal Spectrum, Phytochemical Screening And Antioxidant Potentials Of *Alchornealaxiofolia* And *Piliostigmareticulatum* Leaf On Pathogenic Isolates. 2016.
- [24] Osuntokun OT, Olajubu FA. Antibacterial and Phytochemical Properties of some Nigerian Medicinal Plants on *Salmonella typhi* and *Salmonella paratyphi* Isolated from Human Stool in Owo local Government, Ondo State, Nigeria. *Journal of Scientific Research & Reports*. 2015; 4(5): 441-449.
- [25] Akhtar M, Awais M, Anwar M, Ehtisham-ul-Haque S, Amar N, Saleemi M, Ashraf K. The effect of infection with mixed *Eimeria* species on haematology and immune responses following Newcastle disease and infectious bursal disease booster vaccination in broilers, *Veterinary Quarterly*. 2015; 35(1): 21-26.
- [26] Davies RH, Carrique-Mas JJ. *Salmonella* Enteritidis in commercial layer flocks in Europe: legislative background, on-farm sampling and main challenges. *Rev. Brasileira de Ciência Avícola*. 2008; 10 (1): 01-09.
- [27] Patra G, Khan MD, Ayub A, Chanu V, Jonathan L, Joy LK, Prava M, Ravindran R, Das G, Inaotombi DL. PCR Based Diagnosis of *Eimeriatenella* Infection in Broiler Chicken. *International Journal of Poultry Science*. 2010; 9 (8): 813-818.
- [28] Mondal DK, Chattopadhyay S, Batabyal S, Bera AK. Plasma biochemical indices at various stages of infection with a field isolate of *Eimeriatenella* in broiler chicken. *Veterinary World*. 2011; 4(9):404