

Original Research Article

Baby corn based intercropping system in enhancing the growth and production of winter season baby corn in Chhattisgarh plains

improve the title, it seems redundant

Formatted: Font color: Red

Abstract

An experiment was carried out during winter season (*rabi*) for a period of two years 2019-20 and 2020-21 on Instructional cum Research Farm, Indira Gandhi Krishi Vishwavidyalaya, Raipur (C.G) with the objective to enhance the production of baby corn through intercropping. The experiment laid out in randomized block design (RBD) with eight treatments consisting of sole and paired row baby corn along with a combination of various legumes. The treatments were T1 : sole baby corn (45 x 20 cm), T2 : paired row baby corn (30/60 cm) T3 : sole fenugreek (30 x 10 cm), T4 : sole pea (30 x 20 cm), T5 : sole cowpea (30 x 20 cm) T6 : paired row baby corn + pea (2:2), T7 : paired row baby corn + fenugreek (2:2) and T8 : paired row baby corn + cowpea (2:2). The result of the experiment was concluded that growth attributes was highest under the treatment of T6 : paired row baby corn + pea which was closely followed by T8 : paired row baby corn + cowpea. Also, T6 : paired row baby corn + pea maximized the baby corn cob and fodder yield and therefore pea proved to be a valuable intercrop with baby corn in enhancing production of baby corn.

Keywords : baby corn, cob yield, fodder yield, intercropping, legumes, RBD

Introduction

it is necessary to carry out a review of current and extensive literature, it is necessary to strengthen this introductory part

Formatted: Font color: Red

Baby corn (*Zea mays* L.) is the unfertilized young cob of the corn which is mainly used for vegetables and soup. Baby corn is dehusked immature maize ear, harvested within 2-3 days of silking but prior to fertilization (Pandey *et al.*, 1998). The valuable size of baby corn is 6-11 cm length and 1-1.5 cm diameter with regular row/ovule arrangement. Generally, creamish - yellow to very light - yellow colored baby corn is top most priority to the consumer (Pal, 2011). Corn is

starchy in nature due to its carbohydrate content, but baby corn constitutes non-starchy one. It appends several health benefits as it is highly nutritive crop having capacity to convert more nutrients to food (Yasu, 2016). It contains folic acid, vitamin A, B, C, phosphorus, calcium and zinc which is used as supplements in various preparations such as vegetable, salad, soup, pickle, kheer, murabba, chutney and manchurian. The nutritive value of Baby corn; 100 g of baby corn consists of 89.1% moisture, 0.2 g fat, 1.9 g protein, 8.2 mg carbohydrate, 0.06 g ash, 28.0 mg calcium, 86.0 mg phosphorus, 11.0 mg ascorbic acid (Das *et al.*, 2009).

Therabi season is generally dominated by crops like wheat / chickpea / safflower / vegetable / pulse crops. In order to feed the growing population, there is an alarming rate to enhance the productivity per unit area per unit time which can be excelled by enhancing the cropping intensity especially under irrigated area to fulfill the optimum food requirement of the country. Corn is the most popular crop among the farmers in the state, however baby corn may also be adopted by farmers to raise their farm income and standard of living. Intercropping system is believed to enhance production per unit land area per unit time. Baby corn is a wide spaced crop however, possibility for introduction of inter row crops could be more income fetching. Verge of crop combination is applicable to maximize the utilization of growth resources per unit area and to improve the yield as well as to keep the soil in fruitful conditions (Shennan, 2008). Hence, sustainable productivity of crops is the need of the hour in the present context of Indian farming. So, we must explore every possibility for crop intensification with sustainable nutrition management for achieving the sustainability. It is necessary to introduce the reader to the crops that can be used in combination and why the choice of these.

Formatted: Font color: Red

METHODS AND MATERIALS

In this section it is necessary to clarify exactly the treatments, as well as the number of repetitions, it is also necessary to clarify what parameters were measured, in the field and in the laboratory (if that were the case), and put what techniques were used for this. Likewise, indicate which statistical analyzes were used to better understand the differences between the treatments. It is also necessary to clarify how the sampling was carried out, how many plants were used to make the measurements, among others,

Formatted: Font color: Red

The experiments were conducted at Instructional cum Research Farm of Indira Gandhi Krishi Vishwavidyalaya, Raipur (C.G.) India. The Instructional cum Research Farm is situated at 21°4'N latitude, 81°39'E longitude and 298 m altitude from the sea level. The region falls under the eastern plateau and hill region (Ago-climatic zone-7) of India.

The soil of the experimental field was *Vertisol*. The soil was neutral in reaction. The available nitrogen, phosphorus and potassium content were low, medium and high, respectively. Baby corn G5414 hybrid, Green pea variety Dhanya green pearl, Cowpea variety Ankur Gomati and fenugreek variety Meghna were taken as test crop during the course of investigations.

Detasseling, is an important practice in baby corn production. It was done immediately after the appearance of the tassel to avoid pollination and fertilization. Detasseling was done by holding the tassels firmly with hand and giving an upward jerk. This operation is necessary for getting good quality baby corn. Harvesting of baby corn was done after 3 – 4 days of silk emergence. Harvesting was done manually treatment wise by removing the cobs from the plants. In all there was 3 pickings at an interval of 1 – 3 days for complete removal of cobs. Harvesting was usually done in the morning when moisture percentage was high and temperature was low.

Result

By not knowing the number of repetitions, the results do not show a standard deviation or standard error, therefore it is risky to know if the differences are due to the treatments and not for some other reason.

Formatted: Font color: Red

Growth attributes

The intercropping with different pulses had a significant effect on the growth attributes of baby corn. This was marked by a gradual increase in plant height, number of leaves, leaf area and LAI and dry matter accumulation plant^{-1} . Treatment T6 : paired row baby corn + pea produced significantly the taller plants and higher number of leaves in comparison with the other treatments. Also the other growth attributes viz. leaf area plant^{-1} , Leaf Area Index and dry matter accumulation plant^{-1} were highly significant with the treatment of T6 : paired row baby corn + pea. The treatment of T8 : paired row baby corn + cowpea and the treatment of T2 : paired row baby corn was statistically at par to T6 : paired row baby corn + pea and was significantly superior over other treatments. The treatment of T1 : sole baby corn was found to

Formatted: Highlight

Formatted: Highlight

be inferior with the least growth attributes i.e. plant height, number of leaves plant⁻¹, leaf area plant⁻¹, LAI and dry matter

Formatted: Highlight

It is not possible to speak of significance, there is no standard deviation and Tukey's test

Formatted: Font color: Red

accumulation. This seems possible due to proper allocation of space with optimum plant population combined with increased plant height provided more number of nodes and thus more number of leaves were generated under the treatment of T6 : paired row baby corn + pea, T8 : paired row baby corn + cowpea and T2 : paired row baby corn. Similarly, higher leaf area plant⁻¹ and LAI were as a result of optimum utilization of growth resources due to proper space allocated for individual plants. The dry matter increased due to the cumulative effect of growth attributing features like plant height, number of leaves, leaf area which plays a very important role in proper growth of plant and thereby increasing dry matter. Leaf area index and dry matter of baby corn plant were highly correlated with the coefficient of 0.87 (Graph 1). Crop Growth Rate in general increased with the plant growth with the advancement of crop age upto 60 DAS and declined thereafter. The similar findings have been also reported by Singh et al. (2000), Ennin et al. (2002), Eskandari and Ghanbari (2009) and Hekmat (2016). The treatment of T6 : paired row baby corn + pea resulted in more number of baby cobs per plant with higher cob length and cob weight plant⁻¹ and was at par to treatment T8 : paired row baby corn + cowpea and T2 : paired row baby corn. Whereas the yield of intercrops like pea, cowpea and fenugreek were found to be superior in their sole planting as compared to intercrop treatments as the plant population under optimal condition leads to increase in yield. Mohan, *et al.* (2018), Mohan, *et al.* (2021).

It is necessary to add and strengthen a section for discussion of results before the conclusions that contrasts the results obtained here with other works.

Formatted: Font: Bold, Font color: Red

CONCLUSION

From the above study, it could be concluded that baby corn is a good crop for Chhattisgarh plains and when sowing in winter season, baby corn in paired row along with pea as intercrop could be a beneficial practice for achieving the sustainability and profitability in Chhattisgarh plain regions.

tables and graphs are required to edit

Formatted: Font color: Red

Formatted: Font color: Red

The tables and graphs must also contain the standard error or the standard deviation, in order to know the heterogeneity of the results.

Formatted: Font color: Red

Table 1: Growth attributes of baby corn as influenced by different intercrop

Treatment	Plant height (cm)	LAI	DMA (g plant ⁻¹)
T1: Sole baby corn	202.21	4.54	144.6
T2 : paired row baby corn	205.8	4.69	154.2
T6 : paired row baby corn + pea	210.4	5.14	157.3
T7 : paired row baby corn + fenugreek	203.9	4.71	149.3
T8: paired row baby corn + cowpea	206.3	5.07	156.3
SEm±	1.56	0.13	1.58
CD (P = 0.05)	5.10	0.44	5.14

I think it's important not only to show the height data, weight of the treatments that were measured. It is necessary to measure some other variables that strengthen what was found

Formatted: Font color: Red

which is the object of making the correlation, it is neither in the materials nor described in the results

Formatted: Font color: Red

Graph 1: Correlation between LAI and dry matter accumulation plant⁻¹




Table2 :yield attributes and yield of baby corn as influenced by different intercrop

Treatment	Cob length (cm)	Cob weight (g)	Baby corn cob yield (t ha ⁻¹)
T1: Sole baby corn	7.72	11.21	1.60
T2 : paired row baby corn	8.15	12.40	1.79
T6 : paired row baby corn + pea	8.80	13.69	1.92
T7 : paired row baby corn + fenugreek	8.03	11.72	1.63
T8: paired row baby corn + cowpea	8.51	12.49	1.85
SEm±	0.22	0.54	0.81
CD (P = 0.05)	0.71	1.76	0.26

References

It is essential to carry out a review of the current and extensive literature

Formatted: Font color: Red

Ennin, S. A., Clegg, M. D. and Francis, C.A. 2002. Resource utilization in maize/ soybean intercrops, African Crop Science Journal 10(3) : 251 – 261.

Eskandari and Ghanbari. 2009. Intercropping of maize and cowpea as whole crop forage effect of different planting pattern on total dry matter production and maize (*Zea mays* L.) forage quality. NotulaeBotanicaeHortiAgrobotanicalClujNapoca, 37(2): 152-155.

Hekmat, A. W., Abraham, T., Joyan, A., Kumar, M. and Ahmadzai, M.A. 2018. Productivity and economics of certified organic baby corn (*Zea Mays L.*) as influenced by different form of organic manure and intercropping with pulses. International Journal of Agriculture Sciences, 10(21): 7468-7471.

Mohan, M. Khajanji, S. N. Pandey, N. Effect of crop geometry and integrated nutrient management on growth and yield of winter season baby corn. The pharma innovation. 10(9): 253-255

Mohan, M.Rathore A. L. and Jha S, K. 2018 Response of drip fertigation, intrarow seeding of legume and planting geometry on net return of summer maize. International Journal of Pure and Applied Bioscience. 6(2) 964-967.

Pandey, A.K., Prasad, Kamta, Mani, V.P., Chauhan, V.S. and Singh, P. 1998. Improved maize production technology for mid-hills of N-W Himalayas. Technical Bulletin, 11: 1-46.

Singh D. P., Rana N.S. and Singh R.P. 2000. Dry matter production and nitrogen uptake in winter maize (*Zea mays*) – based intercropping system under different levels of nitrogen. Indian Journal of Agronomy, 45 (4): 676.

UNDER PEER REVIEW