

Comparative Morphology, Anatomy and Phytochemistry of *Cyrtosperma senegalense* (Schott) Engl. and *Alocasia macrorrhizos* L. (Araceae)

ABSTRACT

Morphological, anatomical and epidermal studies were carried out on two species of Araceae, *Cyrtosperma senegalense* (Schott) Engl. and *Alocasia macrorrhizos* L. to investigate the taxonomic value of their similarities and differences. Morphological features were visually observed. Fresh specimens were dehydrated, wax embedded, mounted, microscopically observed and micrographed. Basic similarities were found in their leaf shape, venation, inflorescence and spathe, seeded fruits, scattered vascular bundles, possession of aerenchyma, and presence of stomata on the abaxial and adaxial leaf surfaces. Conversely, the presence of prickles and sparseness of raphide idioblasts containing a raphide bundle each in *C. senegalense* distinguishes it from *A. macrorrhizos* which has abundant raphides. Phytochemical screening shows differences in their alkaloids, saponin, triterpenoids, steroids and glycosides contents. Though these preliminary studies yielded data that revealed their relationship, further investigations of their cytology using electron microscopy and molecular biology are needed for more diagnostic data to distinguish them from each other.

Key words: Aerenchyma, *Alocasia macrorrhizos*, Araceae, *Cyrtosperma senegalense*, raphide bundles,

INTRODUCTION

Araceae, also called the aroid, arum or cocoyam family comprises monocotyledonous flowering plants with palmately, pinnately or pedately dissected commonly arrow-shaped leaves with sheathing base petioles. Their unique characteristic is their inflorescence spadix, an internal spike of crowded flowers on fleshy axis, which is partially enclosed in a spathe, an external modified leaf (Osuji, 2013). They are pollinated by small flies that get temporarily trapped in the spadix. They bear yellow or red juicy berry fruits with seeds. Most of their body parts, especially leaves,

petioles and tubers have milky saps with calcium oxalate crystals which make them irritable. (Mayo et al. 1997; Prychid and Rudall 1999; Osuji 2013).

They diverged from the Order Alismatales, most of which got extinct during the Oligocene climate cooling and some reaching Africa, Australia, South America and South-East Asia (Wilson and Morrison 2000; Nauheimer et al. 2012). They are divided into subfamilies; Aroideae, Lasioideae, Zamioculcadoideae, Monsteroideae, Pothoideae, Lemnoideae, Orontioideae and Gymnostachyoideae. *Cyrtosperma senegalense* is in the subfamily Lasioideae while *A. macrorrhizos* is in Aroideae. The genera *Alocasia*, *Colocasia* and *Xanthosoma* species have monoecious flowers while *Amorphophallus* and *Cyrtosperma* species have hermaphrodite flowers. (Ivancic and Lebot 2000; Renner 2014).

Cyrtosperma senegalense (Scott) Engl. (homotypic synonym of *Lasimorpha senegalensis* (Scott); herbaceous and found in swamps) and *A. macrorrhizos* (herbaceous and commonly called the giant taro) represent Araceae species growing in different environmental conditions with similarities and differences in many parameters. *Cyrtosperma senegalense* is indigenous to Nigeria (Tarawou and Young 2016) and is noted for its possession of prickles on petioles, peduncles and underside of main leaf veins. It grows up to 6 – 12 ft (1.83 - 3.66 m) tall (Petersen 1989). In southern Nigeria, swamp arum fruits are used to treat dysentery and gonorrhoea while the young leaves are used as ulcer remedy and vegetable in Gabon (Burkil 1985). According to Tarawou and Young (2016), the powdered seeds proved effective in the removal of Mercury ions from aqueous solutions.

Alocasia macrorrhizos which gets as big as 15 feet (4.57 m) tall and 10 feet (3.05 m) wide in optimum terrestrial growth conditions is native to Phillipines and Asian mainland and can be found along roads, waste places and forests. Its elephant ear-like leaves, jute (tilting) skywards instead of drooping (Mayo et al. 1997; Mayo and Bogner 2011). Though it is regarded as a weed in Africa and tropical America, it is cultivated for its edible tubers in Malaysia, Sri Lanka and Bangladesh (Space and Flynn 2000; Wagner et al. 2008; Lebot 2008). The tubers are thoroughly boiled to dematerialize the irritable calcium oxalate crystals before consumption (Nauheimer et al. 2012). It is frequently grown at home and outdoors as ornamental plant. The rootstock, which is

seen as cooling and diuretic is a mild laxative useful in treatment of inflammations, piles, rheumatism, constipation, jaundice and as an astringent (Moghal et al. 2014).

Vegetative and reproductive morphological information constitute important and diagnostic sources of evidence for classification (Smith 1996). Hence they are traditionally used as sources of discriminatory evidence at all taxonomic levels, particularly at the specific and generic ranks (Taia and El-Olayan 2003; Taia 2004). In the Araceae, most diagnostic morphological information include: floral and inflorescence forms, leaf shape and coloration pattern. Taia (1998) and Osuji and Nwala (2015) underscored the value of morphological and anatomical features in plant classification, especially the aroids. Architecture of the vascular system, idioblasts and epidermal structure (Osuji 2006), presence or absence of ergastic substances (Nyananyo and Osuji, 2007), trichomes, stomata, cuticles and leaf architecture (Osuji and Nwala, 2015; Aziagba *et al.*, 2016; Ekeke *et al.*, 2021) have been frequently used to draw systematic conclusions.

Calcium oxalate crystals, which irritate the skin, and inflame the oral cavity and mucous membranes are common in Araceae (Osuji 2006). Though Prychid and Rudall (1999) reported that the druse, a type of Calcium oxalate crystal, may function as main irritant in toxic organs of plants, Konyar et al. (2014) stated that the presence and type of calcium oxalate crystals is not absolutely correlated to the toxicity of plant organs. However, Osuji and Nsaka (2014) showed the presence of raphide bundles of calcium oxalate in the edible Nigerian aroids. This finding agrees with Prychid and Rudall (1999) and Osuji (2013) that crystals of calcium oxalate play a taxonomic role as their location, type, quantity, shape, frequency of occurrence and distribution may be used for both taxonomic classification and delimitation. These crystals have been found in specific tissues such as epidermis, airspaces, cortex, or distributed in all parts of the plant (Osuji and Ndukwu, 2005; Konyar et al. 2014); and have been implicated as waste product in plants, objects of defensive mechanism, stored products or by products of metabolism (Osuji and Ndukwu 2005).

MATERIALS AND METHODS

Plant Material

Samples of *A. macrorrhizos* were collected from the Bioresources Conservation area in the University of Port Harcourt while samples of *C. senegalense* were collected along New Calabar River in Rivers State. Voucher specimens of both plants UPH/P/240 (*A. macrorrhizos*) and UPH/238 (*C. senegalense*) were deposited at the University of Port Harcourt Herbarium.

Morphology

Vegetative and reproductive morphology of the two species were carefully observed, photographed and described.

Epidermal studies

Following the method of Osuji and Nwala (2015), upper and lower epidermal membranes of leaves of the two species were peeled. The peeled epidermes were kept in absolute ethanol till needed. The epidermes were rinsed with distilled water and stained with 0.1 % safranin solution. The peels were then mounted with glycerin on clean glass slides, covered with cover slip and sealed with nail hardener. They were observed and micrographed under the microscope.

Anatomy

Following the modified method of Ekeke *et al.* (2021), fresh parts (petiole, midrib, leaf and root) were fixed in formalin acetic acid (FAA) for 24 hours after which they were passed through alcohol series (30, 50, 70 and 95 % v/v) solution and stored in 100 % ethanol until use. The specimens were embedded in paraffin wax and sectioned. Thin sections of the petiole, midrib and root were obtained by free-hand sectioning using a new blade. Selected thin sections were de-waxed, stained with safranin, rinsed with distilled water and mounted on clean glass slides each with a drop of glycerin and covered with a cover slip. The slides were microscopically studied and micrographed.

Phytochemical screening

Phytochemical screening was done on fresh leaves, roots and fruits of *C. senegalense* and leaves, roots and tubers of *A. macrorrhizos*.

Test for alkaloids: A total of 5g of plant samples was pulverized and heated in 10 ml of 10 % H_2SO_4 for 5 minutes on water bath. The mixture was filtered, and to three different 2 ml of filtrate, 3 drops of Dragendorff's reagent, Mayer's reagent and Hager's reagent were added respectively. Precipitation indicates presence of alkaloids.

Test for saponin: A total of 2g of plant sample was pulverized and warmed in 5 ml of distilled water for about 5 minutes on water bath and filtered. The filtrate was shaken vigorously for 20 seconds and allowed to stand. Observation of persistent frothing indicates the presence of saponin.

Test for carbohydrates (Molisch's test): Approximately 2g of plant sample was pulverized, warmed in 5 ml of water on a water bath for 5 minutes and filtered while warm. A total of 1 ml of alpha naphthol solution was added. The test tube was slanted and 1 ml of conc H_2SO_4 was added. A violet-purple-brown colour at the interface indicates the presence of carbohydrates.

Test for triterpenoids (Lieberman-Burchard's test): A total of 2g of plant sample was pulverized and macerated in 5 ml of chloroform and filtered. About 1 ml of acetic anhydride was added to the filtrate followed by 2 ml of conc. H_2SO_4 down the side of the test tube. The appearance of pink-red color at the interface indicates the presence of triterpenoids.

Test for steroids (Salkowski's test): Plant sample weighing 2g was pulverized; macerated in 5 ml of chloroform and filtered; then 2 ml of conc. H_2SO_4 was carefully added down the side of the test tube. A brown color at the interface indicates the presence of steroid.

Test for glycosides (Kedde's test): Plant sample weighing 2g of was pulverized, macerated in 5 ml of chloroform and filtered. A total of 3 drops of Kedde A solution followed by one drop of Kedde B solution were added. An immediate purple or violet color indicates the presence of glycosides.

RESULTS

Morphology

Leaf Morphology

The two species have palmate leaves held by conspicuous sheathing base petioles, inflorescences that are enclosed in a spathe, and fruits borne in cross-pattern arrangements on their spadix. They share few leaf features but show distinct variation in leaf margin, base, colour and lamina orientation (Table 1). Whereas both *Alocasia macrorrhizos* and *Cyrtosperma senegalense* have entire leaf margin when young, the margin becomes undulate in *A. macrorrhizos* at maturity. Colour of the leaves are very distinct between the two species. The lamina of *A. macrorrhizos* is auriculate while that of *C. senegalense* is overlapped and acutely pointed. The leaves are yellow green in *A. macrorrhizos* but brown in younger and dark green in mature leaves of *C. senegalense*. The lamina of *A. macrorrhizos* is nearly vertical in orientation with leaf apex pointing upwards while that of *C. senegalense* is horizontal with apex bent downwards.




Plate 1: Morphological features of *A. macrorrhizos* (A) showing (Ai) leaf structure (Aii) smooth Petiole (Aiii) inflorescence and infructescence and (Aiv) off-white-coloured fruits; and *C. senegalense* (B) showing (Bi) leaf structure (Bii) prickly Petiole and (Biii) Infructescence and (Biv) maroon-coloured fruits.

The petiole (leaf stalk) shows distinct variation between the two species. The petiole of *A. macrorrhizos* is tubular without prickles while that of *C. senegalense* is angular with numerous prickles arranged along the angles. Both species have variable leaf sizes on one plant. The leaves of *A. macrorrhizos* reach 30 cm in length, are yellow-green in colour, wavy along secondary veins, lance-ovate and are arranged in rosette ascending order (Plate 1a) while leaves of *C.*

senegalense are horizontal and reach 20 cm in optimum growth. Some leaves of *C. senegalense* (Plate 1b) are sagitate, slightly hastate.

The fruits of both plants are similar in appearance to maize (*Zea Mays*) when stripped of the spadix. *A. macrorrhizos*' (Plate 4a) are milky green in colour, with pinkish star-like ornaments at the tips. The fruits were embedded in a whitish liquid. The fruits of *C. senegalense* (Plate 4b) are reddish brown and bigger. In the wild, the fruits of *C. senegalense* were observed to be eaten by animals. The seeds range from three to five in each fruit. At the time of plant collection for this study, seeds of *C. senegalense* were found to be light yellow and hardened while that of *A. macrorrhizos* had cream colour and were fragile. *A. macrorrhizos* has underground tubers while *C. senegalense* has rhizomes, with which they propagate.

Table 1: Morphological features of *A. macrorrhizos* and *C. senegalense*.

Morphological Feature	<i>Alocasia macrorrhizos</i>	<i>Cyrtosperma senegalense</i>
Habitat	Mesophytic	Swamp
Foliar Features		
Leaf Type	Simple	Simple
Leaf Venation	Palmate	Palmate
Leaf margin	Undulate (older leaves) /entire (younger leaves)	Entire
Leaf Shape	Sagitate	Sagitate/ hastate
Leaf Apex	Acute	Acute
Leaf base	Auriculate	Overlapped, acutely pointed
Phylotaxy	Spiral/ whorled	Spiral/ whorled
Leaf Colour	Yellow Green	Dark-green (older leaves)/ brown (younger leaves)
Leaf Lamina	Vertical with apex pointing up	Horizontal with apex pointing down
Other Morphological Features		
Prickles	Absent	Present
Petiole shape	Tubular	Angular
Lacunae in Petiole	Present	Present
Underground		

System	Tubers	Rhizomes
Type of flower	Imperfect	Perfect
Colour of	Yellowish brown to white	Dark Purple
Inflorescence		
Number of inflorescence on one plant	3-5	1
Lacunae	Laticiferous	Non-Laticiferous
Milky sap around		
Fruits	Present	Absent
Fruit smell	Weak	Strong
Fruits colour	Milky Green	Reddish Brown
Accessories on		
Fruits	Present	Absent
Colour of Spathe	Green	Purple/white/green streaks
Nature of seeds in		
Fruit	Soft	Hard
Number of seeds in fruit	3-5	3-5

Anatomy

Epidermal Anatomy: Epidermal structures were very prominent in both adaxial (upper) and abaxial (lower) epidermes of the leaves of both species (Plate 2). There was clear similarity in upper and lower epidermal features of each of the species; and clear distinction between the epidermes of the two species. The ordinary epidermal cells of the upper and lower epidermes of *A. macrorrhizos* were elongated with angular edges while those of *C. senegalense* were elongated but slightly more robust and rounded at the edges. There was clear absence of trichomes on the epidermal structures of both species. However, stomata in *A. macrorrhizos* were of various types which include: tetracytic, brachytetracytic and cyclocytic while stomata in *C. senegalense* were mostly anomocytic.




Plate 2: Epidermes of *A. macrorrhizos* and *C. senegalense*. a) Upper epidermis of *A. macrorrhizos*, b) Upper epidermis of *C. senegalense*, c) Lower epidermis of *A. macrorrhizos* and d) Lower epidermis of *C. senegalense*.

Leaf Anatomy: The leaves of both species are clearly dorsiventral (Plate 3) with well-defined single layer of upper and lower epidermes. The leaf lamina of both species has 2-3 layers of palisade mesophyll and 3-5 layers of spongy mesophyll cells. In transverse section, the leaf of *A. macrorrhizos* has less number and size of lacunae in the midrib (Plate 3). However, there are more laticifers in *A. macrorrhizos* than in *C. senegalense*. Several raphide bundles are present in *A. Macrorrhizos* where they are mostly jutting out from the interior walls of the laticifers than in *C. senegalense*. There are several vascular bundles scattered but larger and more numerous aerenchyma in *C. senegalense* than in *A. macrorrhizos* midrib.




Plate 3: Midrib Anatomy of *A. macrorrhizos* and *C. senegalense*. a) T.S. midrib of *A. macrorrhizos* showing fewer laticifers with raphide bundles and b) T.S. midrib of *C. senegalense* showing three large lacunae running through the midrib without raphide bundles.

Petiole Anatomy: The petioles of *A. macrorrhizos* and *C. senegalense* (Plate 4) are uniseriate (Plate 4). However, the epidermal cells in the petioles of *A. macrorrhizos* are more laterally extended than those of *C. senegalense* which are more angular inwards. Their vasculature is collateral close. There are more vascular bundles in the outer ground tissue of *A. macrorrhizos* than *C. senegalense*. The vascular bundles in *A. macrorrhizos* are also closer distributed than in *C. senegalense*. Larger aerenchyma were found in the petiole and midrib of *C. senegalense*. In both plants, there are well defined sclerenchymatous tissues at the xylem but phloem fibre is isodiametric in cross-section and more massive in *A. macrorrhizos* while that of *C. senegalense* is like V-shape inverted inward. Plate 4d shows the prickles at the angle of the petiole of *C. senegalense*. The petioles have laticifers, which in *A. macrorrhizos* contain several raphide idioblasts attached to their inside walls but none in *C. senegalense*.




Plate 4: Anatomical features of the petiole and leaves of *A. macrorrhizos* and *C. senegalense*. a) Transverse section of *A. macrorrhizos* with the presence of raphide bundles in raphide ideoblasts (black arrow head), b) T.S. petiole of *C. senegalense*, c) T.S. petiole *A. macrorrhizos* showing presence of raphide bundles each in a raphide idioblast (black arrow heads) sitting on the inside wall of a laticifer, d) T.S. petiole of *C. senegalense* showing angular edge of the petiole and absence of raphide bundles in lacunae.

Root Anatomy: The root, in transection, is round in both species. The roots of both plants show presence of epiblema, cortex, pith, meta xylem, proto xylem and phloem. *A. macrorrhizos*' (Plate 5) shows the presence of root hairs, casparian strip, higher number of meta- and proto-xylem.




Plate 5: TS Root of *A. macrorrhizos* and *C. senegalense* showing the centralized vascular system and variation in their tissue outlay.

Phytochemistry

Though leaves of both species contain saponins, carbohydrates and glycosides, *C. senegalense* has steroids in the leaves while *A. macrorrhizos* lacks steroids. A common phytochemical in their roots was triterpenoids. Saponins, carbohydrates and glycosides were present in the root of *A. macrorrhizos* and absent in *C. senegalense*. Based on this result, it is observed that the two plants show low mutuality in their chemistry.

Table 2: Summary of phytochemical screening results of *A. macrorrhizos* and *C. senegalense*.

S/N	Phytochemical Test	Species	Leaf	Root	Tuber
1	Alkaloids Drangendoff's/ Meyer's/ Hager's	A.m.	-	-	+
		C.s	-	-	-
2	Saponin Frothing	A.m.	+	+	-
		C.s.	+	-	-
3	Carbohydrates Molisch's/ Fehling's/ Charring	A.m.	+	+	+
		C.s.	+	-	+
4	Triterpenoids Liebermann-Buchard's	A.m.	-	+	+
		C.s.	+	+	-

5	Steroids	Salwoski's	A.m.	-	-	-
			C.s.	+	+	-
6	Glycosides	Kedde's	A.m.	+	+	-
			C.s.	+	-	-

A.m. = *Alocasia macrorrhizos*; C.s. = *Cyrthosperma senegalense*; + = presence and - = absence.

DISCUSSION

A. macrorrhizos and *C. senegalense* are informally referred to as wild cocoyams due to shared morphological features with cocoyams, especially *Xanthosoma* and *Colocasia* spp. The similarities found are particular to the Araceae as outlined by Burkil (1985), Petersen (1986), Ray (1990), Nauheimer et al. (2012) and Osuji (2013). A striking difference between the two species is that the leaves of *A. macrorrhizos*, though on the same plant, may differ in details. It is often the case that different species have characteristic leaf shapes which have been used to identify them.

Thermogenesis in the sterile appendix of the spathe, which helps to attract pollinators has been reported (Wagner et al. 2008). Reproductive mechanism is different between the two species as *A. macrorrhizos* reproduces both sexually by seeds and vegetatively through tubers and basal offset divisions (Flach 1996; Paul and Bari 2011). In cultivation, *A. macrorrhizos* rarely produces flowers hence the vegetative features are mostly used for identification (Osuji 2013). Observation of hermaphroditic flowers with fruits formed all over the spadix of *C. senegalense* agrees with the reports of Nauheimer et al. (2012) and Ivancic and Lebot (2000). Consumption of the fruits by animals in the wild can be a pointer to its medicinal or nutritive value.

Obvious presence of large lacunae in these species attest to long exposure to low oxygen levels, which can induce the formation of lacunae or large air spaces in roots, which also affects the anatomy. In large shrubs growing in marshland, transportation of oxygen from the shoot to the submerged roots is not effective through diffusion. For good aeration of these roots, pressurized internal gas flow is employed. Possession of hollowed midribs, petioles and stems by *C. senegalense* enforce easier transportation of oxygen to the roots. This, in addition to elongated

petioles and presence of aerenchyma in roots and rhizomes is an adaptations which it uses to deal with the low oxygen and changing water levels of swamps.

The presence of anomocytic stomata has been reported among Araceae species (Osuji and Nwala 2015). Variation in stomatal types is of taxonomic significance as similarity of stomatal features in related taxa reflect shared genetic background (Osuji and Nwala 2015). Hence occurrence of paracytic and brachyparacytic stomata in *A. macrorrhizos* as in *Xanthosoma maffafa* indicate close relationship between them.

Though Konyar et al. (2014) reported that the presence and type of calcium oxalate crystals is not absolutely correlated to the toxicity of plant organs, the observation of such ergastic oxalate substance in *A. macrorrhizos* explains its acidity and agrees with the report of Osuji, (2013) that connects acidity to occurrence of calcium oxalates in edible aroids.

The midrib and petiole of *A. macrorrhizos* possess laticifers with raphide bundles attached to their inner walls. The laticifers are responsible for the secretion of irritable milky sap or latex, reported to contain calcium oxalate crystals in *A. macrorrhizos*. These are absent in *C. senegalense*. The presence of irritable latex is a feature of the Araceae according to Mayo et al. (1997) and Osuji (2013). It is notable that occurrence of raphide bundles inside the laticifers implicate the raphide structures as storage forms of calcium oxalate. This evidence supports the observation of Okoli and Green (1987) that raphide bundles are storage facilities in which calcium is stored in the form of calcium oxalate. Similarly, Osuji and Ndukwu (2005) reported calcium oxalate as stored and useful products of metabolism, which are regularly translocated from old to young plant parts

Phytochemical contents such as: alkaloids, carbohydrates, saponin, and terpenes observed on crude extracts of *A. macrorrhizos* agrees with the findings of Moghal et al. (2014). According to Moghal et al. (2014), the methanolic extract of *A. macrorrhizos* depicts good anthelmintic potential and could be used for prevention of free radical-mediated diseases. The presence of these bioactive secondary metabolites singly or in combination may imply defensive functionality against microorganisms and insects. Phytochemical screening of *C. senegalense* shows the presence of saponins, carbohydrates, terpenoids, steroids and glycosides. Observations

in this study corroborate the claim of Onwukaeme et al. (2007) of absence of tannins and presence of reducing sugar in *C. senegalense*. The presence of phenolic compounds and flavonoids in this plant suggests antioxidative property and explains the usefulness of this plant in herbal medicine.

The habitats of the two species have, no doubt, induced structural and biochemical response in them to develop characters that support their survival. Large aerenchyma in *C. senegalense* has very significant value in keeping it buoyant in its aquatic habitat whereas relatively smaller aerenchyma and slightly more sclerenchyma in *A. macrorrhizos* reflect its mesophytic habitat that requires physical strength and less buoyancy. The constituent phytochemicals in the two species indicate their potential ethno-medicinal values.

REFERENCES

- Aziagba B, Okeke CU and Iiodibia C (2016) Taxonomic importance of morphology of seven varieties of *Vigna unguiculata* (L.) Walp. Cultivated in Awka Anambra State, South Eastern Nigeria. American Journal of Life Science Research 4(4): 131-135.
- Burkil HM (1985) *The useful plants of West Tropical Africa*. Royal Botanic Gardens, Kew. 319 pp.
- Ekeke C, Nichodemus CO and Ogazie CA (2021) Morphological and Anatomical Studies on *Ipomoea coccinea* L. (Convolvulaceae): A New Record from Nigeria. Asian Journal of Research in Botany 6(1): 1-8.
- Flach M (1996) *Plant resources of South-East Asia; Plants yielding non-seed carbohydrates*. Backhugs Publishers. 15-42
- Ivancic A and Lebot V (2000) *The genetics and breeding of taro*. Cambridge University Press. pp 194 .
- Konyar ST, Necla O and Dane F (2014) Occurrence, types and distribution of calcium oxalate crystals in leaves and stems of some species of poisonous plants. Botanical Studies 55: 32.
- Lebot V (2008) Tropical root and tuber crops: cassava, sweet potato, yams and aroids. Journal of plant research 121(1): 73-82.
- Mayo S and Bogner J (2011) The first evolutionary classification of Araceae; Engler's Natural System. In: Engler A: Early Events in Monocot Evolution.

- Mayo SJ, Bogner J and Boyce PJ (1997). The Genera of Araceae. Royal Botanic Gardens, Kew. U.K. London. pp 370.
- Moghal MM, Alam K, Nsirim S, Rasheed S., Banik S, Ibrahim M and Amin M (2014) Determination of biological properties of *Alocasia macrorrhizos*: A medicinal plant. World Journal of pharmaceutical research 3(9): 193-210.
- Nauheimer L, Metzler D and Renner S (2012) Global history of the ancient monocot family Araceae inferred with models accounting for past continental positions and previous ranges based on fossils. New phytologist 149(4).
- Nyananyo BL and Osuji JO (2007) Biosystematic characterization of *Sphenostylis stenocarpa* (Hochst ex A. Rich.) Harms. Nigerian Journal of Botany 20(2): 411-419.
- Okoli BE and Green BO (1987) Histochemical localization of calcium oxalate crystals in starch grains of yams (*Dioscorea* L.). Annals of Botany 60: 139-142.
- Onwukaeme DN, Ikuegbvweha TB and Asonye CC (2007) Evaluation of Phytochemical Constituents, Antibacterial Activities and Effect of Exudate of *Pycnanthus angolensis* Wedd Warb (Myristicaceae) on Corneal Ulcers in Rabbits. Tropical Journal of Pharmaceutical Research 6(2): 725-730.
- Osuji JO (2006) Microstructural characters of the Inflorescence bracts distinguish between *Musa sapientum* L. and *Musa paradisiaca* L. International Journal of Botany 2(1): 11-16.
- Osuji JO (2013) Probable functions of calcium oxalate crystals on different tissues of the edible aroids (*Xanthosoma* and *Colocasia* spp) in Nigeria. African Journal of Biotechnology 12(25): 3952-3956.
- Osuji JO and Ndukwu BC (2005) Probable functions and remobilization of calcium oxalate crystals *Musa* L. African Journal of Biotechnology 4(10): 1139-1141.
- Osuji JO and Nsaka IJ (2009) Histochemical localization and probable functions of calcium oxalates in Nigerian cocoyams. Nigerian Journal of Plant Protection 26: 91-98.
- Osuji JO and Nwala PC (2015) Epidermal and cytological studies on cultivars of *Xanthosoma* (L.) Schott and *Colocasia* (L.) Schott (Araceae). International Journal of plant and soil science 4(2): 149-155.
- Paul, K.K., and Bari, M.A. (2011). Genetic variability studies in giant taro (*Alocasia macrorrhizos*). Journal of Agriculture 9(2): 107-111.
- Petersen G (1989). Cytology and systematics of Araceae. Norwegian Journal of Botany 9:119-166.

- Prychid CJ and Rudall PJ (1999) Calcium oxalate crystals in monocotyledons: A review of their structure and systematics. *Annals of Botany* 84(6): 725-739.
- Ray TS (1990) Metamorphosis in the Araceae. *American Journal of Botany* 77(12): 1599-1609.
- Renner SS (2014) The relative and absolute frequencies of angiosperm sexual systems; dioecy, monoecy, gynodioecy and an updated online database. *Am J Bot* 101(10):1588-96. doi: 10.3732/ajb.1400196. Epub 2014 Sep 24.
- Smith JF (1996) Tribal relationships within Gesneriaceae: A cladistic analysis of morphological data. *Systematic Botany* 21: 497-513.
- Space JC and Flynn T (2000) Observations on invasive plant species in American Samoa. USDA Forest Service, Honolulu. 51.
- Taia WK (1998) Morphological characters in the leaf of species of Papilionoideae (Leguminosae) in Egypt. *Journal of Arab Biology* 5: 137-158.
- Taia WK (2004) Modern Trends in Plant Taxonomy. *Asian Journal of Plant Sciences* 4: 184-206.
- Taia WK (2004) Leaf characters within tribe Trifolieae (Family Leguminosae). *Pakistani Journal of Biological Science* 7: 1463-1472.
- Taia WK and El-Olayan HA (2003) Effect of habitats on the phenotypic characters of three wild species in El-Riyadh city. *Bioscience Research Bulletin* 19: 171-177.
- Tarawou T and Young E (2016) Studies of Hg(II) ions from aqueous solution using powdered swamp arum (*Larsimorpha senegalense*) seeds. *Journal of applied sciences and Environmental management* 20(2): 221-225.
- Wagner AM, Krab K, Wagner MJ and Moore AL (2008) Regulation of thermogenesis in flowering Araceae: The role of the alternative oxidase. *Biochimica et Biophysica Acta* 1777: 993-1000.
- Wilson K and Morrison D (2000) *Monocots: Systematics and Evolution*. [Edition unavailable]. CSIRO PUBLISHING. <https://www.perlego.com/book/1468700/monocots-systematics-and-evolution-systematics-and-evolution-pdf>.