Original Research Article

Study of nitrogen rates and nano urea effects on physiological parameters and yield of Safflower (*Carthamus tinctorius* L.)

.

UNDER PEER REVIEW

ABSTRACT

Safflower (*Carthamus tinctorius* L.) is an important oil seed crop, essential as vegetable oil for humans and feed for livestock. The experiment was conducted in a split plot design to study the effects of foliar application of nano urea and urea on safflower growth and development. The results showed that application of 100% nitrogen in the form of urea topdressing at vegetative stage and foliar spray of nano urea twice at flowering and seed filling stages significantly improved the physiological parameters *viz*. plant height (cm), primary branches of safflower. The CSI, RWC, Seed yield (kg/ha), 100 seed weight (g) and hull content (%) of safflower also significantly improved with 100% N and foliar spray of nano urea twice at flowering and seed filling stages. The result from the study suggests that application of urea as split doses and foliar spray of nano urea at reproductive stage is beneficial for the crop and will improve seed yield and oil quality of safflower. Foliar application of nano urea improves the nutrient uptake by the crop through its slow and controlled release of nutrients. It also helps in reducing environmental pollution due to leaching of fertilizers.

Keywords: *Nano urea, Safflower, Foliar spray, Chlorophyll stability index, Seed Yield*

1. INTRODUCTION

Safflower (*Carthamus tinctorius* L.) is an important *rabi* oil seed crop belonging to the family Asteraceae. Safflower is a strongly tap rooted, highly branched and thistle like herb with the ability to withstand drought, salinity and cold stress**.** The oil content of the seed ranges from 24-36 per cent.

Safflower oil is preferred for its higher poly unsaturated fatty acid (78% linoleic acid), and it is nutritionally similar to sunflower oil, which is very useful for reducing blood cholesterol content (1,2). Oleic safflower oil is highly preferred in the food, pharmaceutical, biofuel production, paints, lubricants, and cosmetics industries (3).

Plant nutrition is vital for agricultural production and fertilizer application accounts for 40% to 60% of global food production. Fertilizing with nitrogen is important as it determines yield due to its multi-dimensional effects on the growth and development of safflower (4). Nitrogen is required for the manufacture of several growth-promoting enzymes, proteins and so plays an important role in plant growth regulation, particularly during the vegetative phase (5). Excessive fertilizer applications lead to economic losses and discharge of excessive amount of nitrogen in the nitrate form through leaching (6). Agricultural use of inorganic fertilizers in 2019 was about 190 million tonnes of nutrients, of which 57% was nitrogen (7). Farmers were forced to increase the amounts of applied nitrogen fertilizers to achieve better crop production due to the high N loss and low use efficiency, which increased the costs of farming while also increasing the environmental implications (8, 9, 10). As a result, there is a compelling need to increase nitrogen availability for plants while avoiding adverse environmental impacts. Some studies proposed the use of nanoparticle fertilizers, particularly nanoparticle urea (NPU), to avoid the issues associated with the application of bulk urea while not depriving the plant of its benefits (8, 9,11). When compared to traditional fertilizers, nano fertilizers improve growth parameters (plant height, leaf area, number of leaves per plant), dry matter production, chlorophyll production and photosynthesis rate, resulting in increased photosynthesis and translocation to different parts of the plant (12, 13). Nano urea when applied to foliage, easily enters through stomata and distributed through phloem. Unutilized nitrogen is stored in plant vacuole and is slowly released for proper growth and development of plant (14). The lack of synchronisation between nutrient release from the fertilizer and plant need is the fundamental cause behind low nitrogen use efficiency of bulk chemical fertilizer's (10). The controlled release of nutrients provided by nanostructured fertilizers will allow for a longer effective duration of nutrient supply to the plant, (15) ensuring without any negative environmental consequences (16). Nitrogen fertilizer application to the soil may result in nutrient losses, particularly through leaching and denitrification; however, foliar spraying may reduce such losses to provide nutrients and can mitigate nutrient deficiencies (17). Nano Urea is an important tool

UNDER PEER REVIEW

in agriculture to improve crop efficiency, yield and quality parameters with increase nutrient use efficiency, reduce wastage of fertilizers and cost of cultivation (14).

2. MATERIALS AND METHODS

The experiment was conducted at the research farm of All India Coordinated Research Project (AICRP) on Safflower, VNMKV, Parbhani, Maharashtra during *rabi* season of 2021-22 on medium black soil. The experiment was conducted on PBNS 86 variety of safflower in a split plot design with three replications.

Each replication consists of three main plots which include F1 with 100% N (60kg N/ha), F2 with 75% N (45kg N/ha) and F3 with 50% N (30kg N/ha) and each main plot is divided into five sub plots, which include N0 (No spray of nano urea), N1 (nano urea spray at flowering stage), N2 (nano urea spray at flowering and seed filling stages), N3 (2% urea spray at flowering stage) and N4 (2% urea spray at flowering and seed filling stages) thus, each replication has total fifteen treatment combinations.

Table 1. Initial physio-chemical properties of experimental soil

FC%	33.00%	EC at 0-15 cm	0.53 ds/m
BD	1.28 Mg/m^3	EC at 15-30 cm	0.64 ds/m
Soil texture	Clayey	Available N (kg/ha)	157.86
Soil pH	7.71	Available P (kg/ha)	14.17
Organic carbon	0.73%	Available K (kg/ha)	619.81

The net plot size was 5.4 x 6.0m. The spacing is 45 X 20cm. One third of nitrogen was applied at the time of sowing and remaining 2/3rd in 2 equal splits at 30 DAS and 60 DAS as urea top dressing. The nano urea @ 3ml/litre water and 2% urea @ 20g of urea/litre water were applied as foliar spray at flowering and seed filling stages.

Biometric observations were recorded from five randomly selected plants from each plot periodically. The data for studying plant height, number of primary branches was recorded from each plot at 30 DAS, 60 DAS and 90 DAS.

2.1 Leaf Chlorophyll content and Chlorophyll stability index (CSI)

UNDER PEER REVIEW

Fresh leaves from each treatment were collected randomly for chlorophyll studies. Shoaf and Lium (1976) devised an improved method of extraction of chlorophyll by dimethyl sulphoxide (DMSO) (18). Weigh 100mg of fresh leaf and put in a test tube. Incubate the leaf in 7.0 ml of DMSO at 65^0 C for 30 minutes. After 30 minutes, decant the supernatant into a test tube and discard the leaf tissue. The final volume was made up to 10 ml with DMSO. The absorbance of the extract was measured at 645 and 663 nm in spectrophotometer using DMSO as blank.

$$\text{Total chlorophyll} = 20.2\,(A645) + 8.02\,(A663)\ X\ \frac{V}{1000 \times W \times a}$$

Where,

A = Absorbance at specific wavelength (645 and 663 nm)

V = Final volume of the chlorophyll extract (ml)

W = Fresh weight of the sample (g)

a = path length of light (1cm)

The Chlorophyll stability index was computed by using the method given by Dhopte and Livera (1989). The same procedure was followed for stressed leaves but in this case the leaf will be subjected to high temperature stress by placing the leaf tissue in a test tube containing 10 ml of distilled water. Place it in hot water bath and subject it to high temperature of 65^0 C for 1 hour. After 1 hour drain out the water, add 7.0 ml of DMSO and incubate it. Likewise, the unheated sample, same procedure is followed water bath and then record spectrophotometer readings for both leaf samples. The difference in two readings on absorbance (reading without heating and reading after heating of sample at 65^0 C control) is defined as chlorophyll stability index.

CSI (%) = (Chlorophyll content of treated sample/Chlorophyll content of control) X100

2.2 Relative Water Content (%)

Barrs and Weatherley (1962) given the method for estimation of relative water content. A physiologically functional leaf i.e., third from top was selected for RWC estimation. The fresh weight of leaf samples was immediately recorded. These leaf samples were kept floating on water for 4-6 hrs. depending upon the degree of imbibitions to get turgid weight. Finally, the same leaf samples were

UNDER PEER REVIEW

kept in an oven at 75^{O}C for assessing dry weight and the values of RWC were calculated by using following formula.

$$RWC = \frac{[\text{Fresh weight (Wf)} - \text{dry weight (Wd)}]}{[\text{turgid weight (Wt)} - \text{dry weight (Wd)}]} X\ 100$$

2.3 Hull content (%)

Weigh five grams seeds and soak them in water for 24 hours. Drain the water and separate the hull from seed by hand. Collect all the hulls and dry them in oven at 40^{0}C for 8 hours or dry them in sun for 2 days. Find the weight of oven/sun dried hulls. To find the hull content the following formula is used:

$$\text{Hull content (\%)} = \frac{\text{Weight of oven/sun dried hulls (g)}}{5} \times 100$$

The observations for Days to 50% flowering, Chlorophyll stability index, Relative water content were taken once during crop development stage. Yield parameters *viz*. 100 seed weight (g), seed yield per plant (g), seed yield (kg/ha), oil content and hull content were recorded at the time of harvest.

3. Results and Discussion

The results showed that application of nitrogen as two split doses of urea had significantly increased the physiological parameters. Plant height was recorded maximum with 100% N application at all growth stages *i.e.,* 30, 60, 90 DAS and at harvest. Similarly, number of primary branches/plants also showed a decrease with decreasing the dose of nitrogen up to 50% N and maximum were observed with 100% N application at 60 DAS, 90 DAS and at harvest (Table 3). The plants applied with N2 showed maximum plant height followed by N1 and N4 whereas, the plants treated with no foliar spray (N0) recorded minimum plant height. Highest number of primary branches were recorded in N2 treatment and lowest was recorded in N0 treatment (Table 3). Nitrogen fertilization helps in preventing premature death of leaves due to nutrient deficiencies (21).

3.1 Effect of nitrogen on Chlorophyll content :

UNDER PEER REVIEW

The results showed that split application of nitrogen in the form of urea had significant effect on the chlorophyll a, chlorophyll b and total chlorophyll content of safflower. Among the given nitrogen doses, 100% N (F1) application showed the highest (2.21 mg/g) chlorophyll content of leaves as compared to 50% N (F3) application (1.49 mg/g) as presented in table 2.

Similarly, maximum chlorophyll stability index of safflower was recorded with 100% N (F1) application followed by 75% N (F2) and 50% N (F3) application respectively as presented in table 2.

The maximum leaf chlorophyll content and CSI was recorded with foliar spray of nano urea twice at flowering and seed filling stages (N2) while, the lowest leaf chlorophyll content (1.48 mg/g) and CSI was recorded with no foliar spray of nano urea (N0). However, other treatments showed the effect lower than N2, but were higher when compared to the control. Application of urea as split doses and foliar spray of nano urea at reproductive stage showed higher chlorophyll content as compared to the control where only soil applied nitrogen was given which may be due to the decrease in applied dose of nitrogen as it is important for protein and carbohydrate metabolism and can enhance the plant height, production of photo-assimilates and distribution of assimilates to the reproductive organs (22) and ultimately greater production of dry matter (23) indicating a direct relation of nitrogen content with dry matter production. Nitrogen is essential for synthesizing the amino acids, nucleic acids, and chlorophyll. The result from the current study also indicates a positive relationship between chlorophyll content and nitrogen.

3.2 Effect of nitrogen on Relative water content (%) :

The effect of split application of nitrogen was significant on relative water content of safflower. Maximum RWC was observed with 100% N (F1) followed by 75% N (F2) and 50% N (F3) respectively as presented in table 2. Foliar application of nano urea and 2% urea at reproductive stage of crop significantly affected the relative water content of leaf. Among the given treatments, foliar spray of nano urea twice at flowering and seed filling stages (N2) recorded maximum RWC and lowest was recorded for control (N0). However, the other treatments recorded comparatively lower values than N2 but higher than control (N0).

With increase in nitrogen content, RWC also increased. Osmatic adjustment (OA) is a powerful mechanism of conserving cellular hydration under drought stress and RWC expresses the

UNDER PEER REVIEW

effect of OA in this respect. Hence, RWC is an appropriate estimate of plant water status in terms of cellular hydration under the possible effect of both leaf water potential and OA (20).

Table 2: Mean data for Chlorophyll content, Chlorophyll Stability Index (%), Relative Water content (%) and Hull content (%).

TREATMENTS	Chlorophyll content (mg/g)	CSI (%)	RWC (%)	Hull content (%)
F1	2.21	85.00	89.49	41.58
F2	1.87	84.48	87.63	40.63
F3	1.49	81.69	82.81	40.36
SE ±	0.02	0.44	0.14	0.16
CD at 5%	0.09	1.73	0.55	0.62
FOLIAR SPRAY				
N0	1.48	74.21	78.42	39.27
N1	2.06	87.73	89.38	41.51
N2	2.17	91.39	91.22	42.35
N3	1.65	80.98	86.34	40.21
N4	1.93	84.32	87.86	40.94
SE ±	0.03	0.58	0.31	0.19
CD at 5%	0.07	1.68	0.89	0.56
INTERACTION				
SE ±	0.04	1.00	0.53	0.33
CD at 5%	0.13	NS	1.55	NS
G. MEAN	1.86	83.73	86.64	40.85

UNDER PEER REVIEW

3.3 Effect of nitrogen on hull content (%) :

Application of different doses of nitrogen in the form of urea as split doses at vegetative stage showed significant effects on hull content of safflower seeds. The data obtained from the present

study showed that maximum hull content was recorded with 100% N (F1) while lowest was recorded with 50% N (F3) application as shown in table 2. Similarly, foliar spray of nano urea twice at flowering and seed filling stages (N2) recorded maximum hull content while, the lowest was recorded for no foliar spray of nano urea (N0). However, the other treatments recorded comparatively lower values than N2 but higher than control (N0).

3.4 Effect of nitrogen on Seed yield

The results showed that enhanced yield of a crop is often associated with optimum nutrient uptake, particularly nitrogen. Among the given treatments, the plants applied with 100% N at vegetative stage showed significantly maximum growth followed by 75% N and 50% N application.

Table 3: Mean data for 100 seed weight (g), Seed yield/plant and per plot, oil yield (kg/ha) and hull content (%).

TREATMENTS Nitrogen Fertilizer	100 seed weight (g)	Seed yield/plant (g)	Seed yield (kg/ha)	Oil yield (Kg/ha)
F1	6.42	86.47	14.69126	478
F2	6.03	81.27	10.45673	343
F3	5.82	78.60	9.89911	324
SE ±	0.11	0.73	0.622	23.60
CD at 5%	0.44	2.86	2.44	92.50
Foliar spray				
N0	4.88	73.33	9.458097	309
N1	6.56	84.56	11.8552	397
N2	7.10	91.00	16.11444	520
N3	5.76	79.67	10.54176	333
N4	6.14	82.00	10.44233	350
SE ±	0.09	0.69	0.372	15.60
CD at 5%	0.27	2.00	1.086	45.50
Interaction				
SE ±	0.16	1.19	0.644	27.00

UNDER PEER REVIEW

CD at 5%	NS	NS	NS	NS
G. Mean	6.09	82.11	11.68237	382

Foliar application of nano urea at the time of flowering and seed filling stages of safflower had increased the returns by promoting maximum yield through its slow and controlled release of nitrogen which is very beneficial for the plant in utilizing the nutrients and improving the yield attributes. Maximum mean values were recorded with N2 (Table 3) for the 100 seed weight. Seed yield (kg/ha) had significantly increased with N2 over N0. Some studies reported that higher yield in safflower is related to increase in number of heads/plant and number of seeds/heads in safflower (24). It has been reported that split application of nitrogen can improve the seed yield by maximising the number of capitula/plant (25) and the responding level of plant to urea fertilizer can be attributed to the needed nitrogen in soil (26).

It is evident that applying nano nitrogen fertilizers in conjunction with reduced dosses of conventional nitrogen fertilizers can boost the productivity of several cereal crops (6, 27) by their smart nutrient delivery system. Applying nitrogen more than required may reduce crop yields and is toxic to plants. So, providing optimum doses of nitrogen is essential for obtaining maximum yields. Through foliar application, nutrients can be supplied as per plant demand and can reduce fertilizer losses.

CONCLUSION

Application of 100% nitrogen fertilizer improved the growth and yield of safflower. When the nitrogen supply is reduced to half there is reduced growth and yield of safflower. The supply of nitrogen through foliar application of nano urea and 2% urea at reproductive stage provided the plant with required nutrients and enhanced the yield attributes and seed yield of safflower. Foliar application of nutrients is an effective way to mitigate nutrient deficiencies when plants are unable to absorb the nutrients directly from soil. Due to lack of synchronization between nutrient release from soil applied urea and its demand by the plant, resulted in less nutrient use efficiency by the crop. This can be managed by applying the nano urea which will release the nutrients in a slow and control manner to meet the plant's nutrient needs. Application of nano urea will reduce the cost of urea for crop production and will minimize pollution caused by conventional fertilizers.

UNDER PEER REVIEW

REFERENCES

1. Belgin C, Bilal G. and Mustafa K. Oil content and fatty acid composition of some safflower (*Carthamus tinctorius* L.) varieties sown in spring and winter. International Journal of Natural and Engineering Science. 2007;1(3), 11-15.
2. Kadu P. S. and Ismail S. Impact of organics and biofertilizers on yield, quality and uptake of major nutrients by safflower grown on vertisols. Annals of Plant Physiology. 2008; 22(2), 214-216.
3. Nogales-Delgado, S., Encinar, J. M., & Cortés, Á. G. High oleic safflower oil as a feedstock for stable biodiesel and bio lubricant production. Industrial Crops and Products. 2021;170, 113701. https://doi.org/10.1016/j.indcrop.2021.113701
4. Grant, C. Enhancing nitrogen use efficiency in dry land cropping systems on the Northern Great Plains. 18th World Congress of Soil Science, Philadelphia, USA; 2006.
5. Iqbal, M. A. Nano-fertilizers for sustainable crop production under changing climate: a global perspective. Sustainable crop production. 2019;8, 1-13. DOI:10.5772/intechopen.89089
6. Henke, J., Breustedt, G., Sieling, K. and Kage, H. Impact of uncertainty on the optimum nitrogen fertilization rate and agronomic, ecological and economic factors in an oilseed rape-based crop rotation. The Journal of Agricultural Science. 2007; 5, 455 - 468. DOI:10.1017/S0021859607007204
7. FAO. World Food and Agriculture - Statistical Yearbook 2021. Rome.
8. Rathnayaka, R. M. N. N., Iqbal, Y. B., & Rifnas, L. M. Influence of urea and nano-nitrogen fertilizers on the growth and yield of rice (*Oryza sativa* L.) cultivar 'Bg 250'. Influence of Urea and Nano-Nitrogen Fertilizers on the Growth and Yield of Rice (*Oryza sativa* L.) Cultivar 'Bg 250', International Journal of Research Publications. 2018;5(2), 7-7.
9. Chhowalla, M. Slow-release nano fertilizers for bumper crops. ACS Central Science. 2017;3(3), 156-157. https://doi.org/10.1021/acscentsci.7b00091
10. Marchiol, L. Nanofertilisers. An outlook of crop nutrition in the fourth agricultural revolution. Italian Journal of Agronomy. 2019;14(3), 183-190. DOI:10.4081/ija.2019.1367

UNDER PEER REVIEW

11. Kottegoda, N., Sandaruwan, C., Priyadarshana, G., Siriwardhana, A., Rathnayake, U. A., Berugoda Arachchige, D. M., and Amaratunga, G. Urea-hydroxyapatite nanohybrids for slow release of nitrogen. ACS nano. 2017;11(2), 1214-1221.
12. Ali, N. S., & Al-Juthery, H. W. A. The application of nanotechnology for micronutrient in agricultural production. The Iraqi Journal of Agricultural Science. 2017;48(4), 984.
13. Singh, M. D. Nano-fertilizers is a new way to increase nutrients use efficiency in crop production. International Journal of Agriculture Sciences. 2017;9(7), 3831-3833.
14. Prem Baboo. Nano Urea, the Philosophy of Future. Retrieved from - https://www.researchgate.net/publication/353260539 ;2021.
15. Liu, R. and Lal, R. Potentials of engineered nanoparticles as fertilizers for increasing agronomic productions. Science of the total environment. 2015;514, 131-139.
16. Kopittke, P. M., Lombi, E., Wang, P., Schjoerring, J. K. and Husted, S. Nanomaterials as fertilizers for improving plant mineral nutrition and environmental outcomes. Environmental Science: Nano. 2019;6(12), 3513-3524.
17. Samadhiya, V. K. Response of micronutrients and urea foliar spray on yield and nutrient uptake of safflower (*Carthamus tinctorius* L.) in Chhattisgarh plan. International journal of chemical studies. 2017;5, 325-35.
18. Shoaf, W. T., & Lium, B. W. Improved extraction of chlorophyll a and b from algae using dimethyl sulfoxide. Limnology and oceanography. 1976;21(6), 926-928.
19. Dhopte, A. M., & Livera-Muñoz, M. Useful techniques for plant scientists. Useful techniques for plant scientists; 1989.
20. Barrs, H. D., & Weatherly, P. E. Physiological indices for high yield potential in wheat. Indian Journal of Plant Physiology. 1962;25, 352-357.
21. Abdel-Hafiz, M., and Hassan, A. Morph physiological traits of safflower as affected by plant densities and nitrogen fertilization. Journal of Productivity and Development. 2019;24(1), 1-14.
22. Dordas, C. A. and Sioulas, C. Dry matter and nitrogen accumulation, partitioning, and retranslocation in safflower (*Carthamus tinctorius* L.) as affected by nitrogen fertilization. Field Crops Research. 2009;110(1), 35-43.

UNDER PEER REVIEW

23. Saitheja, V., Senthivelu, M., Prabukumar, G. and Prasad, V. B. R. Maximizing the Productivity and Profitability of Summer Irrigated Green gram (*Vigna radiata* L.) by Combining Basal Nitrogen Dose and Foliar Nutrition of Nano and Normal Urea. International Journal of Plant & Soil Science. 2022;109-116. DOI:10.9734/ijpss/2022/v34i2231362

24. Shahrokhnia, M. H. and Sepaskhah, A. R. Physiologic and agronomic traits in safflower under various irrigation strategies, planting methods and nitrogen fertilization. Industrial Crops and Products. 2017;95, 126-139. DOI:10.1016/j.indcrop.2016.10.021

25. Goud, G. M. K., Sudhakar, K. S., Pasha, M. L. and Madhavi, A. Evaluation of the Foliar Application of Nano Urea on the Performance of Rabi Sunflower (*Helianthus annuus* L.). International Journal of Environment and Climate Change. 2022;12(11), 2700-2706. **DOI:** 10.9734/ijecc/2022/v12i1131258

26. Mahmoodi, P., Yarnia, M., Rashidi, V., Amirnia, R. and Tarinejhad, A. Effects of nano and chemical fertilizers on physiological efficiency and essential oil yield of *Borago officinalis* L. Applied Ecology and Environmental Research. 2018; 16, 4773-4788. DOI:10.15666/aeer/1604_47734788

27. Benzon, H. R. L., Rubenecia, M. R. U., Ultra Jr, V. U. and Lee, S. C. Nano-fertilizer affects the growth, development, and chemical properties of rice. International Journal of Agronomy and Agricultural Research. 2015;7(1), 105-111

UNDER PEER REVIEW

UNDER PEER REVIEW