

Original Research Article

EFFECT OF FOLIAR NUTRITION ON YIELD, QUALITY, NUTRIENT CONTENT AND UPTAKE OF *KHARIF* COWPEA

ABSTRACT

A field experiment was conducted at the Agronomy Instructional Farm, C. P. College of Agriculture, SardarkrushinagarDantiwada Agricultural University, Sardarkrushinagar to investigate the “effect of foliar nutrition on yield, quality, nutrient content and uptake of *kharif* cowpea” during *kharif* season of 2020. In Randomize Block Design, eight treatment combinations were tested in four replications. Cowpea variety Gujarat cowpea 5 was planted at a distance of 45 cm × 10 cm. Application of 75% of RDF + 2% spray of each urea and urea phosphate at 15 and 30 DAS gave significantly higher yield (1422 kg/ha), protein content (22.93 %), nutrient content of nitrogen and phosphorus in seed (3.67 % and 0.446 %) and in stover (0.73% and 0.273% respectively) and nutrients uptake of nitrogen (71 kg/ha), phosphorus (13 kg/ha) and potassium (56 kg/ha) by cowpea crop it was at par with application of 75% of RDF + 2% spray of NPK 19:19:19 at 15 and 30 DAS.

KEYWORD :Yield, Quality, Nutrient content, Nutrient uptake and Cowpea

INTRODUCTION

According to **Sebethaet al. (2010)**, a 100g edible green cowpea pod contains 84.6 g of moisture, 4.3 g of protein, 0.2 g of fat, 0.9 g of minerals, 2.0 g of fiber, and 8.0 g of carbohydrates. and its seed serves as a nutritional supplement to human diets as well as animal feed. In India, pulses are grown nearly in 29.99 million ha with an annual production of 23.15 million tons and average productivity of 841 kg/ha (**DES, 2018**). The per capita per day availability of pulses in 2017 was 52.9 g per day. The total area under *kharif* and summer pulses in Gujarat is 9.08 lakh ha and production of 9.43 MT with average productivity of 1037 kg/ha in 2018-19. Gujarat has 5.2 Lakh hectares area under cowpea cultivation and generates 3.5 lakh tons with the productivity of 665 kg per hectare (**DAG, 2019**).

On comparison to other common grain legumes, this crop is more resistant to of low fertility because of its significant of nitrogen fixation, efficient symbiosis with *mycorrhizae*, and capacity to grow on soils with a wider range of pH.Higher or lower yields, elevated or reduced development of the crop, and changes in the crop's physiology are all caused by

variations in the availability of nutrients to the crop. Quality and seed yield decline when nitrogen level decreases.

The application of phosphorus increased the number of pods per plant, the number of seeds per pod, and the mean seed weight, all of which boosted cowpea yield (**Ayodele and Oso, 2014**). Legume benefits from potassium application in terms of growth, nodulation, nitrogen fixation, and yield (**Giller, 2000**).

Cowpea productivity in our country is quite poor, so it is necessary to establish appropriate agronomic strategies to increase cowpea output. The most crucial of these activities is the foliar application of nutrients, which fully utilizes the crop's genetic potential. Foliar application is thought to have the advantages of quick and effective nutrient consumption, elimination of leaching losses, fixing, and regulation of nutrient uptake by the plant.

MATERIALS AND METHODS

A field experiment was carried out at Agronomy Instructional Farm, Department of Agronomy, C. P. College of Agriculture, Sardarkrushinagar Dantiwada Agricultural University, S. K. nagar. **Located in the** North Gujarat Agro-climatic region at 24° 19' North latitude and 72° 19' East longitude with an elevation of 154.52 meter above the mean sea level. This area's climate is sub-tropical monsoon type and falls under semi-arid region. The soil of research field was loamy sand in texture low levels in organic carbon (0.27 %) available nitrogen is 158 kg per hectare, available phosphorus is 38.9 kg per hectare, and available potash is 295.2 kg per hectare, having Electrical Conductivity 0.14 (dSm⁻¹), pH value of 7.58.

Total eight treatment combinations were laid out in (RBD) randomized block design with four replications. Treatments details are T₁: Control, T₂:RDF (Recommended dose of fertilizer), T₃:75% of RDF + 1% spray of each urea and urea phosphate at 15 and 30 DAS, T₄:75% of RDF + 1% spray of NPK 19:19:19 at 15 and 30 DAS, T₅:75% of RDF + 2% spray of each urea and urea phosphate at 30 DAS, T₆:75% of RDF + 2% spray of NPK 19:19:19 at 30 DAS, T₇:75% of RDF + 2% spray of each urea and urea phosphate at 15 and 30 DAS, T₈:75% of RDF + 2% spray of NPK 19:19:19 at 15 and 30 DAS. Cowpea variety (GC 5) was sown at a spacing of 45 cm × 10 cm spacing on 7th July and the RDF was 20:40 kg N: P₂O₅/ha. The crop was harvested at physiological maturity on 1st Oct, 2020.

Representative samples were taken from the output of each net plot in order to estimate

the amounts of nitrogen, phosphorous, and potassium in the seed and stover. The materials were mechanically ground into powder and oven dried at 60°C for 24 hours before the nutrients were calculated using Micro Kjeldhal's method, Vanadomolybdo phosphoric yellow color method and Flame photometric method respectively (**Tandon, 1998**).

Results and discussion

Effect on yield

The increase in seed yield (kg/ha) under T₇ and T₈ over T₂ (RDF) was 19.92 and 13.70 per cent and over T₁ (control) was 61.59 and 53.29 per cent, respectively. **stover** yield due to T₇ over T₁ (control) and T₂ (Recommended dose of fertilizer) was to the extent of 64.39 and 31.79 per cent and T₈ over T₁ and T₂ was to the accordance of 29.10 and 16.66 per cent, respectively. **Bute et al. (2019)**

Whereas Treatment T₁ (control) produced significantly lowest seed yield and **stover** yield.

The improvement in the overall seed production was mostly brought about by the enhanced nitrogen supply and decreased nutrient losses. Spraying of either 2 per cent urea and urea phosphate or 2 per cent NPK (19:19:19) aided with fast nitrogen and phosphorus absorption when the need for nutrients is at its highest due to foliar spray at 15 DAS and 30 DAS of the crop. As a result, it decreased flower drop, improved pod setting and increased seed production. The outcomes support the conclusions made by **Choudhary and Yadav (2011)** in cowpea who indicated that foliar spray of 2% DAP recorded significantly higher growth, yield parameters and yield at branching and flowering stages over other treatments **Pradeepa et al. (2017)** observed similarly result in cowpea crop.

Effect on quality parameter

The set of data concerning to the protein content in seeds of cowpea which are provided in Table 1. illustrated that the various treatments succeed to produced their significant effect on protein content. At 15 and 30 DAS, 75% RDF+2% spray of each urea and urea phosphate treatment (T₇) resulted in significantly greater protein content (22.93%), However, it was statistically equivalent to all other treatments examined except T₆ (75% of RDF + 2% spray of NPK 19:19:19 at 30 DAS) and T₁(Control). Increased protein content in seed could be attributed to the fact that higher nitrogen content in seed is directly related to increased nitrogen availability to plants. Because nitrogen is a vital component of amino acids, which constitute the basis of protein content, higher nitrogen levels in seeds were directly responsible for higher protein levels. **Gupta et al. (2014)** found similar results.

Effect on nutrients content

Summarized in Table 1. it appeared that different treatments exerted their significant influence on nitrogen content (%) in seed and stover. Treatment T₇ is recorded significantly higher N content but Except for T₆ and T₁, all other treatments remained on par in terms of nitrogen content (%) in seed. In case of nitrogen content in stover treatment T₇ observed significantly higher N content which was at par with T₈ and T₃.

A close examination of data indicated that significantly higher phosphorus content in seed (0.446%) and stover (0.273%) were recorded with treatment T₇, but it was at par with Treatments T₈, T₂, T₃, T₄ and T₅ in case of phosphorus content in seed and with treatments T₈, T₂ and T₃ in case of phosphorus content in **stover**. Treatment T₁ (control) recorded lower content of phosphorus in seed (0.399%) and stover (0.189%).

Potassium content in cowpea seed **and stover** was not found significant with respect to various **treatments** of fertilizer and foliar spray sources.

This increase in nutritional content in seed and stover might be due to favorable and synergistic effect on availability of nutrients at different rate of fertilizer at basal and foliar sprays. Similar observation was reported by **Vikrant *et al.* (2005), Pandya and Bhatt (2007), Venkatesh and Basu (2011) and Vighnesh (2021).**

Effect on nutrients uptake

A close examination of data (Table 2.) revealed that various treatment manifests their significant impacts on the N, P and K uptake by cowpea plant. Among all treatments, T₇ (75% of RDF + 2% spray of each urea and urea phosphate at 15 and 30 DAS) observed significantly higher nitrogen (N) uptake (71 kg/ha), Phosphorus (P) uptake (12 kg/ha) and Potassium (K) uptake (56 kg/ha) by the crop which was statistically equivalent to the application of Treatment T₈ (75% of RDF + 2% spray of NPK 19:19:19 at 15 and 30 DAS). Though, treatment T₁ (control) noted significantly lower nitrogen uptake (38 kg/ha), Phosphorus uptake (7 kg/ha) and Potassium uptake (32 kg/ha) by the cowpea crop respectively.

The enhanced nutrient intake may be attributed to increased nutrient availability to the plant and higher biomass production and retarded loss of chlorophyll and leaf nitrogen with boost photosynthesis and increased N, P, and K provide during flowering and pod filling stages, results collaborated with the findings of **Choudhary and Yadav (2011), Venkatesh and Basu (2011), and Vighnesh (2021).**

Data substantiated that different treatments did not significantly improve the soil fertility in terms of available N, P₂O₅ and K₂O (kg/ha) in soil after harvest of cowpea.

Conclusion

In light of results obtained from present investigation, as a result, it is recommended that the kharif cowpea crop be fertilized with 75% of RDF (15:30 kg N: P/ha) and 2% of either urea and urea phosphate or NPK 19:19:19 at 15 and 30 DAS to achieve good quality and greater nutrient content and uptake in loamy sand.

References

- Ayodele, O. J. and Oso, A. A. (2014). Cowpea responses to phosphorus fertilizer application at Ado-Ekiti, South-West Nigeria. *Journal of Applied Science and Agriculture*. **9**(2): 485-489.
- Bute, V.A; Thaokar, A; Gawli, K.A.; Sarda, A.; Nagmote, A. and Jatav S.K. (2019). Growth and yield of cowpea influenced by foliar application of nutrient. *J. pharmacogn. Phytochem.* **8**(5): 2034-2037.
- Choudhary, G.L and L .R. Yadav (2011). Effect of fertility levels and foliar nutrition on cowpea productivity. *Journal of food Legumes*. **24**(1): 67-68.
- DAG (2019). Annual Report, Directorate of Agriculture, Govt. of Gujarat, Gandhinagar.
- DES (2018). Directorate of Economics and Statistics. Department of Agriculture and Co-operative New Delhi.
- Giller, K. E. (2000). Nitrogen fixation in tropical cropping systems. *Agricultural Science*. **2**(1): 1-8.
- Gupta, B. R., Pathak, R. K., Bhan, S. and Singh, A. (1972). Effect of NPK on yield, nutrient and quality of toria (*Brassica compenstris* var. toria). *Indian Journal of Agronomy*. **17**(2): 88-91.
- Gupta, K.C. and Saxena, R. (2014). Resource management in cowpea (*Vigna unguiculata* L. Walp.) for yield maximization under rainfed conditions. *Journal Crop and Weed*, **11**:146-148.
- Gupta, S.C., Sangeev K. And Khandwe. (2011). Effect of biofertilizer and foliar spray of urea on symbiotic traits, nitrogen uptake and productivity of chickpea. *Journal of Food legumes*. **24**(2): 155-157.
- Kumar, C.V., Vaiyapuri, K., Amanullah, M.M. and Gopala swamy, G. (2013). Influence of foliar spray of nutrients on yield and economics of soybean (*Glycine max* L. Merrill). *Journal of Biological Sciences*. **13**: 563-565.

- Pandya, C.B. and Bhatt, V.R. (2007). Effect of different nutrient levels on yield and nutrient content of fodder cowpea. *Legume Research*. **30** (3): 218-220.
- Pradeepa, T.M. and Ganajaximath (2017). Effect of mode of fertilization on growth, yield and economics of cowpea genotypes. *Journal of Farm Science*. **30**(1): 129-131.
- Sebetha, E. T.; Ayodele, V. I.; Kutu, F. R. and Mariga, (2010). Yields and protein content of two cowpea varieties grown under different production practices in Limpopo province, South Africa. *African Journal of Biotechnology*. **5**: 628–634.
- Singhal, Vivekkumar.; Patel, G.G.; Patel, D.H. Uttamkumar and Saini, L. (2015). Effect of foliar application of water-soluble fertilizers on growth, yield and economics of vegetable cowpea production. *An International Quarterly Journal of Environmental Sciences Special, Issue*. **VII**: 79-83.
- Tandon, H. L. S. (1998). Methods of analysis of soils, plant, water and fertilizers. Fertilizer development and consultation organization. **31**:9-16.
- Venkatesh, M.S. and Basu, P.S., (2011). Effect of foliar application of urea on growth, yield and quality of chickpea under rainfed conditions. *Journal of Food Legumes*. **24**(2): 110-112.
- Vighnesh; C, Seenappa; S, Harisha and KN, Kalyanamurthy (2021). Effect of foliar nutrition on yield and nutrient uptake of cowpea (*Vigna unguiculata*). *Pharma innov*.**10**(9): 361-364.
- Vikrant, V.R.; Singh, H.; Malik, C.V.S. and Singh, B.P. (2005). Influence of farm yard manure and phosphorus application on grain yield and economic of cowpea. *Crop Research*. **29**(1): 87-90.

Table 1. Effect of inorganic fertilizers and foliar spray on yield, quality and content of *kharif* cowpea

Treatments	Yield (kg/ha)		Protein content (%)	Nitrogen Content (%)		Phosphorus content (%)		Potassium content (%)	
	Seed	Stover		Seed	Stover	Seed	Stover	Seed	Stover
T ₁	880	1553	20.30	3.25	0.62	0.399	0.189	1.30	1.33
T ₂	1186	1937	21.98	3.52	0.69	0.434	0.259	1.35	1.38
T ₃	1214	2083	22.05	3.53	0.71	0.431	0.258	1.37	1.39
T ₄	1194	2002	22.03	3.53	0.69	0.420	0.236	1.41	1.39
T ₅	1098	1892	21.94	3.51	0.66	0.420	0.222	1.35	1.36
T ₆	1079	1937	20.93	3.33	0.64	0.409	0.227	1.39	1.35
T ₇	1422	2553	22.93	3.67	0.73	0.446	0.273	1.35	1.44
T ₈	1349	2260	22.75	3.64	0.71	0.439	0.255	1.45	1.40
S.Em. ±	59	106	0.52	0.08	0.01	0.010	0.008	0.03	0.04
C.D. at 5 %	173	313	1.52	0.24	0.03	0.029	0.025	NS	NS
C.V. %	10.01	10.50	4.74	4.74	3.16	4.69	6.95	4.27	6.01

Table: 2 Effect of inorganic fertilizers and foliar spray on total nutrient uptake and available nutrients after harvest in soil

Treatments	Nitrogen uptake (kg/ha)	Phosphorus uptake (kg/ha)	Potassium uptake (kg/ha)	Available Nutrient (kg/ha)		
				N	P ₂ O ₅	K ₂ O
T ₁	38	7	32	156	46	258
T ₂	55	10	42	165	49	263
T ₃	58	11	46	172	48	270
T ₄	56	10	44	160	50	269
T ₅	51	9	40	176	51	263
T ₆	48	9	41	176	48	268
T ₇	71	13	56	171	52	262
T ₈	65	12	51	169	51	274
S.Em. ±	3.0	0.5	2.0	4.8	1.3	6.3
C.D. at 5 %	9.0	1.5	6.7	NS	NS	NS
C.V. %	11.1	10.38	10.38	5.77	5.08	4.72