

Integrated Nutrient Management for Sustainable Sprouting Broccoli (*Brassica Oleracea L. Var. italica*) Production

ABSTRACT

Aim: The objective of the research was to study the effect of Integrated Nutrient Management on the vegetative growth of sprouting broccoli (*Brassica Oleracea L. Var. italica*) Var. Green Magic.

Study design: The field experiment was conducted in Randomised Block Design (RBD).

Place and duration of study: The experiment was conducted at the Research Farm of the Department of Vegetable Science, Chandra Shekhar Azad University of Agriculture & Technology, Kanpur U.P during the *Rabi* seasons of 2021-22 and 2022-23.

Methodology: During bed preparation, the required quantity of farmyard manure (FYM), vermicompost and biofertilizers was mixed into the soil in individual specified plots according to the treatment. The treatments included **T₀**: Control, **T₁**: 100% RDF @ 120: 80: 60, N: P: K (kg/ha), **T₂**: 75% RDF + 25% N through FYM, **T₃**: 75% RDF + 25% N through Vermicompost, **T₄**: 75% RDF + 25% N through FYM + Biofertilizer, **T₅**: 75% RDF + 25% N through Vermicompost + Biofertilizer, **T₆**: 75% RDF + 12.5% N through FYM + 12.5% N through Vermicompost, **T₇**: 75% RDF + 12.5% N through FYM + 12.5% N through Vermicompost + Biofertilizer, **T₈**: 50% RDF + 50% N through FYM, **T₉**: 50% RDF + 50% N through Vermicompost, **T₁₀**: 50% RDF + 50% N through FYM + Biofertilizer, **T₁₁**: 50% RDF + 50% N through Vermicompost + Biofertilizer, **T₁₂**: 50% RDF + 25% N through FYM + 25% N through Vermicompost and **T₁₃**: 50% RDF + 25% N through FYM + 25% N through Vermicompost + Biofertilizer. Observations were recorded on plant height (cm) and plant spread (cm) at 30 DAT, 60 DAT and at maturity and on stalk length (cm) and stalk diameter (cm) at maturity.

Results: The treatment **T₁₁**: 50% RDF + 50% N through Vermicompost + Biofertilizer outperformed all the other treatments in terms of plant height (cm), plant spread (cm), stalk length (cm) and stalk diameter (cm).

Conclusion: The study emphasizes the vital necessity of transitioning to INM with organic and inorganic fertilizers to preserve soil fertility, ecosystem health, and sustainable food production for future generations.

Keywords: Biofertilizer, *Brassica oleracea L. Var. italica*, FYM, INM, vermicompost,

1. INTRODUCTION

Broccoli, scientifically known as *Brassica oleracea* var. *italica*, holds a significant place among cruciferous vegetables. In India, broccoli cultivation is prominent in Himachal Pradesh, the hilly areas of Uttar Pradesh as well as Jammu and Kashmir and the northern plains. Broccoli is a nutritional powerhouse, boasting high levels of essential vitamins. It is notably rich in vitamin C, known for its immuno-boosting properties and antioxidant benefits. Furthermore, broccoli serves as an excellent source of vitamin A, essential for vision and skin health, vitamin B2 (riboflavin), which supports various metabolic processes in the body, and calcium, crucial for strong bones and teeth [1]. These

39 key factors attribute to the surge in popularity of broccoli among the affluent health-conscious
40 individuals of the country.

41 In today's market, the growing demand for vegetables has pushed farmers to produce higher yields
42 thus intensifying competition in the agricultural sector. To achieve this, farmers have turned to the
43 application of heavy doses of chemical fertilizers to boost growth and yield. This overreliance on
44 chemical fertilizers, in an attempt to increase food and fiber production has paradoxically resulted in
45 diminishing productivity over time, despite the apparent nutrient abundance. Hence, there arises a
46 pressing need for farmers to adopt a more balanced approach to plant nutrition management. This
47 involves integrating organic and inorganic fertilizers in a judicious manner, a practice known as
48 integrated nutrient management (INM). **Integrated nutrient management not only enhances and
49 maintains soil fertility but also benefits the physical, chemical, and biological properties of the soil [2].**
50 Moreover, it helps prevent deficiencies in secondary and micronutrients, ensuring a more sustainable
51 agricultural system. Accordingly, the present study was undertaken to know the effect on Integrated
52 Nutrient Management on vegetative growth of sprouting broccoli.

53 **2.METHODOLOGY**

54 The experiment was conducted during *Rabi* seasons of 2021-22 and 2022-23 at the Research Farm
55 of the Department of Vegetable Science, Chandra Shekhar Azad University of Agriculture &
56 Technology, Kanpur U.P. **Geographically, Kanpur is situated in the alluvial belt of Gangetic Plains of
57 Central Zone of U.P. It is positioned at 25.26" and 26.58" North latitude and 79.31" and 80.34" East
58 longitude at an altitude of 127 meter above the mean sea level (MSL).** This investigation was aimed at
59 knowing the effect of Integrated Nutrient Management on the vegetative growth of sprouting broccoli
60 Var. Green Magic. The experiment was laid out in a Randomized Block Design (RBD) with fourteen
61 treatments in three replications. The treatments included T_0 : Control, T_1 : 100% RDF @ 120: 80: 60,
62 N: P: K (kg/ha), T_2 : 75% RDF + 25% N through FYM, T_3 : 75% RDF + 25% N through Vermicompost,
63 T_4 : 75% RDF + 25% N through FYM + Biofertilizer, T_5 : 75% RDF + 25% N through Vermicompost +
64 Biofertilizer, T_6 : 75% RDF + 12.5% N through FYM + 12.5% N through Vermicompost, T_7 : 75% RDF
65 + 12.5% N through FYM + 12.5% N through Vermicompost + Biofertilizer, T_8 : 50% RDF + 50% N
66 through FYM, T_9 : 50% RDF + 50% N through Vermicompost, T_{10} : 50% RDF + 50% N through FYM +
67 Biofertilizer, T_{11} : 50% RDF + 50% N through Vermicompost + Biofertilizer, T_{12} : 50% RDF + 25% N
68 through FYM + 25% N through Vermicompost and T_{13} : 50% RDF + 25% N through FYM + 25% N
69 through Vermicompost + Biofertilizer. Initially, seedlings were raised under a polyhouse in Pro-trays,
70 and they were ready for transplanting to the main field in approximately 28-30 days. Meanwhile,
71 during bed preparation, the required quantity of farmyard manure (FYM) and vermicompost was
72 broadcasted and mixed into the soil in individual specified plots according to the treatment
73 combinations to ensure proper mineralization before seedling transplanting. Nonetheless, biofertilizers
74 like PSB and *Azotobacter* were mixed into the soil a mere 2-3 days prior to transplanting. Nitrogen,
75 phosphorus, and potash were applied in the form of urea (46% N), di-ammonium phosphate (46%
76 P_2O_5), and muriate of potash (60% K_2O), respectively. As per treatment combinations, the entire
77 recommended dose of phosphorus, potassium, and one-third of the nitrogen was applied during

78 transplanting. The remaining half of the nitrogen was split into two doses, one 30 days after
 79 transplanting and the other just before head initiation. The remaining half dose of nitrogen was split
 80 into two applications, with the first occurring at 30 days after transplanting and the second just before
 81 head initiation. Four weeks old healthy and uniform seedlings having 4-5 leaves with an average
 82 height of about 8-10 cm were transplanted in well prepared experimental plots in straight lines during
 83 evening hours at a spacing of 60 cm × 45 cm. Subsequently, the recommended package of practices
 84 for cultivation was diligently followed. Observations were recorded on five randomly selected plants
 85 per replication for plant height (cm) and plant spread (cm) at 30 DAT, 60 DAT and at maturity and on
 86 stalk length (cm) and stalk diameter (cm) at maturity. The field data recorded for agronomic traits were
 87 statistically analyzed using randomized block design (RBD), as per [3].

88 3. RESULTS AND DISCUSSION

89 The data recorded on vegetative parameters clearly indicated that the plant height (cm), plant spread
 90 (cm), Stalk length (cm) and stalk diameter (cm) were significantly influenced by the integrated nutrient
 91 management practices.

92 Plant Height (cm)

93 The data illustrating the impact of integrated nutrient management on plant height (cm) is presented in
 94 Table 1 and Figure 1. At 30 days after transplantation (DAT), the highest plant heights of 27.13 cm
 95 and 28.90 cm were observed in T₁₁: 50% RDF + 50% N through Vermicompost + Biofertilizer. In
 96 contrast, the lowest plant heights of 19.04 cm and 19.92 cm were recorded in T₀: Control.
 97 Furthermore at 60 DAT, the tallest plants, measuring 49.49 cm and 50.58 cm, were found in T₁₁: 50%
 98 RDF + 50% N through Vermicompost + Biofertilizer. Conversely, the shortest plants at this stage were
 99 in T₀: Control, with heights of 42.10 cm and 44.38 cm. Upon reaching the harvesting stage, the
 100 maximum plant heights were observed in T₁₁: 50% RDF + 50% N through Vermicompost +
 101 Biofertilizer, reaching 51.03 cm and 52.31 cm. Meanwhile, T₀: Control exhibited the minimum plant
 102 heights at this stage, measuring 46.63 cm and 47.66 cm. The observed effect can be attributed to the
 103 combined benefits of Vermicompost and biofertilizers. Vermicompost supplies essential
 104 micronutrients like zinc, calcium, copper, and iron at optimal levels, aids in nutrient retention from
 105 inorganic fertilizers, enhances soil properties, and boosts microbial activity, making both micro and
 106 macro nutrients readily available. Biofertilizers, on the other hand, serve as microbial inoculants,
 107 facilitating the decomposition of organic matter. This synergistic combination of 50 % RDF,
 108 Vermicompost and biofertilizers collectively contribute to improved plant growth. This finding is in
 109 confirmity with the results of [4] [5] [6] [7] [8].


110 **Table 1. The effect of Integrated Nutrient Management of Plant height (cm) of sprouting broccoli**

Treatment Details		Plant height (cm)					
		At 30 DAT		At 60 DAT		At Maturity	
		2021-22	2022-23	2021-22	2022-23	2021-22	2022-23
T ₀	Control	19.04	19.92	42.10	44.38	46.63	47.66
T ₁	100% RDF @ 120: 80: 60, N: P: K (kg/ha)	23.01	23.72	45.70	46.59	48.85	49.10

T ₂	75% RDF + 25% N through FYM	23.43	24.01	45.18	45.60	46.01	46.62
T ₃	75% RDF + 25% N through Vermicompost	24.14	25.01	44.97	45.73	48.05	49.57
T ₄	75% RDF + 25% N through FYM + Biofertilizer	23.78	24.54	47.11	47.98	48.99	49.30
T ₅	75% RDF + 25% N through Vermicompost + Biofertilizer	25.21	25.54	48.06	48.74	48.86	49.52
T ₆	75% RDF + 12.5% N through FYM + 12.5% N through Vermicompost	22.83	23.71	46.69	47.62	48.21	48.83
T ₇	75% RDF + 12.5% N through FYM + 12.5% N through Vermicompost + Biofertilizer	21.74	22.75	46.74	47.73	48.67	48.56
T ₈	50% RDF + 50% N through FYM	22.75	25.25	47.09	48.44	50.20	51.14
T ₉	50% RDF + 50% N through Vermicompost	23.15	25.85	46.39	47.78	49.79	49.68
T ₁₀	50% RDF + 50% N through FYM + Biofertilizer	24.25	26.11	48.23	48.85	48.68	49.61
T ₁₁	50% RDF + 50% N through Vermicompost + Biofertilizer	27.13	28.90	49.49	50.58	51.03	52.31
T ₁₂	50% RDF + 25% N through FYM + 25% N through Vermicompost	24.11	25.34	44.83	45.53	48.11	48.60
T ₁₃	50% RDF + 25% N through FYM + 25% N through Vermicompost + Biofertilizer	25.79	27.28	49.13	49.30	50.13	50.78
SE(m) ±		0.569	4.863	1.093	0.851	0.655	0.602
CD (P=0.05)		1.66	0.69	3.18	2.47	1.90	1.75

111

112 Figure 1. The effect of Integrated Nutrient Management of Plant height (cm) of sprouting broccoli



113

114 **Plant spread (cm)**

115 The data in Table 2 and Figure 2 showcases the effects of integrated nutrient management on plant
116 spread. At 30 DAT, the most extensive plant spread, measuring 37.13 cm and 37.53 cm, was
117 observed in T₁₁: 50% RDF + 50% N through Vermicompost + Biofertilizer. In contrast, the minimum


118 plant spread, 25.87 cm and 26.32 cm, was recorded in T₀: Control. Subsequently at 60 DAT, the
 119 maximum plant spreads of 58.67 cm and 59.91 cm were found again in T₁₁: 50% RDF + 50% N
 120 through Vermicompost + Biofertilizer. Conversely, the smallest plant spread at this stage was in T₀:
 121 Control, with measurements of 48.23 cm and 49.03 cm. Upon reaching the harvesting stage, the
 122 maximum plant spread was observed in T₁₁: 50% RDF + 50% N through Vermicompost + Biofertilizer,
 123 reaching 68.80 cm and 70.05 cm. Meanwhile, T₀: Control exhibited the minimum plant spread at this
 124 stage, measuring 53.19 cm and 55.22 cm. The plant spread and various growth parameters were
 125 notably amplified when biofertilizer and vermicompost were applied in combination, as demonstrated
 126 in studies by [9] [10] [11] [12] [13] [14]. Nitrogen is recognized as a pivotal nutrient for optimal plant
 127 growth and development. The improved plant spread achieved by applying nitrogen through
 128 vermicompost and biofertilizer can be attributed to the favourable soil conditions they foster, including
 129 enhanced moisture retention capabilities and increased availability of major nutrients.

130 **Table 2. The effect of Integrated Nutrient Management of Plant spread (cm) of sprouting broccoli**

Treatment Details		Plant spread (cm)					
		At 30 DAT		At 60 DAT		At Maturity	
		2021-22	2022-23	2021-22	2022-23	2021-22	2022-23
T ₀	Control	25.87	26.32	48.23	49.03	53.19	55.22
T ₁	100% RDF @ 120: 80: 60, N: P: K (kg/ha)	31.20	30.82	50.53	51.79	59.46	60.07
T ₂	75% RDF + 25% N through FYM	32.31	33.66	54.52	55.48	55.19	56.94
T ₃	75% RDF + 25% N through Vermicompost	28.89	29.99	48.37	49.34	58.46	59.11
T ₄	75% RDF + 25% N through FYM + Biofertilizer	34.34	34.14	49.65	51.52	67.23	67.30
T ₅	75% RDF + 25% N through Vermicompost + Biofertilizer	34.26	34.96	54.47	56.85	67.36	67.69
T ₆	75% RDF + 12.5% N through FYM + 12.5% N through Vermicompost	35.81	36.78	56.25	53.34	67.53	67.93
T ₇	75% RDF + 12.5% N through FYM + 12.5% N through Vermicompost + Biofertilizer	34.81	35.61	54.84	55.46	65.00	66.10
T ₈	50% RDF + 50% N through FYM	34.70	34.26	56.08	56.51	67.34	67.85
T ₉	50% RDF + 50% N through Vermicompost	33.92	33.42	54.50	56.60	66.53	67.20
T ₁₀	50% RDF + 50% N through FYM + Biofertilizer	34.02	34.20	54.73	55.39	67.17	68.00
T ₁₁	50% RDF + 50% N through Vermicompost + Biofertilizer	37.13	37.53	58.67	59.91	68.80	70.05
T ₁₂	50% RDF + 25% N through FYM + 25% N through Vermicompost	30.14	31.36	57.42	57.80	59.73	60.46
T ₁₃	50% RDF + 25% N through FYM + 25% N through Vermicompost + Biofertilizer	35.37	35.54	55.00	55.93	66.30	67.78
SE(m) ±		1.247	1.222	1.461	1.198	0.434	0.576
CD (P=0.05)		3.62	3.55	4.27	3.50	1.26	1.68

131
 132
 133

Figure 2. The effect of Integrated Nutrient Management of Plant spread (cm) of sprouting broccoli



134

135 Stalk length (cm)

136 The data depicting the influence of integrated nutrient management on stalk length (cm) have been
 137 displayed in Table 3 and Figure 3. At the final harvest the maximum stalk length (cm) of 20.81 cm and
 138 22.10 cm was recorded with the application of T₁₁ : 50% RDF + 50% N through Vermicompost +
 139 Biofertilizer. Whereas the minimum stalk length of 13.52 cm and 14.57 cm was recorded in T₀:
 140 Control. Similar results were reported by [15]. Accordingly, the best treatment increased broccoli stalk
 141 length by optimizing nutrient availability, improving moisture retention, enhancing soil conditions and
 142 promoting the activity of beneficial microbes in the soil. This treatment provides essential nutrients like
 143 nitrogen and micronutrients, supporting plant growth and cell elongation for longer stalks.


144 **Table 3. The effect of Integrated Nutrient Management of stalk length (cm) of sprouting broccoli**

Treatment Details		Stalk length (cm)	
		At 30 DAT	
		2021-22	2022-23
T ₀	Control	13.52	14.57
T ₁	100% RDF @ 120: 80: 60, N: P: K (kg/ha)	15.35	16.02
T ₂	75% RDF + 25% N through FYM	16.12	16.42
T ₃	75% RDF + 25% N through Vermicompost	17.25	18.22
T ₄	75% RDF + 25% N through FYM + Biofertilizer	19.12	20.15
T ₅	75% RDF + 25% N through Vermicompost + Biofertilizer	17.84	18.35
T ₆	75% RDF + 12.5% N through FYM + 12.5% N through Vermicompost	18.16	19.57
T ₇	75% RDF + 12.5% N through FYM + 12.5% N through Vermicompost + Biofertilizer	19.02	20.42
T ₈	50% RDF + 50% N through FYM	19.06	20.58
T ₉	50% RDF + 50% N through Vermicompost	20.19	20.41
T ₁₀	50% RDF + 50% N through FYM + Biofertilizer	19.28	20.41

T ₁₁	50% RDF + 50% N through Vermicompost + Biofertilizer	20.81	22.10
T ₁₂	50% RDF + 25% N through FYM + 25% N through Vermicompost	17.31	18.39
T ₁₃	50% RDF + 25% N through FYM + 25% N through Vermicompost + Biofertilizer	19.33	20.26
SE(m) ±		0.568	0.712
CD (P=0.05)		1.65	2.07

145
146

Figure 3. The effect of Integrated Nutrient Management of stalk length (cm) of sprouting broccoli



147
148

Stalk diameter (cm)

150 The data illustrating the impact of integrated nutrient management on stalk diameter (cm) is presented
 151 in Table 4 and Figure 4. During the harvesting stage, the plots treated with T₁₁ : 50% RDF + 50% N
 152 through Vermicompost + Biofertilizer exhibited the maximum stalk diameter of 4.03 cm and 4.62 cm.
 153 In contrast, T₀: Control reported minimum stalk diameter of 2.17 cm and 2.83 cm. results parallel to
 154 the present study were reported by [16]. Accordingly, the most effective treatment combination
 155 according to the present study, delivers ample essential nutrients, including nitrogen and
 156 micronutrients, in a well-balanced manner. This nutrient equilibrium fosters cell expansion and growth,
 157 consequently leading to thicker stalks.


158 Table 4. The effect of Integrated Nutrient Management of stalk diameter (cm) of sprouting broccoli

Stalk diameter (cm)			
Treatment Details		At 30 DAT	
		2021-22	2022-23
T ₀	Control	2.17	2.83
T ₁	100% RDF @ 120: 80: 60, N: P: K (kg/ha)	2.75	3.73

T ₂	75% RDF + 25% N through FYM	2.63	3.79
T ₃	75% RDF + 25% N through Vermicompost	2.75	3.75
T ₄	75% RDF + 25% N through FYM + Biofertilizer	2.84	3.87
T ₅	75% RDF + 25% N through Vermicompost + Biofertilizer	2.61	3.58
T ₆	75% RDF + 12.5% N through FYM + 12.5% N through Vermicompost	2.73	4.08
T ₇	75% RDF + 12.5% N through FYM + 12.5% N through Vermicompost + Biofertilizer	2.89	3.59
T ₈	50% RDF + 50% N through FYM	3.15	3.87
T ₉	50% RDF + 50% N through Vermicompost	2.85	3.70
T ₁₀	50% RDF + 50% N through FYM + Biofertilizer	2.94	3.75
T ₁₁	50% RDF + 50% N through Vermicompost + Biofertilizer	4.03	4.62
T ₁₂	50% RDF + 25% N through FYM + 25% N through Vermicompost	2.67	3.52
T ₁₃	50% RDF + 25% N through FYM + 25% N through Vermicompost + Biofertilizer	3.01	4.15
SE(m) ±		0.176	0.158
CD (P=0.05)		0.51	0.46

159

160 Figure 4. The effect of Integrated Nutrient Management of stalk diameter (cm) of sprouting
161 broccoli



162

163

164 4. CONCLUSION

165 In this study, it is evident that treatment T₁₁: 50% RDF + 50% N through Vermicompost + Biofertilizer
166 outperformed all the other treatments in terms of plant height (cm), plant spread (cm), stalk length
167 (cm) and stalk diameter (cm). Fundamentally, this study highlights the need to shift from chemical
168 fertilizers to INM with organic and inorganic fertilizers from the imperative to protect soil fertility,

169 maintain ecosystem health, and ensure sustainable food production for the future. This transition isn't
170 just a matter of choice, it's an essential mandate to safeguard the health of ecosystems and secure
171 the welfare of generations to come.

172 **ACKNOWLEDGEMENTS**

173 The authors extend their heartfelt gratitude to the department where this study was conducted for
174 their invaluable support and resources. Your assistance was instrumental in the successful
175 completion of this research.

176 **COMPETING INTRESTS**

177 The authors declare that they have no competing interests.

178 **AUTHOR CONTRIBUTIONS**

179 **Pranjal Singh:** Data collection, wrote the first draft of the manuscript.

180 **Sanjive Kumar Singh:** Designed the research program.

181 **Ram Batuk Singh:** Provided the facilities required during the course of the research.

182 **Ashutosh Upadhyay:** Analysis of the study.

183 **Bankey Lal:** Managed the literature searches.

184 All the authors read and approved the final manuscript.

185 **REFERENCES**

186 1. Sanwal SK, Laxminarayana K, Yadav DS, Rai N and Yadav RK. Growth, yield, and dietary
187 antioxidants of broccoli as affected by fertilizer type. Journal of vegetable
188 Science. 2006;12(2):13-26.

189 2. Aulakh MS and Grant CA. Integrated nutrient management for sustainable crop production.
190 2008. (No Title).

191 3. Panse VG and Sukhatme PV. Statistical methods for agricultural workers (4th edition). Indian
192 Council of Agricultural Research, New Delhi, India. 1985.

193 4. Kumar P, Kumar S, Meena RK, Kumar R and Rawat R. Efficacy of bio-fertilizers on growth,
194 yield and quality of sprouting broccoli (*Brassica oleracea* var. *italic plank*), cv. Pusa Broccoli
195 KTS-1. Plant Archives. 2017;17(2):1647-1650.

196 5. Mohanta R, Nandi AK, Mishra SP, Pattnaik A, Hossain MM and Padhiary AK. Effects of
197 integrated nutrient management on growth, yield, quality and economics of sprouting broccoli
198 (*Brassica oleracea* L. var. *italica*) cv. Shayali. Journal of Pharmacognosy and Phytochemistry.
199 2018;7(1): 2229- 2232.

- 200 6. Singh S, Chamroy T, Malik S and Kundu S. Integrated nutrient management (INM) studies on
201 growth and yield of broccoli (*Brassica oleracea* var. *italica* Plenck). *Plant Archives*. 2020;20(2):
202 9651-9654.
- 203 7. Walling I, Kanaujia SP and Changini M. Response of broccoli (*Brassica oleracea* var, *italica*) to
204 integrated nutrient management. *Annals of Plant and Soil Research*. 2022;24(1):106-109.
- 205 8. Sagar K, Kumar D, Singh N and Pathania A. Response of Integrated Nutrient Management on
206 Growth and Yield of Cauliflower (*Brassica oleracea* var. *botrytis*). *Environment and Ecology*.
207 2023;41(2), 772-780.
- 208 9. Manivanan MI and Singh JP. Effect of biofertilizer on the growth and yield of sprouting broccoli.
209 *Indian journal of applied science research*. 2004;15(2): 33-36.
- 210 10. Dass A, Lenka NK, Patnaik US and Sudhishri S. Integrated nutrient management for
211 production, economics, and soil improvement in winter vegetables. *Journal of Vegetable*
212 *Science*. 2009;14(2): 104-120.
- 213 11. Singh V, Khan NS and Rana DK. Combined effect of organic manures and bio-fertilizers on
214 growth and yield of broccoli under Garhwal Himalayan Region. *Hort Flora Research Spectrum*.
215 2017;5(4).
- 216 12. Gogoi P, Barua M, Barua PK, Gogoi S and Borah N. Nutrient management for quality seed
217 production of broccoli in Assam. *Indian Journal of Horticulture*. 2021;78(4):409-416.
- 218 13. Tiwari P, Mishra SK, Roy S and Kumar N. Study on integrated nutrient management in broccoli
219 (*Brassica oleracea* L. var. *italica* Plenck) for better growth and curd quality. *The Pharma*
220 *Innovation Journal*. 2021;10(3): 161-163.
- 221 14. Chaudhary RP, Kumar J, Thapa P, Singh A, Giri B, Kishore B. Effect of integrated nutrient
222 management on growth and yield parameter of cauliflower (*Brassica oleracea* var. *Botrytis*)
223 variety Madhuri. 2023
- 224 15. Sharma C, Kang BS, Kaur R, Singh SK and Aulakh K. Effect of integrated nutrient management
225 on growth, yield and quality of broccoli (*Brassica oleracea* L. var. *italica*). *International Journal*
226 *of Chemical Studies*. 2018;6(2): 1296-1300.
- 227 16. Atal MK, Dwivedi DH, Narolia SL, Bharty N and Kumari R. Influence of bio-fertilizer (*Rhizobium*
228 *radiobacter*) in association with organic manures on growth and yield of broccoli (*Brassica*
229 *oleracea* L. var. *italica* Plenck) cv. Palam Samridhi under Lucknow conditions. *Journal of*
230 *Pharmacognosy and Phytochemistry*. 2019;1: 604-608.

231

232

233

234

235

236

237

238

239