

Studies on Storability of Manual and Mechanically Harvested and Threshed Sunnhemp Seeds (*Crotalaria juncea* Linn.)

ABSTRACT

Background: The present investigation is an attempt to study the effect of different harvesting and threshing methods on storability of sunnhemp seed. The use of a combine harvester to harvest sunnhemp will be an effective alternative method. However, work on different methods of harvesting and threshing of sunnhemp is limited and storage studies of machine harvested seed is scanty. Hence, a study was conducted to evaluate the impact of various harvesting and threshing methods on storability of sunnhemp seeds.

Methods: This study was conducted at Agricultural Engineering College and Research Institute, TNAU, Kumulur, Tiruchirappalli, Tamil Nadu. The sunnhemp seed crop was harvested and threshed using four different methods viz., manual harvesting and manual threshing, manual harvesting and mechanical threshing, manual harvesting and threshing by tractor treading and harvesting and threshing by combine harvester. The resultant seeds were cleaned and graded. The graded seeds were stored under ambient conditions in both cloth bag (C1) and super grain bag (C2) containers with 12.0 per cent and 8 per cent moisture content respectively. The experiment was designed adopting FCRD with eight replications. The following quality parameters were recorded initially and at monthly intervals for a period of 12 months to assess the storability of seeds. Treatment wise the seed samples were taken and the seed moisture content was estimated by low constant temperature oven method. Germination was tested by roll towel method using 50 seeds in eight replications. Germination percentage, root and shoot length were measured in ten days after sowing from ten randomly selected seedlings in each replication. For the estimation of dry matter production, ten seedlings were selected at random and kept in a hot air oven maintained at 85°C for 24 hours after measuring their root and shoot length and vigour index was calculated. At the end of each every month from one to 12th month, samples were collected and the seeds were subjected to health test. Seed health tests were done by Blotter Incubation method.

Result: The result revealed that there was a significant difference found among the different harvesting and threshing methods on storability of sunnhemp seed. Irrespective of harvesting and threshing methods seeds stored in cloth bag recorded 12% moisture content for initial month and increased up to 13.76% in 12th month whereas seeds stored in super bag recorded 8% initial month and increased up to 8.78% in 12th month. In this study, seeds obtained by manual harvesting and manual threshing method registered maximum germination percentage (93% & 94%), followed by seeds obtained by combine harvesting (91% & 92%). The minimum germination percentage was recorded in the seeds obtained by manual harvesting and mechanical threshing (89% & 90%). The maximum germination percentage was observed during the initial period of storage and reached the minimum at 12 months of storage. Between the containers, super grain bag maintained the highest germination percentage (92%) while the lowest germination percentage was observed in the seeds stored in cloth bag (91%). Seed health test revealed that the seed borne pathogens were identified after four months of storage period and no incidence was noticed up to four months of storage. From this study it could be concluded that sunnhemp seed crop harvested and threshed by different methods and reduced to the seed moisture content of 8 percent and packed in super grain bag maintained seed quality above minimum seed certification standards up to twelve months of storage. Hence it is recommended that combine can be used to harvest the sunnhemp seed crop to minimise the cost of labour as well as saving time and found to be maintaining the longevity of sunnhemp seed under ambient condition without deterioration of viability.

Key words: *Crotalaria juncea*, Containers, Germination, Harvesting and threshing methods, Seed borne pathogen, Seed storage.

INTRODUCTION

Farm mechanization is one of the realistic approaches to improve agriculture production and has the obvious advantages of reduction in human drudgery, reduction in cost of cultivation, enhancement of working efficiency and timeliness of working (Masilamani and Tajuddin 2012; Masilamani *et al.*, 2021). Different types of farm operations starting from land preparation to seed storage are involved in agriculture. All these operations are labour intensive and time consuming when carried out with manual labourers. Failure to complete farm operations within the stipulated time may sharply reduce the crop production. Seed quality is the most important factor influencing crop growth, development and yield processes and could increase yield by 5-20% (Rickman *et al.*, 2006). Harvesting sunnhemp seed crop using combine has been accepted and practiced to overcome the peak demand of farm labourers and to minimize the field losses incurred during manual harvesting. Sunn hemp is a tropical legume that has been used as a green manure for soil improvement and efficient nitrogen fixer (Masilamani and Sivasubramaniam, 2016; Rajendra Prasad *et al.*, 2017). It is a potential tropical cover crop, reduces erosion, and improves soil fertility and tilth (Alwell, 2015). It produces biomass yield of 15-20t/ha. (Lates and Mabbayad, 1983). It is resistant to nematodes (Mc sorely *et al.*, 1994) and it can grow on dry zone soil with low fertility. It suppresses weeds (Rotar and Joy, 1983) through smothering effect and conserves soil moisture by reducing evaporation from soil. It promotes biological transformation in soil leading to improved soil structure, fertility and increased crop yields (Pradhan *et al.*, 2001; Ulemale *et al.*, 2002). It is mostly raised for biomass and is normally incorporated in to the soil during pre-flowering phase. The crop is traditionally used for making ropes, strings, twines, floor mat, fishing nets, hand-made paper, *etc.* in cottage industry (Desai *et al.*, 2023). Hence there should be enough buffer stock of seed to meet the season-season seed requirement (Masilamani *et al.*, 2017). Efficient seed technological interventions encompassing seed production, processing and storage are essential. Seed deterioration during storage is a gradual and inevitable process causing considerable losses. Seeds tend to lose viability and vigour during storage and information on storability of seed lots from harvest until the next planting season and also for carry over purposes is of immense importance in any seed production programme (Johny subakarivinet *et al.*, 2021). In storage, viability and vigour of the seeds is regulated by many physico-chemical factors such as moisture content of the seed, atmospheric humidity, temperature and initial seed quality, physical and chemical composition of seed, gaseous exchange, storage structure and packaging materials. Till date there was no work so far carried out to find out the storability of machine harvested sunnhemp seeds. Against the stalemate the study was initiated to assess the storability of manual and mechanically harvested and threshed sunnhemp seeds.

MATERIALS AND METHODS

An experiment was conducted at Agricultural Engineering College and Research Institute, Tamil Nadu Agricultural University, Kumulur, Tiruchirappalli, Tamil Nadu during 2019-2020 to find out the

influence of harvesting and threshing methods on seed storability of sunnhemp seeds. The treatments are manual harvesting and manual threshing (T1), manual harvesting and mechanical threshing (axial flow thresher) (T2), manual harvesting and tractor treading (T3) and combine harvesting (with pneumatic wheel) (T4)(Fig. 1).. The seeds collected from different harvesting and threshing methods were cleaned and graded. The graded seeds were stored in both cloth bag (C1) and super grain (C2) containers with 12 per cent and 8 per cent moisture content respectively and stored under ambient conditions. The experiment was designed adopting FCRD with eight replications. The following quality parameters were recorded initially and at monthly intervals for a period of 12 months to assess the storability of seeds.

Seed moisture content (%)

Treatment wise the seed samples were taken and the seed moisture content was estimated by low constant temperature oven method at 103 ± 1 °C for 16 ± 1 h with known weight of seed samples. After drying, the seed samples were placed in desiccators containing calcium chloride for 30 min and weighed. The per cent of moisture content was calculated using the following formula (ISTA 1985).

$$\text{Moisture content (\%)} = \frac{M_2 - M_3}{M_2 - M_1} \times 100$$

Where,

M_1 = Weight of empty moisture bottle along with lid (g)

M_2 = Weight of moisture bottle along with sample before drying (g)

M_3 = Weight of moisture bottle along with sample after drying (g)

Germination Test

Treatment wise, the seeds were placed for germination in roll towel method. Under each treatment, 400 seeds were sown with eight replications of 50 seeds each. Seed germination was expressed as the percentage of seeds producing normal seedlings (Central Seed Certification Board. 2013). Ten days after sowing ten seedlings from each replication were randomly selected and the root and shoot lengths were measured and the mean value was recorded. Ten random seedlings were dried in a hot air oven at 85^0 C for 24 h. and the dry weight was recorded and expressed as g.seedling⁻¹⁰. The vigour index I and II was calculated using the following formula (Abdul – Baki and Anderson, 1973) by following the below given formula.

$$\text{Vigour index - I} = \text{Germination (\%)} \times \text{Total seedling length (cm)}$$

$$\text{Vigour index - II} = \text{Germination (\%)} \times \text{Dry matter production (g/10 seedlings)}$$

Seed Health Test

At the end of the each every month from one to 12th month, samples were collected from cloth and super grain bag and the seeds were subjected to health test. Seed health tests were done by Blotter Incubation method following ISTA (1999) procedure. Blotter incubation test was done on Whatman No. 1 blotter paper contained in petridish. Three layer of water soaked blotter papers were placed on each petridish and 25 seeds/plate were placed. The seeds were incubated in the incubation chamber. After five days, number of seeds with mycelia colonies was counted.

Statistical analysis

The data obtained from the experiment were analyzed by the 'F' test of significance following the methods described by Panse and Sukhatme, 1995. Wherever necessary, the per cent values were transformed to angular (Arc-sine) values before analysis. The critical differences (CD) were calculated at 5 per cent probability level. The data were tested for statistical significance.



Manual Harvesting



Manual Threshing



Mechanical Threshing



Tractor Treading



Combine Harvesting

Fig.1. Different methods of Harvesting and threshing of Sunnhemp

RESULTS AND DISCUSSION

Moisture Content

The results

of the different harvesting and threshing methods on storability of sunn hemp revealed that the moisture content was significantly influenced by harvesting and threshing methods, container used for storage and period of storage in sunn hemp seed tested (Fig. 2). In this study, there was a slight increase in the seed moisture content over the storage period, irrespective of container and harvesting and threshing methods. Irrespective of harvesting and threshing methods seeds stored in cloth bag recorded 12% moisture content for initial month and increased up to 13.76% in 12th month whereas seeds stored in super bag recorded 8% initial month and increased up to 8.78% in 12th month. Increase in the moisture content is associated with decline of seed quality. These results are in agreement with the results obtained by Govindarajet al., 2017 in rice. Higher the moisture content and temperature, lesser the shelf life of paddy seed reported by Kaliyanet al., 2006. Ramanadane and Ponnusamy (2004) reported that the moisture content is associated with decline of seed quality. Harringtons thumb rule says that storability increases as the moisture content decreases. For every one percent reduction in moisture content, the shelf life is doubled within a range of 5 to 15%. It was reported that, there was a negative logarithmic relation between moisture content and longevity (Ellis et al., 1990).

The super grain bag played significant role in preventing vapour entry from the surrounding air. This becomes effective strategy in regulating lower moisture content in the seeds using vapour impervious containers. Lower respiration rate and metabolic activity are governed by lower moisture content and temperature during the storage period (Muangkaeo et al., 2005). Doijode (1995) reported that the seeds packed in polythene bags exhibited higher germination; seedling length and seedling dry weight. Similar observations have been reported by Padma and Reddy (2002) in maize and green gram. Seeds packed in polythene bag and acted as vapour proof barrier in regulating lower moisture content in the seeds.. This is in accordance with the findings of Azad et al., (2014) who have observed higher vigour when wheat seeds dried to 12 per cent moisture content and preserved in polythene bags stored for nine months. Similar findings have been reported by Saxena et al., (1987) in cereals.

Germination Test

Physiological parameters were significantly influenced by harvesting and threshing methods, container used for storage and period of storage in sunn hemp seed tested. In this study, seeds obtained by manual harvesting and manual threshing method registered maximum germination percentage (93% & 94%), followed by seeds obtained by combine harvesting (91% &

92%)(Table1).Theminimumgermination percentagewasrecorded inthe seedsobtained bymanualharvestingandmechanicalthreshing(89% & 90%).Themaximum germination percentagewasobservedduringtheinitialperiodof storage and reached the minimum at 12 monthsofstorage.Betweenthecontainers,supergrainbag maintained the highest germinationpercentage (92%)while the lowestgermination percentagewas observed in the seeds stored in cloth bag (91%). The root length and shoot length of the seedling reflected the same trend as on germination percentage. Regarding dry matter production, the seedsobtainedbymanualharvestingandmanualthreshing method registered maximum drymatterproduction(0.192g & 0.231g), seedling vigour I (3024 & 3585) and seedling vigour II (18 & 22)followed by seedsobtained by manual harvesting and tractor treading (2837 & 3317) (16 &19)(Table2,3,4,5and 6).

Theminimumdrymatterproduction, seedling vigour I and seedling vigour IIwasrecorded inthe seedsobtained bymanualharvestingandmechanicalthreshing(0.164g & 0.197g). The maximum drymatterproduction and seedling vigour I and vigour index IIwasobservedduringtheinitialperiodof storage and reached the minimum at 12 monthsofstorage.Betweenthecontainers,supergrainbag maintained the highest drymatterproduction, seedling vigour I and seedling vigour II while the lowestdrymatterproduction, seedling vigour I and II was observed in the seeds stored inclothbag. Between the harvesting method, manual harvesting and threshing maintained the highest drymatterproduction, seedling vigour I and II was observed and the lowest drymatterproduction, seedling vigour I and II was observed in manual harvesting and mechanical threshing. These results are in agreement with the results obtained by Akter *et al.* (2014) in soybean, Sharon *et al.*, 2015 in blackgram.

The decline ingerminationpercentage maybe attributedtoageingeffect. Ageinghasdamagingeffectonenzymesthat are necessary to convert reserve food in theembryo to usable form and ultimately production of normal seedling (Iqbal *et al.*, 2002). Alternatively, the reduction in germination, dry matter productionandseedlingvigourmightbeduetodegradationof mitochondrial membrane leading to reduction inenergy supply necessary for germination (Gidroletal., 1998). Loss of germination and seedling vigour during storage were reported by Ajayet *al.*, (2017) in soybean and Htweb *et al.*,(2018) ingreen gram and chickpea. The decline in the seedling dry matter production might be attributed to DNA degradation with ageing which leads to impaired transcription causing incomplete or faulty enzyme synthesis essential for earlier stages of germination (Kapoor *et al.*, 2002). All seeds undergo ageing process during long-term storage which leads to deterioration in seed quality, however, the rate of seed deterioration can vary among various plant species (Merritt *et al.*, 2003).Rajasekaran *et al.* (2005) observed that niger seeds packed in polylined cloth bags maintained higher germination and vigour index even after six months of storage. Van Chin and Kieu (2006) reported in rice seeds that germination percentage under super bag and vietnamese bag are similar at three and six months after storage. However, at 9 and 12

months storage, IRRI Super bag was superior to Vietnamese bag statistically. Singh and Dadlani (2003) stated that soybean seeds packed in 700 gauge poly ethylene bag could be stored for fourteen months whereas, the seeds packed in cloth bag could be stored only up to eight months.

Seed Health Test

The seed health studies revealed that the genera of fungi were identified under the compound microscope at 40x. Fungal species identified were *Aspergillus flavus* and *Fusarium* sp. In all the treatments, fungal incidence were noticed in seeds collected from cloth bag where as no incidence was found in seeds samples collected from super grain bag. Seed borne pathogens were identified after four months of storage period and no incidence was noticed up to four months of storage. Incidence of *Fusarium* sp. ranged from 1.5 to 3% at fifth month and from 14 to 16.5% at 12 months of storage. Incidence of *Aspergillus flavus*, ranged from 0 to 0.5% at four months of storage and from 10 to 10.5% at 12 months of storage (Fig.3). Similar type of seed mycoflora association was also reported by Sadhu (2014) and Devamani *et al.*, (2017) in green gram and Biswal *et al.*, 2019 in black gram

As seed deterioration is unavoidable and irreversible process it cannot be stopped completely but the extent of deterioration can be slowed down to certain extent. Similarly in our study, irrespective of the harvesting and threshing methods, the seed quality parameters declined progressively with an increase in storage period. However, seed harvested and threshed by manual method- manual harvesting and manual threshing, manual harvesting and mechanical threshing, manual harvesting and tractor treading and combine harvesting and stored in both cloth bag and super grain containers all are found to be maintaining the longevity of sunnhemp seed under ambient conditions. However, fungal incidence were noticed in seeds collected from cloth bag where as no incidence was found in seeds samples collected from super grain bag.

CONCLUSION

From this study it could be concluded that sunnhemp seed crop harvested and threshed by different methods and reduced to the moisture content of 8 percent and packed in super grain bag maintained seed quality above minimum seed certification standards up to twelve months of storage without any pathogen incidence. Hence it is recommended that combine harvester can be used to harvest the sunnhemp seed crop to minimise the cost of labour as well as saving time and found to be maintaining the longevity of sunnhemp seed under ambient storage.

REFERENCES

Abdul Baki AA and JD Anderson (1973). Vigour determination in Soybean seed by multiple criteria. *Crop Science*, 13: 630 -633.

- Alwell, A (2015). Sunn hemp gains popularity as stress-tolerant cover crop. *Organic Broadcaster*, 23:1&6
- Ajay P.G., K.Vishunavat, C.Mohan, S.Ravi.2017.Effect of seed coating, storage periods and storage containers on soybean (*Glycine max* (L.) Merrill) seed quality under ambient conditions.*Journal of Applied and Natural Science*. Vol. 9(1). DOI <https://doi.org/10.31018/jans.v9i1.1237>.
- Akter, N., M. M. Haque, M. R. Islam and K. M. Alam. 2014.Seed Quality of Stored Soybean (*Glycine max* L.) as Influenced by Storage Containers and Storage Periods. *The Agriculturists*, 12(1): 85-95
- Azad A. W., T. Jaya Joshi, T. Anurag and D. S. Tomar. 2014. Effect of seed treatments and packing materials on seed quality parameters of maize (*Zea mays* L.) during Storage. *Indian J. Appl. Res.*, 4:40-44.
- Biswal, K., N. Ranasingh, K. C. Sahu, R. L. Moharana and Behera, S. 2019. Seed Health Status of Farmers' Saved Black Gram (*Vigna mungo* (L.) Hepper) Seeds in Western Undulated Zones of Odisha. *Int.J.Curr.Microbiol.App.Sci*. 8(10): 2738-2742.
doi: <https://doi.org/10.20546/ijcmas.2019.810.316>
- Central Seed Certification Board (2013). Indian Minimum Seed Certification Standards. Ministry of Agriculture, GOI, New Delhi. pp. 183-184.
- Desai, T.B., Madhu Bala and R.K. Patel (2023). Genetic Divergence in Sunnhemp (*Crotalaria juncea*L.), *Legume Research*. 46(4): 413-416. doi: 10.18805/LR-4397.
- Devamani, B.D., Saifulla, M., Jayappa and Sab, J. (2017). Prevalence of Seed mycoflora of Mung bean in Karnatak, India, *Int. J. Microbiol. App. Sci*. 6(6): 844-852
- Doijode, S. D. 1995. Effect of silica gel and storage containers on viability and vigour in onion. *Seed Res.*, 18:163-165.
- Ellis, R.H., Hong, T.D. and Roberts, E.H. 1990. Moisture content and the longevity of seeds of *Phaseolus vulgaris*L. *Ann. Bot.*, 66: 341-348.
- Gidrol, X., Noubhani, A., Mocquot, B., Fournier, A. and Pradet, A. 1998. Effect of accelerated aging on protein synthesis in two legume seeds. *Plant Physiol.Biochem.*, 26: 281-288.
- Govindaraj, M., P. Masilamani and V. Alex Albert. 2017. Influence of Harvesting and Threshing Methods on Storability of Rice Varieties. *Madras Agricultural Journal*. 104: 395-400.
- International Seed Testing Association (ISTA). 1985. Determination of moisture content. *Seed Science and Technology* 13: 338-343.
- International Seed Testing Association (ISTA). ISTA. 1999. International Rules for Seed Testing. *Seed Science and Technology Supplement*, 27: 39.
- Lates,J.C. and B.B. Mabbayad.1983. The potential and establishment method of *Crotalaria juncea* L. as a green manure for corn (*Zea mays* L.) .*Philippines Journal of Crop Science*,8:145-147
- Iqbal, N., Shahzad, A., Basra, M. and Khalil Rehman, U. 2002. Evaluation of vigor and oil quality in cotton seed during accelerated aging. *Int. J. Agric. Biol.*, 4(3): 318-322.
- Johnny subakarivin,J., Anbuselvam, M. Surendhar and J.R. Jerish.2021. Effect of storage containers and seed treatments on germination and vigour of Black gram(*Vigna mungo*(L)Happer). *Plant Archives*. 21(1):2360-2362
- Kaliyan, N., Alagusundaram, K. and Gayathri, P. 2006.Effect of temperature and moisture content on shelf life of paddy. The American Society of Agricultural and Biological Engineers. 066-193.
- Kapoor, N., Arya, A., Siddiqui., Mohd Asif., Kumar, H. and Amir, A. 2002. Physiological and biochemical changes during seed deterioration in aged seeds of rice (*Oryza sativa* L.). *Am. J. Plant Physiol.*, 6(1): 28-35.
- Masilamani, P and A. Tajuddin 2012. Can we use combine for seed purpose?.*Kisan World*. December2012.39:38-39.
- Masilamani, Pand K. Sivasubramaniyam (2016). *Green Manure Seed Production*. Kalyani Publishers, New Delhi. pp.163.
- Masilamani., P., V. Alex albert, S. Vallalkannan and M. Govindaraj. 2017. Influence of harvesting and threshing Methods on Seed Quality of Sunn hemp (*Crotalaria juncea* L.). *Seed Research*. 45: 12-15.
- Masilamani, P. T. Eevera., T. Ramesh and S.Venkatesan. 2021. Harvesting and Threshing

- methods on Seed Quality of Dhaincha (*Sesbania aculeate*) Legume Research. DOI: 10.18805, 1-7.
- Mc Sorley, R., D.W. Dickson., J.A. De Brito., T.E. Hewlett and J.J. Frederic. 1994. Effects of tropical rotation crops on *Meloidogyne arenaria* population densities and vegetable yields in microplots. *Journal of Nematology*, 26: 175-181
- Merritt, D.J., Senaratna, T., Touchell, D.H., Dixon, K.W. and Sivasithamparam, K. 2003. Seed ageing of four Western Australian species in relation to storage environment and seed antioxidant activity. *Seed Sci.Res.*, 13: 155-165.
- Muangkaeo, R., S. Srichuwong and S. Vearasilp. 2005. Influence of packaging materials and storage time on seed viability and chemical component of rice seed. Conference on International Agricultural Research for Development, Stuttgart- Hohenheim, October 11-13, 2005
- Padma, V. and M. B. Reddy. 2002. Study on seed storage ability in rice genotypes. *Oryzae*, 39(1&2):71-75.
- Panse VG and PV Sukhatme (1995). Statistical methods for agricultural workers. Indian Council of Agricultural Research Publications, New Delhi. 175 P.
- Pradhan, S.K., S.K. Sarkar and S. Prakash. 2001. Verital response of Sunnhemp, *Crotalaria juncea* L. to *Rhizobium japonicum*(Cowpea type) with reference to dynamics of nodulation. Legume Research. 24: 164 – 168.
- Rajasekaran, R., P. Balamurugan and C. Reshma. 2005. Effect of eco-friendly seed treatments and containers on storability of niger (*Guizotia abyssinica*L. f. Cass.) cv. Paiyur 1. *The Madras Agric. J.*, 92(1-3):95-100.
- Rajendra prasad, R., P. Masilamani, and K. Balakrishnan, 2017. Effect of pre-sowing seed treatments on dormancy of sunn hemp (*Crotalaria juncea* L.). *Seed Research*. 45: 136-140.
- Ramanadane, T. and A.S. Ponnuswamy.2004. Ageing and anatomical influence on seed storability in rice (*Oryzasativa* L.) hybrids and parental lines. *Tropical Agric. Res.*, 16: 37-50.
- Rickman, J.F., M. Belland D. Shires, 2006. Seed Quality, Available at <http://www.knowledgebank.irri.org>. Accessed 21/12/2014.
- Rotar PP and RJ Joy (1983). ‘Tropic Sun’ sunn hemp (*Crotalaria juncea* L.). Research Extension Series. 36. Hawaii Institute of Tropical Agriculture and Human Resources. University of Hawaii, Honolulu.
- Sadhu. K.A. 2014. Seed borne fungi and their effect on Seed health of Green gram. *Bioscience Discovery*, 5(2):251-255.
- Saxena, O. P., G. Singh, H. Pakeeraiah and N. Pandey. 1987. Seed deterioration studies in some vegetable seeds. *Acta Horticulture*, 215:39-44.
- Sharon M.E.M, C. V. K.Abirami, K. Alagusundaram and J. A.Sujeetha.2015. Safe storage guidelines for black gram under different storage conditions. *Journal of Stored Products and Postharvest Research*. Vol. 6(5):38-47
- Singh, K.K . and Dadlani.2003. Effect of packaging on vigour and viability of Soybean (*Glycine max* (L) Merrill.) seed during ambient storage. *Seed Research*. 31(1): 27-32
- Ulemale, R.B., D.G. Giri, R.S. Shivankar and V.N. Patil. 2002. Effect of Sowing dates, rowspacing and phosphorus levels on yield and yield attributes of sunnhemp (*Crotalaria juncea*.L), *Legume Research*. 25: 273-275.
- Van Chin, D. and T.T. Kieu. 2006. Study on hermetically sealed storage system for rice seeds. *Omonrice*, 14: 64-70.
- Htwe E.M, K.T.One, E.H.Kyaw, K.Ngwe, K.K.Win.2018. Effect of Different Seed Moisture Contents and Storage Containers on Seed Quality of Green Gram(*Vigna radiata* L. Wilczek)and Chickpea(*Cicer arietinum*). *Journal of Agricultural Research*,5(2):67-75

Table 1. Effect of harvesting and threshing methods on germination of Sunn hemp

Treatments(T) Periods (P)	Cloth bag (C ₁)					Super grain bag (C ₂)				
	T ₁	T ₂	T ₃	T ₄	Mean	T ₁	T ₂	T ₃	T ₄	Mean
P ₀	98 (82.80)	98 (82.80)	98 (82.80)	98 (82.80)	98 (82.80)	98 (82.80)	98 (82.80)	98 (82.80)	98 (82.80)	98 (82.80)
P ₁	97 (80.63)	96 (79.46)	97 (80.63)	97 (80.63)	97 (80.63)	97 (80.63)	97 (80.63)	97 (80.63)	97 (80.63)	97 (80.63)
P ₂	96 (79.46)	95 (77.24)	95 (77.24)	96 (79.46)	96 (79.46)	97 (80.63)	96 (79.46)	95 (77.24)	96 (79.46)	96 (79.46)
P ₃	95 (77.24)	93 (76.06)	93 (76.06)	94 (76.94)	94 (76.94)	96 (79.46)	95 (77.24)	93 (76.06)	94 (76.94)	95 (77.24)
P ₄	94 (76.94)	91 (72.88)	92 (75.44)	93 (76.06)	93 (76.06)	95 (77.24)	93 (76.06)	92 (75.44)	93 (76.06)	93 (76.06)
P ₅	94 (76.94)	90 (72.05)	91 (72.88)	92 (75.44)	92 (75.44)	95 (77.24)	92 (75.44)	91 (72.88)	92 (75.44)	93 (76.06)
P ₆	93 (76.06)	89 (70.69)	90 (72.05)	91 (72.88)	91 (72.88)	94 (76.94)	89 (70.69)	91 (72.88)	92 (75.44)	92 (75.44)
P ₇	93 (76.06)	87 (69.25)	88 (70.47)	90 (72.05)	90 (72.05)	93 (76.06)	88 (70.47)	91 (72.88)	91 (72.88)	91 (72.88)
P ₈	92 (75.44)	86 (68.56)	87 (69.25)	89 (70.69)	89 (70.69)	93 (76.06)	85 (67.81)	89 (70.69)	90 (72.05)	89 (70.69)
P ₉	91 (72.88)	85 (67.81)	86 (68.56)	88 (70.47)	88 (70.47)	92 (75.44)	84 (66.77)	89 (70.69)	89 (70.69)	89 (70.69)
P ₁₀	89 (70.69)	84 (66.77)	84 (66.77)	87 (69.25)	86 (68.56)	91 (72.88)	83 (65.76)	88 (70.47)	89 (70.69)	88 (70.47)
P ₁₁	88(70.47)	82 (65.27)	83 (65.76)	86 (68.56)	85 (67.81)	90 (72.05)	83 (65.76)	88 (70.47)	89 (70.69)	88 (70.47)
P ₁₂	86 (68.56)	80 (63.63)	82 (65.27)	85 (67.81)	83 (65.76)	89 (70.69)	82 (65.27)	86 (68.56)	88 (70.47)	86 (68.56)
Mean	93 (76.06)	89 (70.69)	90 (72.05)	91 (72.88)	91 (72.88)	94 (76.94)	90 (72.05)	91 (72.88)	92 (75.44)	92 (75.44)
	C	P	T	CP	PT		CT		CPT	
SEd	0.317	0.807	0.448	1.142	1.614		0.633		2.283	
CD (P=0.05)	0.624	1.591	0.883	NS	NS		NS		NS	

(Figures in parentheses are arc sine transformed values) P- Storage period in months

T₁- Manual harvesting and Threshing

T₂- Manual harvesting and Mechanical threshing

T₃- Manual harvesting and Tractor Treading T₄- Combine harvester

Table 2. Effect of harvesting and threshing methods on root length (cm) of Sunn hemp

Treatments(T) Periods (P)	Cloth bag (C ₁)					Super grain bag (C ₂)				
	T ₁	T ₂	T ₃	T ₄	Mean	T ₁	T ₂	T ₃	T ₄	Mean
P ₀	18.0	16.9	17.5	17.1	17.4	20.1	19.6	20.0	20.0	19.9
P ₁	17.7	16.6	17.3	17.0	17.2	19.8	19.0	19.7	19.5	19.5
P ₂	17.5	16.2	17.1	16.7	16.9	19.4	18.5	19.1	19.0	19.0
P ₃	17.4	15.9	16.9	16.4	16.7	19.1	18.1	18.5	18.3	18.5
P ₄	17.2	15.7	16.6	16.1	16.4	18.7	17.6	18.1	17.8	18.1
P ₅	17.0	15.4	16.3	15.9	16.2	18.3	17.1	17.6	17.4	17.6
P ₆	16.7	15.1	16.1	15.6	15.9	17.9	16.7	17.0	16.9	17.1
P ₇	16.4	14.9	15.9	15.4	15.7	17.6	16.2	16.7	16.5	16.8
P ₈	16.1	14.7	15.6	15.1	15.4	17.2	15.9	16.4	16.1	16.4
P ₉	15.9	14.5	15.2	14.8	15.1	16.9	15.2	16.1	15.8	16.0
P ₁₀	15.7	14.2	15.0	14.6	14.9	16.6	14.7	15.8	15.3	15.6
P ₁₁	15.4	14.0	14.8	14.3	14.6	16.1	14.4	15.3	14.9	15.2
P ₁₂	15.1	13.8	14.6	14.1	14.4	15.8	14.2	14.8	14.4	14.8
Mean	16.6	15.2	16.1	15.6	15.9	18.0	16.7	17.3	17.1	17.3
	C	P	T	CP	PT	CT		CPT		
SEd	0.049	0.127	0.070	0.179	0.253	0.099		0.358		
CD (P=0.05)	0.098	0.250	0.139	0.353	NS	NS		NS		

T₁- Manual harvesting and Threshing

T₂- Manual harvesting and Mechanical threshing

T₃- Manual harvesting and Tractor Treading T₄- Combine harvester

Table 3. Effect of harvesting and threshing methods on shoot length (cm) of Sunn hemp

Treatments(T) Periods (P)	Cloth bag (C ₁)					Super grain bag (C ₂)				
	T ₁	T ₂	T ₃	T ₄	Mean	T ₁	T ₂	T ₃	T ₄	Mean
P ₀	18.9	17.1	17.9	17.7	17.9	22.4	20.5	21.4	21.1	21.4
P ₁	18.9	16.5	17.0	16.9	17.3	22.1	20.0	21.0	20.7	21.0
P ₂	17.8	16.3	16.9	16.5	16.9	21.8	19.6	20.6	20.2	20.6
P ₃	17.1	16.3	16.7	16.5	16.7	21.5	19.1	20.0	19.6	20.1
P ₄	16.0	15.4	15.8	15.6	15.7	20.9	18.7	19.7	19.0	19.6
P ₅	15.6	14.9	15.5	15.4	15.4	20.6	18.2	19.3	18.5	19.2
P ₆	15.3	14.5	14.8	14.7	14.8	20.1	17.6	18.8	18.2	18.7
P ₇	15.1	14.4	14.8	14.7	14.8	19.7	17.0	18.5	17.8	18.3
P ₈	14.9	14.3	14.8	14.6	14.7	19.3	16.6	18.0	17.3	17.8
P ₉	14.9	13.8	14.8	14.2	14.4	19.0	16.3	17.6	16.9	17.5
P ₁₀	14.1	13.9	14.1	14.0	14.0	18.7	15.7	17.1	16.4	17.0
P ₁₁	14.0	13.2	13.9	13.7	13.7	18.2	14.9	16.8	16.0	16.5
P ₁₂	13.9	12.3	13.7	13.6	13.4	17.8	14.5	16.4	15.6	16.1
Mean	15.9	14.8	15.4	15.2	15.4	20.2	17.6	18.9	18.3	18.7
	C	P	T	CP	PT	CT	CPT			
SEd	0.0495	0.1262	0.0699	0.1784	0.2523	0.0989	0.3568			
CD (P=0.05)	0.0976	0.2487	0.1379	0.3517	NS	0.1951	NS			

T₁- Manual harvesting and Threshing

T₂- Manual harvesting and Mechanical threshing

P- Storage period in months

T₃- Manual harvesting and Tractor Treading T₄- Combine harvester

Table4. Effect of harvesting and threshing methods on dry matter production (g) of Sunn hemp

Treatments(T) Periods (P)	Cloth bag (C ₁)					Super grain bag (C ₂)				
	T ₁	T ₂	T ₃	T ₄	Mean	T ₁	T ₂	T ₃	T ₄	Mean
P ₀	0.238	0.211	0.216	0.214	0.220	0.264	0.242	0.251	0.247	0.251
P ₁	0.220	0.190	0.201	0.195	0.202	0.260	0.235	0.247	0.242	0.246
P ₂	0.214	0.183	0.190	0.185	0.193	0.253	0.229	0.241	0.237	0.240
P ₃	0.207	0.175	0.184	0.182	0.187	0.248	0.222	0.236	0.231	0.234
P ₄	0.197	0.170	0.179	0.178	0.181	0.242	0.216	0.229	0.225	0.228
P ₅	0.193	0.167	0.177	0.172	0.177	0.237	0.206	0.220	0.217	0.220
P ₆	0.187	0.157	0.175	0.168	0.172	0.233	0.193	0.212	0.208	0.212
P ₇	0.184	0.156	0.170	0.163	0.168	0.229	0.188	0.205	0.200	0.206
P ₈	0.180	0.155	0.165	0.158	0.165	0.222	0.180	0.196	0.192	0.198
P ₉	0.177	0.150	0.158	0.154	0.160	0.216	0.172	0.189	0.183	0.190
P ₁₀	0.173	0.145	0.155	0.149	0.156	0.208	0.164	0.181	0.176	0.182
P ₁₁	0.168	0.144	0.150	0.145	0.152	0.201	0.157	0.174	0.168	0.175
P ₁₂	0.159	0.134	0.146	0.138	0.144	0.194	0.151	0.168	0.160	0.168
Mean	0.192	0.164	0.174	0.169	0.175	0.231	0.197	0.211	0.207	0.211
	C	P	T	CP	PT	CT	CPT			
SEd	0.0006	0.0015	0.0008	0.0021	0.0029	0.0012	0.0042			
CD (P=0.05)	0.0012	0.0029	0.0016	0.0042	NS	0.0023	NS			

T₁- Manual harvesting and Threshing

T₂- Manual harvesting and Mechanical threshing P- Storage period in months

T₃- Manual harvesting and Tractor Treading T₄- Combine harvester

Table 5. Effect of harvesting and threshing methods on seedling vigour index I of Sunn hemp

Treatments(T) Periods (P)	Cloth bag (C ₁)					Super grain bag (C ₂)				
	T ₁	T ₂	T ₃	T ₄	Mean	T ₁	T ₂	T ₃	T ₄	Mean
P ₀	3616	3332	3469	3410	3457	4165	3930	4057	4028	4045
P ₁	3550	3178	3327	3288	3336	4064	3783	3948	3899	3924
P ₂	3389	3088	3230	3187	3223	3996	3658	3772	3763	3797
P ₃	3278	2995	3125	3093	3122	3898	3534	3581	3563	3644
P ₄	3121	2830	2981	2948	2970	3762	3376	3478	3422	3509
P ₅	3064	2727	2894	2880	2891	3696	3248	3358	3303	3401
P ₆	2976	2634	2781	2757	2787	3572	3053	3258	3229	3278
P ₇	2930	2549	2702	2709	2722	3469	2922	3203	3121	3179
P ₈	2852	2494	2645	2643	2659	3395	2763	3062	3006	3056
P ₉	2803	2406	2580	2552	2585	3303	2646	2999	2910	2965
P ₁₀	2652	2360	2444	2488	2486	3212	2523	2895	2821	2863
P ₁₁	2587	2230	2382	2408	2402	3087	2432	2825	2750	2773
P ₁₂	2494	2088	2321	2355	2314	2990	2353	2683	2640	2667
Mean	3024	2685	2837	2825	2843	3585	3094	3317	3266	3315
	C	P	T	CP	PT	CT	CPT			
SEd	8.997	22.938	12.724	32.439	45.876	17.994	64.879			
CD (P=0.05)	17.737	45.221	25.084	63.952	90.442	35.474	NS			

T₁- Manual harvesting and Threshing

T₂- Manual harvesting and Mechanical threshing P- Storage period in months

T₃- Manual harvesting and Tractor Treading T₄- Combine harvester

Table 6. Effect of harvesting and threshing methods on Seedling vigour index II of Sunn hemp

Treatments(T) Periods (P)	Cloth bag (C ₁)					Super grain bag (C ₂)				
	T ₁	T ₂	T ₃	T ₄	Mean	T ₁	T ₂	T ₃	T ₄	Mean
P ₀	23	21	21	21	22	26	24	25	24	25
P ₁	21	18	19	19	19	25	23	24	23	24
P ₂	21	17	18	18	18	25	22	23	23	23
P ₃	20	16	17	17	18	24	21	22	22	22
P ₄	19	15	16	17	17	23	20	21	21	21
P ₅	18	15	16	16	16	23	19	20	20	20
P ₆	17	14	16	15	16	22	17	19	19	19
P ₇	17	14	15	15	15	21	17	19	18	19
P ₈	17	13	14	14	15	21	15	17	17	18
P ₉	16	13	14	14	14	20	14	17	16	17
P ₁₀	15	12	13	13	13	19	14	16	16	16
P ₁₁	15	12	12	12	13	18	13	15	15	15
P ₁₂	14	11	12	12	12	17	12	14	14	15
Mean	18	15	16	16	16	22	18	19	19	20
	C	P	T	CP	PT		CT		CPT	
SEd	0.065	0.165	0.092	0.233	0.330		0.129		0.466	
CD (P=0.05)	0.128	0.325	0.181	0.460	0.651		0.255		0.920	

T₁- Manual harvesting and Threshing

T₂- Manual harvesting and Mechanical threshing P- Storage period in months

T₃- Manual harvesting and Tractor Treading T₄- Combine harvester

Figure 2. Influence of harvesting and threshing methods on moisture content of Sunn hemp seed storage





Figure 3. Influence of harvesting and threshing methods on fungal growth of Sunn hemp seed storage in cloth bag



T₁- Manual harvesting and Threshing

T₂- Manual harvesting and Mechanical threshing

P- Storage period in months

T₃- Manual harvesting and Tractor Treading T₄- Combine harvester