

Evaluation of the Nephroprotective and Anti-hyperlipidic Potential of Methanol Leaf Extract of *Gongronemalatifolium* (Utazi) in Wistar Rat

Abstract

The aim of this study was to evaluate the nephroprotective and anti-hyperlipidemic potential of methanol leaf extract of *Gongronemalatifolium* (utazi) in albino rats. Freshly harvested leaf of *G. latifolium* was processed into fine powder and then extract. Twenty (20) adult male albino rats were divided into five groups of five rats per group. Group I was the normal control and was administered with 2 ml of distilled water. Group III and IV were pretreated with 200 and 400 mg/kg of extract respectively for 28 days prior to oral administration of 100 mg/kg of TAA on groups II, III and IV. Animals were denied food overnight and subsequently sacrificed by cervical dislocation. Collected blood samples were analyzed using standard procedures. The result obtained from the qualitative phytochemical analysis performed on the leaf of *G. latifolium* revealed that saponins and flavonoids are the most abundant of the phytochemicals reportedly present. Oral administration of methanol leaf extract of *G. latifolium* significantly ($P < 0.05$) demonstrated the potential to normalize serum lipid profile, as well as indices of renal functions. In conclusion, *Gongronemalatifolium* leaf is nephroprotective and anti-hyperlipidemic.

Keywords: Nephroprotective, Anti-hyperlipidemic, *Gongronemalatifolium*, Flavonoids

Introduction

The liver, likewise the kidneys, is one of the crucial organs of the body saddled with the task of ridding the body of metabolic waste products [1]. Very often, human beings are intentionally or unintentionally exposed to a wide array of chemical agents that harm these delicate and sensitive organs, such as the liver and kidneys, which results translates to injury to such as organs and thereby impairing their ability to optimally perform deliver their metabolic functions. An estimated 10% of the global population are victims of this problem [2]. On the other hand, the role of the liver in the synthesis of lipids cannot be overemphasized, and when injured, it can result in translate to impaired lipid homeostasis and its attendant consequences. Conventionally, dialysis and transplant are the most frequently procedures adopted for individuals with the these said problems. Unfortunately, there are characterized by shortcomings such as immunological rejection of kidney grafts, immune suppression, and theirs its attendant consequences [3].

Comment [CNBMT1]: Change the title. Suggestion title : Nephroprotective and Anti-Hyperlipidemic Effects of Methanol Leaf Extract of *Gongronema latifolium* (Utazi) against Thioacetamide-Induced Renal Injury in Wistar Rats

Formatted: Highlight

Comment [CNBMT2]: four

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Comment [CNBMT3]: TAA stand for????

The use of plant-based therapies in the treatment of human ailments dates back to prehistoric times. Its use is widespread, as evident by the fact that an estimated 80% of the population in of developing countries depend solely on it to meet their health needs [4]. *Gongronemalatifolium* is a climbing plant known for its broad, heart-shaped leaves with a characteristic sharp, bitter, and slightly sweet taste, especially when eaten fresh. The stems have soft and hairy, that yielding a milky latex or exudates [5]. It belongs is a member of the to the *Asclepiadaceae* family. *Gongronemalatifolium*, locally known to the South-Easterners of Nigeria as Utazi and commonly known as amaranth globe leaf, is an edible rainforest plant [5]. In folk medicine, it is considered a medicinal spice and vegetable, owing to the fact that it has been successfully used successfully in the treatment of diseases such as diabetes [6]; [7]. In Eastern Nigeria, the leaf is employed in soup-making for mothers who that have recently given birth, as put to bed as it is believed to stimulate appetite, reduce post-partum contraction, and enhance the return of the menstrual cycle [6]. The crude leaf extract of *G. latifolium* crude leaf extract is used in the treatment of malaria, hypertension, and as a laxative. Research efforts have shown that the leaf of said this plant contain is an embodiment of essential oil, fibre, and essential phytochemicals such as saponins, alkaloids, flavonoids, among others [8]. The plethora of existing data in existence on the therapeutic values points to the fact that the plant could be further explored further for more therapeutic benefits. Hence, the importance imperativeness of this study is defined.

Comment [CNBMT4]: Add reference here

Materials and Methods

Collection of Plant Material

Fresh leaves of *Gongronemalatifolium* (Utazi) were harvested from a farm in Uturu in Abia State, Southeast Nigeria. The leaves were subsequently identified at the herbarium unit of the Department of Forestry, Michael Okpara University of Agriculture, Umudike Abia State Southeast Nigeria.

Comment [CNBMT5]: Add reference number for id

Processing and Extraction of Plant Material

Leaves of *G. latifolia* were thoroughly washed with tap water. The leaves were dried at room temperature and afterwards, dried and ground into fine powder. 500 g of powdered *G. latifolia* leaf sample was steeped in one litre of 50% methanol for a period of 72 hours. The mixture was shaken twice daily. The solvent was filtered over a layer of gauze, and then the filtrate evaporated to dryness in vacuo at 55°C.

Comment [CNBMT6]: Be standardized. Use want to hour or h

Formatted: Highlight

Phytochemical analysis

Extract derived from leaf of *Gongronemalatifolium* (Utazi) was assayed to identify the quantity of phytochemicals present in accordance with the method described by Trease et al[9].

Animals

Adult albino rats of both sexes weighing 120-150 g were purchased from the Animal House of the Department of Science Laboratory Technology, Akanulbiam Federal Polytechnic, Uwana, Afikpo. The rats were housed in aluminium cages under standard laboratory conditions. They were given food and water *ad-libitum*. Acclimatization lasted for 14 days.

Comment [CNBMT7]: Change to wistar

Comment [CNBMT8]: Can you stage the age of rats because too thin

Formatted: Highlight

Formatted: Highlight

Median Lethal Dose 50% (LD50%)

Determination of ~~Median~~ ~~median~~ ~~L~~ethal ~~D~~ose 50% (LD50) involved two phases of experiment. At the initial phase, nine (9) adult male albino rats were divided into three groups of three rats each of which was separately administered with 10, 100 and 1000 mg/kg of extract orally. Animals were observed for 24 h for signs of toxicity. Owing to the fact that mortality was not observed after the first phase, the second phase comprising another three groups of one rat each was separately administered with 1600, 2900 and 5000 mg/kg of extract, after which animals were observed for 48 h for signs of toxicity according to Lorke [10].

Formatted: Highlight

Formatted: Highlight

Experimental design

Twenty adult albino rats were starved of food for 24 h prior to the commencement of experiment. The rats were divided into five groups of five rats per group.

Formatted: Font: 11 pt, Highlight

Formatted: Font: 11 pt, Highlight

Group 1: (Normal control) rats were administered 2 ml of distilled water

Group 2: Rats were administered TAA without treatment

Formatted: Highlight

Group 3: Rats were pretreated with 200 mg/kg of MEGLU

Group 4: Rats were pretreated with 400 mg/kg of MEGLU

Pretreatment with extract lasted for 28 days during which the body weight was determined on weekly basis, while TAA was administered on the 28th day by a single dose subcutaneous injection of 100 mg/kg of TAA. Animals were denied food overnight, and subsequently sacrificed by cervical dislocation. Collected blood samples collected in plain bottles for analysis.

Comment [CNBMT9]: plain tubes

Biochemical analysis

To determine creatinine and urea, 2 mL of blood introduced into plain tube was subjected to centrifugation at 4,000 rpm for 15 min and the plasma obtained was stored for biochemical analysis. Kits were used to determine the levels of urea and creatinine.

Formatted: Highlight

Comment [CNBMT10]: Can you add company name for the kits that use here.

Formatted: Highlight

Determination of lipid profile

Cholesterol, HDL and triacylglyceride levels were estimated from serum by CHOD-PAP. LDL and HDL were calculated. While the atherogenic index was calculated using the method described by Muruganandan et al [11].

Statistical Analysis

Data obtained from the study were expressed as Mean \pm standard deviation using SPSS (Ver. 23). Data were analysed using one way analysis of variance (ANOVA). Variation in mean values was compared using Turkey Test. *P*-values less than 0.05 was considered statistically significant.

Table 1: Qualitative phytochemical composition of *Gongronemalatifolium* (Utazi)

Phytochemicals	Abundance
Saponins	+++
Tannins	+
Flavonoids	+++
Alkaloids	++
Glycosides	+
Phenol	+

+ [abundant], ++[more abundant], +++[most abundant]

Table 2: Lipid Profile of Rats administered Aqueous Stem Bark *Gongronema Latifolium* (Utazi)

Treatment	TC (mg/dl)	TG (mg/dl)	HDL (mg/dl)	LDL (mg/dl)
Normal Ctrl (2 ml of distilled H ₂ O)	200.00±5.22 ^a	60.00±5.51 ^a	41.32±3.01 ^a	127.34±5.02 ^a
Negative control (induction without treatment)	270.00±6.64 ^d	100.00±1.15 ^e	70.23±5.60 ^d	203.76±2.08 ^c
MEGLU ₂₀₀ mg/kg TAA ₁₀₀ mg/kg	226.37±0.89 ^{cd}	77.01±4.80 ^d	63.03±2.82 ^c	132.32±2.42 ^{bc}
MEGLU ₄₀₀ mg/kg+TAA ₁₀₀ mg/kg	223.86±5.28 ^c	73.42±2.30 ^c	60.00±3.72 ^b	129.76±4.82 ^b

Results are expressed as mean ± standard deviation from five determinations. Values with the same superscripts in a column are not significantly different ($p < 0.05$).

Comment [CNBMT11]: Add abbreviation for TC, TG, HDL and LDL.

Comment [CNBMT12]: Add abbreviation for TC, TG, HDL and LDL.

Table 3: Renal function markers of Rats administered with *G. Latifolia* (Utazi)

Groups	Urea	Creatinine
Normal Ctrl (2 ml of distilled H ₂ O)	5.63 ±0.73 ^a	74.22±6.280 ^a
Negative control (induction without treatment)	6.34±0.70 ^c	88.20±4.3 ^c
MEGLU ₂₀₀ mg/kg+TAA ₁₀₀ mg/kg	5.92±0.91 ^b	77.00±3.52 ^b
MEGLU ₄₀₀ mg/kg+TAA ₁₀₀ mg/kg	5.88±0.61 ^{ab}	77.01±5.541 ^b

Results are expressed as mean ± standard deviation from five determinations. Values with the same superscript in a column are not significantly different ($p < 0.05$).

Table 4: Body Weight Changes in Rats administered with Methanol Leaf Extract of *Gongronema latifolium*

Groups	Body weight			
	WK 1	WK 2	WK 3	WK 4
Control	150.0±3.48 ^a	159.0±5.83 ^{ab}	165.6±6.82 ^b	168.4±6.73 ^b
Negative control	148.3±2.56 ^c	147.2±4.78 ^c	144.6±3.45 ^b	140.3±5.34 ^a
MEGLU ₂₀₀ mg/kg TAA ₁₀₀ mg/kg	156.2±4.23 ^a	164.1±7.18 ^b	172.0±4.34 ^c	175.2±6.62 ^c
MEGLU ₄₀₀ mg/kg+TAA ₁₀₀ mg/kg	156.4±3.39 ^a	158.0±2.72 ^a	160.6±4.21 ^a	166.0±3.37 ^b

Results are expressed as mean ± standard deviation from five determinations. Values with the same superscripts in a column are not significantly different ($p < 0.05$).

Formatted: Font: 11 pt

Formatted: Font: 11 pt

Formatted: Font: 11 pt

Formatted: Font: 11 pt

Formatted: Font: 11 pt

Formatted: Font: 11 pt

Formatted: Font: 11 pt

Discussion & Conclusions

The liver likewise the kidneys is one of the crucial organs of the body saddled with the task of riding the body of metabolic waste products [1]. Very often, human beings are intentionally or unintentionally exposed to a wide array of chemical agents that harm these delicate and sensitive organs such as the liver and kidney which translates to injury to such as organs. Table 1 shows

Formatted: Font: 12 pt, Font color: Black, Highlight

the qualitative phytochemical composition of methanol leaf extract of *Gongronema latifolia* indicating that saponins and flavonoids are most abundant of all the phytochemicals reportedly present, while glycosides and phenols are the least abundant phytochemicals in the leaf of *G. latifolia*. Table 2 shows the lipid profiles of rats administered with the aqueous methanol leaf extract of *G. ongronema latifolium*, indicating that the oral administration of thioacetamide (TAA) increased the levels of Triacylglyceride (TG), Total cholesterol (TC), High-density Lipoprotein (HDL), and Low-density Lipoprotein (LDL). However, administration of 400 mg/kg of the methanol leaf extract of *G. ongronema latifolium*, resulted in a significant ($P < 0.05$) reduction in the levels of the aforementioned lipids, although they were still which though were significantly ($p < 0.05$) higher than those reported for the normal control.

Comment [CNBMT13]: Rephrase the sentences because it similar with introduction.

Formatted: Font: 12 pt, Font color: Black, Highlight

The ability of the extract to maintain a stable lipid profile could be attributed to its phytochemical constituents, some of which may possess have antioxidant properties. This finding is consistent with the results conducted in tandem with the with the outcome of a work by by Rosemary et al. [12], demonstrated which proved the hypoglycemic and hypolipidemic effect of *G. ongronema latifolium* extract on healthy subjects. Biochemically, a pronounced elevation in the levels of renal parametres is suggestive of renal alterations. Table 3 presents shows the renal function markers of rats administered with *G. latifolia* leaf extract, showing that serum urea and creatinine levels were significantly ($P < 0.05$) increase raised in Group II following the induction of renal damage with oral administration of TAA. However, a contrary

observation was made in Groups III and IV. ~~pretreated with the said~~ the extract was prior to ~~administ~~ administered ~~ration~~ prior to the administration of TAA. The observed decreased in serum creatinine and urea levels ~~observed~~ in groups III and IV could be attributed to the effect of the reactive oxygen species (ROS) generated by TAA. This finding is consistent with the study conducted ~~finding of~~ by Omodale et al. [13], which demonstrated established that the protective effect of *G. latifolium* root extract ~~of Gongronemalatifolium root extract protected~~ against kidney damage. Table 4 shows the changes of body weight ~~changes~~ in rats administered with the methanol leaf extract of *G. latifolia*, indicating that the body weight of rats at the 4th week of feed intake was significantly ($p < 0.05$) higher than that reported at week 1 for Groups I, III and IV. However, a contrary observation was made in Group II.

References

- [1] Chaturvedi R, George S, John A. Preventive and protective effect of wild basil in ethanol-induced liver toxicity in rats. *British J. Biomedical Sci.* 2007; 64:10-12.
- [2] World Kidney Day; Chronic Kidney Disease, 2015. Available from: <http://www.worldkidneyday.org/faqs/chronic-kidney-disease/>. Accessed December 10, 2017.
- [3] Sands JM, Verlanders JW. Anatomy and Physiology of the Kidney. In: Toxicology of the Kidney 3rd ed (Tarloff JB, Lash, LH eds). CRC Press, Boca Raton, FL, 2005, 3-56.
- [4] Akinyemi, KO, Oladapo O, Okwara CE, Ibe CC, Fasura KA. Screening of crude extracts of six medicinal plants used in South-West Nigerian unorthodox medicine for anti-methicillin resistant *Staphylococcus aureus* activity. *BMC Complement Alternat Med.* 2005; 5:6-10.
- [5] Dalziel JM. *Gongronemalatifolium*. In: Dalziel JM, eds. The Useful plants of Tropical West Africa. 1st ed. London: Crown Overseas Agents Colonies; 1937: 461.
- [6] Morebise O, Fafunso MA, Makinde JM, Olajide OA, Awe EO. Anti-inflammatory property of the leaves of *Gongronemalatifolium*; *Phytother Res.* 2002;16:S75-7.

[7] Okafor JC. Tropical plants in health care delivery in Nigeria: contributions of some members of Compositae (Asteraceae). J Coll Med. 2002; 7(2):129-31.

[8] Ugochukwu NH, Babady NE. Antihyperglycemic effect of aqueous and ethanolic extracts of *Gongronemalatifolium* leaves on glucose and glycogen metabolism in livers of normal and streptozotocin-induced diabetic rats. Life Sci. 2003; 73(15):1925-38.

[9] Trease GE, Evans WC. A Textbook of Pharmacognosy. Baillière Tindall, London; 2001.

[10] Lorke D. A new approach to practical acute toxicity testing. Arch Toxicol. 1983;54 (4):275-287.

[11] Muruganandan S, Scrivivasan K, Gupta S, Gupta PK, Lal J. Effect of mangiferin on hyperglycemia and atherogenicity in streptozotocin diabetic rats. J Ethnopharmacol. 2005;97:497- 501

[12] Rosemary AA and Joseph EA. Effects of *Gongronemalatifolium* on blood lipids, lipoproteins and glucose values in adult Nigerians. Int J Res Med Sci. 2015; 3(4):891-895.

[13] Omodale PE, Enitan SS, Adejumo EN, Akinleye WA and Bella AS. Protective effect of aqueous root extract of *Gongronemalatifolium* against paracetamol induced hepatotoxicity and chloroquine induced nephrotoxicity in rats. International Journal of Herbal Medicine 2017; 5(3): 115-120.