

Report on the occurrence of Cicada, *Platypleura octoguttata* (Hemiptera: Cicadidae) on [Eucalyptus](#) in Gujarat

Formatted: Font: Not Italic, Font color: Blue

Formatted: Font color: Blue

Abstract:

In order to determine the cicadas spotted on the [Eucalyptus plants](#) grown nearby to the college, Anand Agricultural University, Jabugam, between April and June 2022, an investigation was conducted. Adults were brought to the Department of Entomology lab at the College of Agriculture, Anand Agricultural University, Jabugam, for identification in addition to do further research. It is confirmed that the pest is cicadas on the Eucalyptus plant that have been reported from Gujarat, India, based on the morphological traits of the adults and DNA fingerprinting.

Formatted: Font: Not Italic, Font color: Blue

Formatted: Font color: Blue

Introduction

The genus *Platypleura* Amyot & Audinet-Serville, 1843 (Order Hemiptera; Family Cicadidae; Subfamily Cicadinae; Tribe Platycleurini) contains a group of homopterous insects, of which the cicada is the most diverse. It is found throughout Africa (Villet 1997), the Indian subcontinent (Distant, 1906), Southeast Asia (Lee 2008, 2009, 2010), and East (Hayashi 1974; Lee and Hayashi 2003). According to Sanborn (2014), the genus contains 101 taxonomically distinct species worldwide, 21 of which have been found on the Indian subcontinent. India has the largest global diversity of cicadas in terms of generic diversity. Like many other insects in the area, the cicadas of the Indian subcontinent have not received enough research since the early 20th century (Price *et al.*, 2016).

Distant's description of over 100 cicada species that are found on the Indian subcontinent is now accepted as accurate. With 148 cicada species recognized in 44 genera, "The wildlife of British India, including Ceylon and Burma" (Distant, 1906) gave the first treatment of the region as a whole. There were 172 cicada species in 47 genera after this, according to an addendum (Distant, 1916), which identified an additional 24

species and three new genera for the area. The two described species of Mata kama (Distant, 1881), which was initially recorded from Darjeeling in West Bengal (Distant, 1906), and Mata rama (Distant, 1912), which was described from Bhutan, are both found in the India region (Sanborn 2013). Mata kama is distributed in India, Java, Malay states (Metcalf 1963) and Nepal (Duffels and Van der Laan 1985) and Mata rama is known only from Bhutan and China (Sanborn 2013; Price *et al.*, 2016).

The cicadas are big insects that prefer different types of trees as hosts, but they have also caused damage to other woody plants when they emerge periodically. They naturally enjoy the sun. The males typically make their calls in the morning sun from the highest branches of large trees. As the day wears on, they descend to the undergrowth and frequently make their calls from the little shrubs next to the stream (Sanborn, 2008). Cicada "songs" can be heard whenever there are adults around, from early in the morning to late at night. The male cicadas are the ones who sing. The male cicadas' hollow abdomen acts as a resonating chamber and amplifies the sound. The tymbal's sound it produced. A row of four flexible ribs is present on each tymbal. A pulse of sound is produced as the cicadas flex his tymbals, causing the ribs to bend sequentially. There are no sound-producing organs in the female cicadas.

The cicadas (Hemiptera: Cicadidae) were observed in Gujarat between April and June 2022 at the Eucalyptus plants cultivated close to the College of Agriculture, Anand Agricultural University, Jabugam. Adults were brought in for identification in the Department of Entomology laboratory at the College of Agriculture, Anand Agricultural University, Jabugam for further study. It is confirmed that the pest is cicadas, which are on Eucalyptus plants and have been reported from Gujarat, India, based on the morphological characteristics of the adults and DNA fingerprinting. According to a research of the literature from the accessible sources, Gujarat's Eucalyptus plants infestation by cicadas has not been documented earlier.

Method

The molecular characterization of cicada was carried out to generate information on species authentication. The cicadascicadae's specimens (10 Nnos.) were collected

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font: Not Italic, Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font: Not Italic, Font color: Blue

Formatted: Font: Not Italic, Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

from the infested Eucalyptus trees situated in experimental plot of College of Agriculture, Anand Agricultural University, Jabugam and preserved in 70% ethyl alcohol and stored at -20 °C until further use. Wings of the specimen were used to isolate genomic DNA of the specimen using DNA extraction kit (QIAGEN DNeasy blood and tissue kit, Germany) following the manufacturer's protocol. The extracted DNA was quantified using Nanodrop spectrophotometer. The concentration of DNA was ranged between 30-110 ng μl^{-1} . The good quality genomic DNA free from protein/RNA contamination was used for further analysis.

Formatted: Font: Not Italic, Font color: Blue

Formatted: Font color: Blue

Results and Discussion

The genomic DNA extracted from the specimen was subjected to PCR amplification of COXI (Cytochrome c oxidase subunit 1) gene following the standard protocol (Hebert *et al.*, 2003). Primers used for amplification of COXI gene were forward primer (LCO 1490 5'-GGTCAACAAATCATAAAGATATTGG-3') and reverse primer (HCO 2198 5'-TAAACTTCAGGGTGACCAAAAATCA-3'). The amplified fragments of COXI gene were analyzed on 1.5 % agarose gel electrophoresis. The single amplicon of COXI gene of size 650 bp was visualized during the analysis. The PCR amplified products were purified by using QIAquick Gel Extraction kits to remove residual dNTPs, primers and salts. The purified PCR product was subjected to automated DNA sequencing on ABI 3500 Genetic analyzer (Applied Biosystem, USA). The PCR amplicons were sequenced bidirectionally. Forward and reverse sequences of COXI gene were assembled using BioEdit/clone manager software. Consensus sequence thus obtained was used for BLAST (Basic Local Alignment Search Tool) search in National Center for Biotechnology Information (NCBI). The COXI generated consensus sequence of cicada found infesting eucalyptus tree has been deposited in NCBI GenBank database (NCBI accession no. **OP563725**).

Formatted: Font color: Blue

Formatted: Font color: Blue, Superscript

➤ *Platypleura octoguttata* cytochrome oxidase subunit I (COI) gene, partial cds; mitochondrial

TATTTTCATTTTTGGTATTTGATCCGGGATAGTTGGGACTGCATTAAGATTTT
ATTCTGAATCGAATTAGGAATACCTGGTCTTTTATTGGAGATGATCAAATTTA

TAATGTGATTGTAACAGCTCATGCTTTTATTATAATCTTTTTTATAGTTATGCC
TATTATGATTGGGGGATTGGGAATTGACTGGTTCCTTGATGATTGGTGCC
CTGACATGGCATTCCCTCGTATAAATAATATAAGATTTTGACTTTTACCACCA
TCTTTAACACTTTTATTAATAGGAAGAATAATTGATAGAGGGGCTGGAAGT
GATGAACTGTTTATCCTCCTTTATCAAGAGTTATATATCACTCTGGTTCCTGTG
TTGACATAACTATTTTTTCTTTACATTTAGCAGGAGTTTCTTCAATTTAGGAG
CTGTAAATTTTATTAGAACAACTTCAATATACGATCCACAGGTATATTTTG
GATCGAACACCTTTATTTGTATGGGCTGTTTAATTACAGCTTTTTTGTATTG
CTATCATTACCTGTTTTAGCAGGTGCAATTAATGCTTACGGATCGTAA
TTAAATACATTTTTTTGATCCTGCAGGTGGGGGAGATCCAATTTTGTATC
AACATTTGTTTT

Conclusion

Based on the morphometric and supporting molecular analysis data, the cicada found infesting eucalyptus tree was identified as *Platypleura octoguttata*. This report appears to be the first report of natural occurrence of cicada, *Platypleura octoguttata* on eucalyptus in Gujarat, India.

Formatted: Font color: Blue

UNDER PEER

References:

Click the references in and out of the research and write them as the pattern of the journal.

Formatted: English (India)

- Distant, W. L. (1881). Descriptions of new species belonging to the Homopterous Family Cicadidae. *Transactions of the Royal Entomological Society of London*, 1881, 627–648. <https://doi.org/10.1111/j.1365-2311.1881.tb00885.x>
- Distant, W. L. (1906). The fauna of British India, including Ceylon and Burma. Rhynchota. Volume- III. *Taylor and Francis, London*, pp. 526.
- Distant, W. L. (1912) New genera and species of Oriental Homoptera. *The Annals and Magazine of Natural History, Series 8-9*, 459–471. <https://doi.org/10.1080/00222931208693156>
- Distant, W. L. (1916). The fauna of British India, including Ceylon and Burma. Rhynchota - Volume- VI. *Taylor and Francis, London*, pp. 248.
- Duffels, J. P. & Van der Laan, P. A. (1985). *Catalogue of the Cicadoidea (Homoptera, Auchenorrhyncha) 1956–1980. Series Entomologica 34*. Dr. W. Junk Publishers, Dordrecht, pp. 414
- Duffels, J. P. (1993). The systematic position of *Moana expansa* (Homoptera, Cicadidae), with reference to sound organs and the higher classification of Cicadoidea. *Journal of National History*, 27, 1223-1237.
- Hayashi, M. (1974). The cicadas of the genus *Platypleura* (Homoptera, Cicadidae) in the Ryukyu Archipelago, with the description of a new species, *Kontyû*, 42, 232–253.
- Hebert, P. D., Cywinska, A., Ball, S. L., & deWaard, J. R. (2003). Biological identifications through DNA barcodes. *Proceedings. Biological sciences*, 270(1512), 313–321. <https://doi.org/10.1098/rspb.2002.2218>
- Illiger, J. C. W. (1800). Vierzig neue Insekten aus der Hellwigischen Sammlung in Braunschweig. *Archiv für Zoologie und Zootomie*, 1, 103-150.
- Lee, Y. J. & Hayashi, M. (2003). Taxonomic review of Cicadidae (Hemiptera, Auchenorrhyncha) from Taiwan, part 1. *Platypleurini, Tibicenini, Polyneurini, and Dundubiini (Dundubiina)*. *Insecta Koreana*, 20 (2), 149–185.
- Lee, Y. J. (2008). A checklist of Cicadidae (Insecta: Hemiptera) from Vietnam, with some taxonomic remarks. *Zootaxa*, 1787, 1– 27.

- Lee, Y. J. (2009). Cicadas (Hemiptera: Cicadidae) from Panay, Philippines, with a new species and a new genus. *Journal of Asia- Pacific Entomology*, 12, 293–295. <https://doi.org/10.1016/j.aspen.2009.06.007>
- Lee, Y. J. (2010). Cicadas (Hemiptera: Cicadidae) of Mindanao, Philippines, with the description of a new genus and a new species. *Zootaxa*, 2351, 14–28.
- Metcalf, Z. P. (1963). General catalogue of the Homoptera. Fascicle VIII. Cicadoidea. Part 1. Cicadidae. Section II. Gaeaninae and Cicadinae. *North Carolina State College Contribution*, 1502, 587–919.
- Price, B., Allan, E., Marathe, K., Sarkar, V., Simon, C. & Kunte, K. (2016). The cicadas (Hemiptera: Cicadidae) of India, Bangladesh, Bhutan, Myanmar, Nepal and Sri Lanka: an annotated provisional catalogue, regional checklist and bibliography. *Biodiversity Data Journal* 4: e8051. DOI: [10.3897/BDJ.4.e8051](https://doi.org/10.3897/BDJ.4.e8051)
- Sanborn, A. F. (2008). New records of Brazilian cicadas including the description of a new species (Hemiptera: Cicadoidea, Cicadidae). *Neotropical Entomology*, 37, 685-690.
- Sanborn, A. (2013) *Catalogue of the Cicadoidea (Hemiptera: Auchenorrhyncha). With contributions to the bibliography by Martin H. Villet*. Elsevier/Academic Press, San Diego, pp.1001 <https://doi.org/10.1016/B978-0-12-416647-9.00001-2>
- Sanborn, A. F. (2014) *Catalogue of the Cicadoidea (Hemiptera: Cicadoidea). With contributions to the bibliography by Martin H. Villet*. Elsevier/Academic Press, San Diego, pp. 1001
- Villet, M. H. (1997). Redescription of three species of the genus, *Platypleura* Amyot & Serville 1843 (Hemiptera : Cicadidae). *Tropical Zoology*, 10, 321–332. <https://doi.org/10.1080/03946975.1997.10539345>