

Effect of moisture conservation practices on yield and economics of pigeonpea (*Cajanus cajan* L.) in black soil.

Formatted: Line spacing: single

ABSTRACT

A field experiment was conducted at Main Agricultural Research Station, Raichur in for a period of two seasons viz., 2021 and 2022 was laid out in a Split-Split design with four main, two sub and two sub-sub treatments with three replications to evaluate the effect of conservation agricultural practices on growth and yield of pigeonpea. The trial was laid out in a Split plot design with four main, two sub and two sub-sub treatments with three replications. The treatments with flat bed, Compartment bund, Ridge and furrow and Broad bed and furrow were taken as main treatments. The sub treatments include treatments with mulching and without mulching, an sub treatment and FYM and RDF & FYM an sub sub treatment. Application of recommended dose of fertilizers (RDF) and RDF along with farm yard manure (FYM) are categorized as sub sub treatments. The results clearly indicated that, Among all treatments ridges and furrow (M₃) method along with mulching coupled with application of FYM and RDF and FYM was found to have exhibited positive influence of yield and economics of pigeonpea. Adopting the ridge and furrow method can be approved as an effective strategy for optimizing the grain yield (1555 kg ha⁻¹ and 1599 kg ha⁻¹). The pooled mean of two years data per hectare indicated for the above treatment resulted in lowest cost of cultivation (Rs. 40675.49), highest gross income (Rs. 112864.70) and net returns (Rs. 72189.21) and cost of cultivation, gross returns, net returns and BC ratio (46991.36 Rs ha⁻¹, 116024.7 Rs ha⁻¹, 72189.2 Rs ha⁻¹ and 2.78) in 2021 and 2022 in pigeonpea crop respectively with benefit cost ratio of 2.78.

Formatted: Font: Italic

Keywords: Pigeonpea, growth and yield attributes, grain yield,

Introduction

Conservation agriculture (CA) means is a set of soil management practices that minimizes the soil disruption of the soil structure, compaction and enhances the natural biodiversity. These practices found to improve the fertility status, variation in total nitrogen

content, different fraction of labile carbon pools along with different physical and biological activities of soil which generally contributes the overall quality of the soil (Add reference). The advantage of CA in terms of better soil quality considers through lower bulk density (BD), increased water holding capacity, higher aggregate stability and better soil structure. On the other hand, CA practices influences soil biological parameters which affect the overall soil quality as it improves the quality, quantity and distribution of organic matter in soil which is a vital factor that largely affects biomass, diversity and activity of soil microorganism as it is the main source of food for soil biota (Reicosky and Saxton, 2007).

Conservation agriculture has proven potential to improving the long-term environmental and financial sustainability of farming (Add reference). The technology of conservation agriculture provides opportunities to reduce the cost of production, lowers weed infestation, saves water and nutrients besides, increase yield. It maintains a soil cover through surface retention of crop residue with no tillage or zero tillage and reduced tillage thus acts as. Potential benefits of CA include reduction in cost of production, reduced incidence of weeds, saving water and nutrients, increased yields, resource improvement and providing environmental benefits. Further Also, it improves physical, chemical and biological qualities of the soil.

In pigeonpea, studies on evaluating effect of soil conservation practices are limited and scanty. Hence, assessing effect of soil conservation measures on soil physical, chemical and biological properties, nutrient content and its uptake and yield in sole crop stand of pigeonpea on which present study is aimed occupies its importance.

Considering these above views, the present investigation with the title “Effect of moisture conservation practices on yield and economics of pigeonpea (*Cajanus cajan* L.) in black soil” was undertaken in black soils with the following objective to study the effect of conservation agriculture practices on growth attributes and yield of pigeonpea.

Material and methods

The field experiment was conducted at Main Agricultural Research Station, Raichur, The experiment was laid out in a Split-Split plot design with four main, two sub and two sub-sub treatments with three replications for study vizas detailed in Table 1.,

Chart 1. Treatment details. Table 1. Details of Treatments imposed in Pigeonpea

Treatment Details	Black soil
-------------------	------------

- Formatted: Font: Bold, Not Italic
- Formatted: Centered
- Formatted Table

Main plot: Moisture conservation practices (Interbund management)

M ₁ - Flat bed	-
M ₂ - Compartment bund	5.4 m × 7.5 m
M ₃ - Ridges and furrow	60 cm furrow
M ₄ - Broad bed and furrow	120 cm bed
Subplot:	
S ₁ - Crop residue mulching	18-20 t ha ⁻¹
S ₂ - Without mulching	-
Sub- sub-plot:	
N ₁ - RDF N: P ₂ O ₅ :K ₂ O	25:50:0 kg ha ⁻¹
N ₂ - RDF with FYM	RDF + FYM 6.0 t ha ⁻¹

The soils of the experimental site belong to medium deep black soil and clay texture, neutral in soil reaction (7.6) and low in electrical conductivity (0.22 dSm⁻¹). The soil organic carbon content was 4.6 g kg⁻¹ and available N (263.42 kg ha⁻¹), available phosphorus (28.68 kg P₂O₅ ha⁻¹) and available potassium (451 kg K₂O ha⁻¹), exchangeable calcium and magnesium (24.82 and 7.86 C mol (p⁺) kg), available sulphur (13.30 mg kg⁻¹). ~~The monthly meteorological data for the period of experimentation 2021-2022 and 2022-2023 were collected at Meteorological Observatory, Main Agricultural Research Station, Raichur. Annual mean rainfall received was 613 mm and 474 mm during 2021-22 and 2022-23 respectively. However, the distribution of rainfall was erratic. The seed yield of Pigeonpea as influenced by various treatments was recorded per plot and expressed as kg ha⁻¹. For economic evaluation cost of cultivation, gross returns, net returns and B:C ratio were calculated per plot and expressed as Rs. ha⁻¹~~

Formatted: Superscript

MENTION STATISTICAL TOOLS USED FOR ANALYSIS**Results and discussion**

The yield parameters in pigeonpea as influenced by conservation practices were discussed herewith and economic analysis with respect to net returns and B:C ratio was also interpreted.

Effect of different soil moisture conservation practices Seed-on seed yield (kg ha⁻¹) of Pigeonpea

During the years 2021 and 2022, the analysis of seed yield (kg ha^{-1}) revealed that the ridges and furrow (M_3) method consistently had significantly higher values compared to the broad bed and furrow (M_4) and flat-bed (M_1) methods (Table 42). The seed yield values were 1555 kg ha^{-1} , 1424 kg ha^{-1} and 1212 kg ha^{-1} for the ridges and furrow (M_3) method in 2021, and 1599 kg ha^{-1} , 1460 kg ha^{-1} and 1238 kg ha^{-1} in 2022, respectively. ~~When considering the~~The pooled mean data ~~from both years, inferred that,~~ the ridges and furrow (M_3) method consistently recorded higher seed yield values of 1577 kg ha^{-1} , 1442 kg ha^{-1} and 1225 kg ha^{-1} compared to the broad bed and furrow (M_4) and flat-bed (M_1) methods, respectively. ~~Based on these findings, the ridges and furrow (M_3) method is recommended for achieving better seed yield (kg ha^{-1}). The findings suggest that the ridge and furrow method can be a valuable technique for improving the seed yield of pigeonpea. The results indicated that B~~by creating raised ridges and sunken furrows, the method optimizes water distribution and reduces waterlogging, leading to improved plant growth and higher seed yield. The results are in similar with findings of (Palaniappan *et al.*, 2009 and Pandey *et al.*, 2015).

~~Similarly, in the years 2021 and 2022, the analysis showed that t~~The crop residue mulching (S_1) method had significantly higher seed yield values compared to the without mulching (S_2) method (Table 42). ~~The seed yield values were 1509 kg ha^{-1} and 1321 kg ha^{-1} in 2021, and 1548 kg ha^{-1} and 1353 kg ha^{-1} in 2022 for the crop residue mulching (S_1) and without mulching (S_2) methods, respectively.~~The pooled data from both years ~~also~~ consistently showed higher seed yield values of 1528 kg ha^{-1} for the crop residue mulching (S_1) method compared to 1337 kg ha^{-1} for the without mulching (S_2) method (Table 42). Therefore, the crop residue mulching (S_1) method ~~is~~ can be recommended for achieving better seed yield (kg ha^{-1}). This demonstrated that the crop residue mulching method positively impacted the seed yield of pigeonpea. Incorporating crop residues as mulch in pigeonpea cultivation can be a sustainable and effective approach to enhance seed yield (Kumawat *et al.*, 2013).

The analysis of seed yield (kg ha^{-1}) in 2021 and 2022 did not show any significant difference between the FYM with RDF (N_2) and RDF N: P_2O_5 : K_2O (N_1) fertilizers (Table 42). ~~However, the FYM with RDF (N_2) consistently had higher seed yield values compared to the RDF N: P_2O_5 : K_2O (N_1) fertilizer. In 2021, the values were 1440 kg ha^{-1} for FYM with RDF (N_2) and 1391 kg ha^{-1} for RDF N: P_2O_5 : K_2O (N_1), while in 2022, the values were 1475 kg ha^{-1} and 1425 kg ha^{-1} , respectively (Table 1). When the data from both years was~~The

Formatted: Superscript

pooled data, ~~the indicated that~~ FYM with RDF (N₂) ~~still~~ exhibited higher values of 1457 kg ha⁻¹ compared to 1408 kg ha⁻¹ for the RDF N:P₂O₅:K₂O (N₁) fertilizer (Table 42). The non-significant results might indicate that the particular combination of FYM with RDF (N₂) and RDF N:P₂O₅:K₂O (N₁) did not result in noticeable differences in seed yield compared to the control or other treatments.

In the pooled data for 2021 and 2022, the different three-way combinations for seed yield (kg ha⁻¹) did not show any significant difference (Table 5) (Figure 1). The combination of ridges and furrow (M₃), crop residue mulching (S₁) and FYM with RDF (N₂) had the highest seed yield value of 1721 kg ha⁻¹, followed by ridges and furrow (M₃), crop residue mulching (S₁) and RDF N:P₂O₅:K₂O (N₁) with a value of 1646 kg ha⁻¹ and compartment bund (M₂), crop residue mulching (S₁) and FYM with RDF (N₂) with a value of 1617 kg ha⁻¹ (Table 42). Similar trends were observed for individual years. ~~It is important to note that the lack of significance does not necessarily imply that these combinations are ineffective or unimportant. Other factors such as environmental conditions, genetic variability and management practices may also contribute to seed yield variation.~~

Comment [E1]: No figure seen in manuscript

To study the effectEffect of different soil moisture conservation practices on economics of pigeonpea

~~Economic analysis is one of the major criteria for evaluating efficiency and economical effect of conservation agriculture practices in pigeonpea.~~ The data on cost of cultivation, gross returns, net returns and benefit cost ratio of conservation agriculture practices in pigeonpea production ~~for per hectare~~ as influenced by various treatments is depicted in Table 23 to 56.

Cost of Cultivation (₹ ha⁻¹)

The cost of cultivation for different conservation agriculture practices as influenced by various treatments is depicted in Table 23.

Based on pooled data, it was found that M₁ recorded least cost of cultivation among M₁, M₂, M₃ and M₄. Highest cost of cultivation is associated with ridges and furrow method (M₃) followed by compartment bund (M₂), broad bed and furrow (M₄) while flatbed method was found to have least cost of cultivation and hence can be considered as cheapest method with the cost of Rs.39578.77, 43833.42, 43190.67, 42869.36 and 39578.77, respectively.

Among the crop residue mulching (S₁) method and without mulching (S₂) methods, S₁

was found to be costlier than S_2 method. Cost involved in using S_1 and S_2 was Rs. 45048.36 and 39687.75 indicating that using without mulching can reduce the cost of the cultivation.

Significantly higher cost of cultivation was recorded with the application of RDF $N:P_2O_5:K_2O$ (N_1) compared to FYM with RDF (N_2). N_1 has the highest cost of Rs. 45543.78 while it was 39192.33 for N_2 indicating its cost efficiency over other method to adopt among conservation agriculture practices.

Among the combinations of treatments or conservation agriculture practices, flat-bed method (M_1), without mulching (S_2) method and RDF $N:P_2O_5:K_2O$ (N_1) recorded lowest cost of cultivation (36301.11) while ridges and furrow (M_3) method, crop residue mulching (S_1) method and FYM with RDF (N_2) was found to be costliest approach among them. Though this combination among conservation agriculture practices involves high cost of cultivation, it can incur profit as it contributed high yield of pigeon compared to other methods.

Gross Returns (₹ ha⁻¹)

The data pertaining to the gross returns for different conservation agriculture practices as influenced by various treatments is depicted in Table 34.

~~During 2021-22, highest gross return (Rs. 112871.13) was observed for ridges and furrow (M_3) method and least (Rs. 88459.23) was noticed in flat bed (M_1) among M_1 , M_2 , M_3 and M_4 . During 2022-23, pattern remained same with the values Rs. 116018.33 and Rs. 90388.53, respectively. Based on the pooled data, it can be concluded~~Pooled mean data for two years inferred that ridges and furrow method is feasible approach with high gross return (Rs. 114444.73) while the flatbed method involves lowest gross returns (Rs. 89398.88).

Among N_1 and N_2 , FYM with RDF (N_2) (Rs. 104621.26) was found to have highest gross return against RDF $N:P_2O_5:K_2O$ (N_1) (Rs. 101145.15) when estimated during 2021 while the pattern remained same for the year 2022 with the values of Rs. 107221.61 and Rs. 103678.94, respectively. Pooled data revealed that N_2 (Rs. 105921.43) incurs high gross returns compared to N_1 (Rs. 102412.04).

Comparison of gross returns among crop residue mulching (S_1) method and without mulching (S_2) method indicated that crop residue mulching (S_1) method results in more gross returns with an amount of Rs. 109551.49 and Rs. 96214.92 during 2021 and 2022,

Formatted: English (United States)

Formatted: Body Text

respectively. Gross returns based on pooled data also indicated to draw similar conclusions with a highest gross return of Rs. 110972.69.

Among the combinations of treatments, it was found that ridges and furrow (M_3) method, crop residue mulching (S_1) method and FYM with RDF (N_2) results in highest gross returns of Rs. 124674.55 while flatbed method (M_1), without mulching method (S_2) and RDF N:P₂O₅:K₂O (N_1) has the lowest gross returns. Similar pattern has been observed for the individual years. Hence, ridges and furrow (M_3) method, crop residue mulching (S_1) method and FYM with RDF (N_2) can be followed to attain high gross returns.

Net Returns (₹ ha⁻¹)

The data pertaining to the net returns for different conservation agriculture practices as influenced by various treatments is depicted in Table 45.

During 2021-22, highest net return (Rs. 69037.71) was observed for ridges and furrow (M_3) method and least (Rs. 48880.46) was noticed in flat-bed (M_1) among M_1 , M_2 , M_3 and M_4 . During 2022-23, pattern remained same with the values Rs. 72184.91 and 50759.77, respectively. Based on the pooled data, it can be concluded that ridges and furrow method is feasible approach with high net returns (Rs. 70611.31) while the flatbed method involves lowest net returns (Rs. 49820.12).

Among N_1 and N_2 , RDF N:P₂O₅:K₂O (N_1) (Rs. 61952.82) was found to have highest gross return against FYM with RDF (N_2) (Rs. 59077.48) when estimated during 2021 while the pattern remained same for the year 2022 with the values of Rs. 64486.60 and Rs. 61677.84, respectively. Pooled data revealed that N_1 (Rs. 63219.71) incurs high gross returns compared to N_2 (Rs. 60377.66).

Comparison of net returns among crop residue mulching (S_1) method and without mulching (S_2) method indicated that crop residue mulching (S_1) method results in more net returns with an amount of Rs. 61952.82 and Rs. 64486.60 during 2021 and 2022, respectively. Pooled data also indicated that S_1 is better which results in highest net return (Rs. 63219.71).

Formatted: English (United States)

Formatted: Body Text

Among the combinations of treatments, it was found that ridges and furrow (M_3) method, crop residue mulching (S_1) method and RDF N:P₂O₅:K₂O (N_1) results in highest net returns of Rs. 76052.89 while flatbed method (M_1), without mulching (S_2) and FYM with RDF (N_2) has the lowest net returns (Rs. 49307.54). Similar pattern has been observed for the individual years. Hence, ridges and furrow (M_3) method, crop residue mulching (S_1) method along with RDF N:P₂O₅:K₂O (N_1) can be followed to obtain high net returns.

BC Ratio

The data pertaining to the BC ratio for different conservation agriculture practices as influenced by various treatments is depicted in Table 56.

Among four methods viz., flat-bed (M_1), compartment bund (M_2), ridges and furrow (M_3) method and broad bed and furrow (M_4), during 2021, BC ratio was found to be highest for ridges and furrow (M_3) method followed by compartment bund (M_2), broad bed and furrow (M_4) and least for flat-bed (M_1) with the values 2.59, 2.48, 2.42 and 2.25, respectively. Trend remained same for the year 2022 with BC ratios of 2.66, 2.53, 2.48 and 2.30, respectively. Further, Pooled data also indicated that M_3 is more efficient method obtain high BC ratio (2.62).

With respect to RDF N:P₂O₅:K₂O (N_1) and FYM with RDF (N_2), N_1 was found to have highest BC ratio of 2.58 indicating its economic feasibility compared to FYM with RDF (N_2) with an BC ratio of 2.29. During 2022, N_1 and N_2 had values of 2.64 and 2.35, respectively. The BC ratio was also high for the N_1 (2.61) based on pooled data of the two years indicating that approach is consistent in yielding higher returns.

Comparison of BC ratio among crop residue mulching (S_1) method and without mulching (S_2) method indicated that both the methods had on par with each other for BC ratio during 2021 (2.44 and 2.43) and 2022 (2.50 and 2.49), respectively. However, crop residue mulching (S_1) method had numerical superiority over without mulching (S_2) method in both the years. Pooled data also indicated that S_1 has numerical superiority compared to S_2 with BC ratio 2.47 and 2.46, respectively.

Among the combinations of treatments, it was found that ridges and furrow (M_3) method, crop residue mulching (S_1) and RDF N:P₂O₅:K₂O (N_1) results in highest BC ratio of 2.76 while flatbed method (M_1), without mulching (S_2) method and FYM with RDF (N_2) has

the lowest BC ratio (2.15). Similar pattern has been observed for the individual years. Hence, ridges and furrow (M_3) method, crop residue mulching (S_1) along with RDF $N:P_2O_5:K_2O$ (N_1) can be followed to obtain high benefit cost ratio by the farmers.

Conclusion: The combination of the ridges and furrow method, crop residue mulching and the application of FYM with RDF emerged as the most effective approach, consistently demonstrating superior performance across multiple parameters. By implementing these recommended practices, farmers and agricultural practitioners can optimize pigeonpea crop productivity, improve soil health and ultimately achieve better results.

References

- Kumawat, Narendra., Singh, R.P., Kumar, Rakesh., Yadav, T.P. and Om, H., 2015, Effect of integrated nutrient management on productivity, nutrient uptake and economics of rainfed pigeonpea (*Cajanus cajan*) and blackgram (*Vigna mungo*) intercropping system. *Indian J. Agric. Sci.*, 85(2): 171-176.
- Palaniappan, S. P., Chandrasekaran, A., Kang, D. S., Singh, K., Rajput, R. P., Kauraw, D. L., Velayutham, M. and Lal, R., 2009, Sustainable management of natural resources for food security and environmental quality: case studies from India—a review. *Climate Change, Intercropping, Pest Control and Beneficial Microorganisms*: pp339-372.
- Pandey, I. B., Tiwari, S., Pandey, R. K. and Kumar, R., 2014, Effect of bed configuration, fertilizer levels and placement method on the productivity of long duration pigeonpea (*Cajanus cajan* (L) Millsp) under rainfed condition. *J. Food Legumes*, 27(3): 206-209.
- Reicosky, D. C. and Saxton, K. E., 2007, The benefits of no-tillage. In: *Baker CJ, Saxton KE, Ritchie WR, Chamen WCT, Reicosky DC, Ribeiro MFS, Justice SE, Hobbs PR* (eds) *No-tillage seeding in conservation agriculture*. 2nd edn. CABI, Wallingford, pp 11–20.

Table 1. Effect of different soil moisture conservation practices on seed yield per hectare

M × S × N		Seed Yield (kg/ha)									
		2021			2022			Pooled			
		S ₁	S ₂	Mean (M×N)	S ₁	S ₂	Mean (M×N)	S ₁	S ₂	Mean (M×N)	
M ₁	N ₁	1247.35	1100.75	1174.05	1273.33	1124.10	1198.71	1260.34	1112.43	1186.38	
	N ₂	1269.05	1234.22	1251.63	1295.28	1261.00	1278.14	1282.16	1247.61	1264.89	
M ₂	N ₁	1563.22	1353.15	1458.19	1600.04	1384.55	1492.30	1581.63	1368.85	1475.24	
	N ₂	1595.23	1367.34	1481.29	1639.56	1395.83	1517.69	1617.40	1381.58	1499.49	
M ₃	N ₁	1625.65	1444.98	1535.32	1667.34	1484.29	1575.82	1646.50	1464.63	1555.57	
	N ₂	1694.70	1458.45	1576.58	1747.42	1498.70	1623.06	1721.06	1478.58	1599.82	
M ₄	N ₁	1508.42	1285.96	1397.19	1553.00	1320.80	1436.90	1530.71	1303.38	1417.05	
	N ₂	1571.54	1330.19	1450.87	1611.12	1357.81	1484.46	1591.33	1344.00	1467.66	
Mean S		1509.39	1321.88		1548.39	1353.38		1528.89	1337.63		
M × S			Mean M	M × S			Mean M	M × S			Mean M
M	M ₁	1258.20	1167.49	1212.84	1284.31	1192.55	1238.43	1271.25	1180.02	1225.64	
	M ₂	1579.23	1360.24	1469.74	1619.80	1390.19	1505.00	1599.52	1375.22	1487.37	
	M ₃	1660.17	1451.71	1555.94	1707.38	1491.49	1599.44	1683.78	1471.60	1577.69	
	M ₄	1539.98	1308.08	1424.03	1582.06	1339.30	1460.68	1561.02	1323.69	1442.36	
N × S			Mean N	N × S			Mean N	N × S			Mean N
N	N ₁	1486.16	1296.21	1391.19	1523.43	1328.43	1425.93	1504.79	1312.32	1408.56	
	N ₂	1532.63	1347.55	1440.09	1573.35	1378.33	1475.84	1552.99	1362.94	1457.97	
SOV	S.Em ±	CD at 5 %		S.Em ±	CD at 5 %		S.Em ±	CD at 5 %			
M	36.53	126.42		34.28	118.63		27.29	94.44			
S	23.25	75.81		12.63	41.20		12.50	40.78			
N	24.10	NS		22.78	NS		16.88	NS			
M×S	46.49	NS		25.26	NS		25.01	NS			
M×N	48.20	NS		45.55	NS		33.75	NS			
S×N	34.08	NS		32.21	NS		23.87	NS			
M×S×N	68.17	NS		64.42	NS		47.73	NS			

Comment [E2]: Below all tables mention M: Main plots, S: Sub Plots, N: sub sub plots

Table 23. Cost of cultivation for different soil moisture conservation practices in pigeon pea

Cost of cultivation (Rs ha ⁻¹)				
M × S × N		Pooled		
		S ₁	S ₂	Mean (M×N)
M ₁	N ₁	38822.75	33779.48	36301.11
	N ₂	44982.95	40729.89	42856.42
M ₂	N ₁	42858.72	37347.95	40103.33
	N ₂	49116.18	43439.81	46278.00
M ₃	N ₁	43326.85	38024.13	40675.49
	N ₂	49856.33	44126.38	46991.36
M ₄	N ₁	42494.98	36883.81	39689.39
	N ₂	48928.12	43170.53	46049.32
Mean S		45048.36	39687.75	
M × S			Mean M	
M	M ₁	41902.85	37254.69	39578.77
	M ₂	45987.45	40393.88	43190.67
	M ₃	46591.59	41075.25	43833.42
	M ₄	45711.55	40027.17	42869.36
N × S			Mean N	
N	N ₁	41875.83	36508.84	39192.33
	N ₂	48220.90	42866.65	45543.78
SOV		S.Em ±	CD at 5 %	
MP		-	-	
SP		-	-	
MP*SP		-	-	
SSP		-	-	
MP*SSP		-	-	
SP*SSP		-	-	
MP*SP*SSP		-	-	

Table 34. Gross return for different soil moisture conservation practices in pigeonpea

M × S × N		Gross returns (Rs ha ⁻¹)									
		2021			2022			Pooled			
		S ₁	S ₂	Mean (M×N)	S ₁	S ₂	Mean (M×N)	S ₁	S ₂	Mean (M×N)	
M ₁	N ₁	90884.13	80549.19	85716.66	92839.15	82262.76	87550.95	91861.64	81405.98	86633.81	
	N ₂	92485.75	89917.86	91201.80	94395.74	91856.49	93126.12	93440.74	90887.17	92163.96	
M ₂	N ₁	113357.43	98447.15	105902.29	116039.44	100734.17	108386.80	114698.44	99590.66	107144.55	
	N ₂	115662.70	99471.25	107566.97	118883.31	101553.02	110218.17	117273.00	100512.14	108892.57	
M ₃	N ₁	117870.21	104927.36	111398.79	120889.29	107771.92	114330.61	119379.75	106349.64	112864.70	
	N ₂	122767.55	105919.40	114343.48	126581.54	108830.58	117706.06	124674.55	107374.99	116024.77	
M ₄	N ₁	109439.24	93686.48	101562.86	112682.71	96212.06	104447.39	111060.97	94949.27	103005.12	
	N ₂	113944.88	96800.67	105372.77	116839.91	98832.29	107836.10	115392.39	97816.48	106604.44	
Mean S		109551.49	96214.92		112393.89	98506.66		110972.69	97360.79		
M × S			Mean M	M × S			Mean M	M × S			Mean M
M	M ₁	91684.94	85233.53	88459.23	93617.44	87059.63	90338.53	92651.19	86146.58	89398.88	
	M ₂	114510.06	98959.20	106734.63	117461.38	101143.59	109302.49	115985.72	100051.40	108018.56	
	M ₃	120318.88	105423.38	112871.13	123735.42	108301.25	116018.33	122027.15	106862.31	114444.73	
	M ₄	111692.06	95243.57	103467.82	114761.31	97522.18	106141.74	113226.68	96382.88	104804.78	
N × S			Mean N	N × S			Mean N	N × S			Mean N
N	N ₁	107887.75	94402.54	101145.15	110612.65	96745.23	103678.94	109250.20	95573.89	102412.04	
	N ₂	111215.22	98027.30	104621.26	114175.13	100268.09	107221.61	112695.17	99147.70	105921.43	
SOV		S.Em ±	CD at 5 %		S.Em ±	CD at 5 %		S.Em ±	CD at 5 %		
MP		2553.99	8837.96		2382.70	8245.23		1898.00	6567.93		
SP		1590.83	5187.98		876.04	2856.91		849.37	2769.94		
MP*SP		3181.66	NS		1752.07	NS		1698.73	NS		
SSP		1681.72	NS		1589.61	NS		1181.46	NS		
MP*SSP		3363.44	NS		3179.22	NS		2362.92	NS		
SP*SSP		2378.31	NS		2248.05	NS		1670.84	NS		
MP*SP*SSP		4756.62	NS		4496.09	NS		3341.68	NS		

Table 45. Net return for different soil moisture conservation practices in pigeonpea

M × S × N		Net returns (Rs ha ⁻¹)									
		2021			2022			Pooled			
		S ₁	S ₂	Mean (M×N)	S ₁	S ₂	Mean (M×N)	S ₁	S ₂	Mean (M×N)	
M ₁	N ₁	52061.38	46769.72	49415.55	54016.40	48483.29	51249.84	53038.89	47626.50	50332.70	
	N ₂	47502.79	49187.96	48345.38	49412.79	51126.60	50269.69	48457.79	50157.28	49307.54	
M ₂	N ₁	70498.71	61099.20	65798.95	73180.72	63386.22	68283.47	71839.71	62242.71	67041.21	
	N ₂	66546.52	56031.44	61288.98	69767.13	58113.21	63940.17	68156.82	57072.32	62614.57	
M ₃	N ₁	74543.36	66903.23	70723.30	77562.43	69747.80	73655.11	76052.89	68325.51	72189.21	
	N ₂	72911.22	61793.02	67352.11	76725.21	64704.20	70714.73	74818.21	63248.61	69033.43	
M ₄	N ₁	66944.26	56802.67	61873.52	70187.73	59328.26	64758.01	68565.99	58065.46	63315.71	
	N ₂	65016.76	53630.15	59323.45	67911.79	55661.77	61786.78	66464.27	54645.96	60555.11	
Mean S		64503.12	56527.17		67345.52	58818.92		65924.32	57673.04		
M × S			Mean M	M × S			Mean M	M × S			Mean M
M	M ₁	49782.09	47978.84	48880.46	51714.59	49804.94	50759.77	50748.34	48891.89	49820.12	
	M ₂	68522.61	58565.32	63543.97	71473.92	60749.71	66111.82	69998.27	59657.52	64827.89	
	M ₃	73727.29	64348.13	69037.71	77143.82	67226.00	72184.91	75435.55	65787.06	70611.31	
	M ₄	65980.51	55216.41	60598.46	69049.76	57495.01	63272.38	67515.13	56355.71	61935.42	
N × S			Mean N	N × S			Mean N	N × S			Mean N
N	N ₁	66011.93	57893.70	61952.82	68736.82	60236.39	64486.60	67374.37	59065.05	63219.71	
	N ₂	62994.32	55160.64	59077.48	65954.23	57401.44	61677.84	64474.27	56281.04	60377.66	
SOV	S.Em ±	CD at 5 %		S.Em ±	CD at 5 %		S.Em ±	CD at 5 %			
MP	2553.99	8837.96		2382.70	8245.23		1898.00	6567.93			
SP	1590.83	5187.98		876.04	2856.91		849.37	2769.94			
MP*SP	3181.66	NS		1752.07	NS		1698.73	NS			
SSP	1681.72	NS		1589.61	NS		1181.46	NS			
MP*SSP	3363.44	NS		3179.22	NS		2362.92	NS			
SP*SSP	2378.31	NS		2248.05	NS		1670.84	NS			
MP*SP*SSP	4756.62	NS		4496.09	NS		3341.68	NS			

Table 56. BC ratio for different soil moisture conservation practices in pigeonpea

M × S × N		BC Ratio									
		2021			2022			Pooled			
		S ₁	S ₂	Mean (M×N)	S ₁	S ₂	Mean (M×N)	S ₁	S ₂	Mean (M×N)	
M ₁	N ₁	2.34	2.38	2.36	2.39	2.44	2.41	2.37	2.41	2.39	
	N ₂	2.06	2.21	2.13	2.10	2.26	2.18	2.08	2.23	2.15	
M ₂	N ₁	2.64	2.64	2.64	2.71	2.70	2.70	2.68	2.67	2.67	
	N ₂	2.35	2.29	2.32	2.42	2.34	2.38	2.39	2.31	2.35	
M ₃	N ₁	2.72	2.76	2.74	2.79	2.83	2.81	2.76	2.80	2.78	
	N ₂	2.46	2.40	2.43	2.54	2.47	2.51	2.50	2.43	2.47	
M ₄	N ₁	2.58	2.54	2.56	2.65	2.61	2.63	2.61	2.57	2.59	
	N ₂	2.33	2.24	2.29	2.39	2.29	2.34	2.36	2.27	2.31	
Mean S		2.44	2.43		2.50	2.49		2.47	2.46		
M × S		Mean M			M × S		Mean M		M × S		Mean M
M	M ₁	2.20	2.30	2.25	2.24	2.35	2.30	2.22	2.32	2.27	
	M ₂	2.50	2.46	2.48	2.56	2.52	2.54	2.53	2.49	2.51	
	M ₃	2.59	2.58	2.59	2.66	2.65	2.66	2.63	2.62	2.62	
	M ₄	2.45	2.39	2.42	2.52	2.45	2.48	2.49	2.42	2.45	
N × S		Mean N			N × S		Mean N		N × S		Mean N
N	N ₁	2.57	2.58	2.58	2.64	2.64	2.64	2.60	2.61	2.61	
	N ₂	2.30	2.29	2.29	2.36	2.34	2.35	2.33	2.31	2.32	
SOV	S.Em ±	CD at 5 %			S.Em ±	CD at 5 %		S.Em ±	CD at 5 %		
MP	0.06	0.21			0.06	0.19		0.05	0.16		
SP	0.04	NS			0.02	NS		0.02	NS		
MP*SP	0.08	NS			0.04	NS		0.04	NS		
SSP	0.04	0.12			0.04	0.12		0.03	0.09		
MP*SSP	0.08	NS			0.08	NS		0.06	NS		
SP*SSP	0.06	NS			0.05	NS		0.04	NS		
MP*SP*SSP	0.12	NS			0.11	NS		0.08	NS		