

# Effect of Integrated Nutrient Management on Growth, Yield, Nutrient Uptake and Economics, of *Rabi Sorghum (Sorghum bicolor (L.) Moench)*

## Abstract

A field experiment was conducted at post graduate experimental farm of M.S. Swaminathan School of Agriculture, Centurion University of Technology and Management, Paralakhemundi during the rabi season, 2022. The experimental soil was sandy loam with low in available nitrogen and medium in phosphorus and potassium. The experimental design was randomized block design. The experiment consists of 10 numbers of treatments and three replications. The treatments are T<sub>1</sub>: Control, T<sub>2</sub>: 50% RDF, T<sub>3</sub>: 75%RDF, T<sub>4</sub>: 100%RDF, T<sub>5</sub>: 50%RDF+5t FYM ha<sup>-1</sup>, T<sub>6</sub>: 50% RDF + 10t FYM ha<sup>-1</sup>, T<sub>7</sub>: 75% RDF + 5t FYM ha<sup>-1</sup>, T<sub>8</sub>: 75%RDF+10t FYM ha<sup>-1</sup>, T<sub>9</sub>: 100%RDF + 5t FYM ha<sup>-1</sup>, T<sub>10</sub>: 100%RDF + 10t FYM ha<sup>-1</sup>. Among all the treatments significantly highest growth, yield attributes and nutrient uptake were recorded from the treatment 100%RDF + 10t FYM ha<sup>-1</sup> (T<sub>10</sub>). Application of 100%RDF + 10t FYM recorded highest sorghum grain yield (2416 kg ha<sup>-1</sup>) and gross return (₹ 85108/ha<sup>-1</sup>). However, maximum net return (₹ 48010/- ha<sup>-1</sup>) was recorded from 100%RDF + 5t FYM ha<sup>-1</sup> but B:C ratio (1.49) from 75%RDF.

**Keywords:** RDF, FYM, sorghum, growth, yield, nutrient uptake, economics


## 1. Introduction

Sorghum, also known as 'Jowar' in India, which is considered as the most important among all millet crops. It is the fifth most important cereal in the world followed by wheat, rice, maize and barley. Worldwide, sorghum is grown on 40.25 million hectares of land, with an average yield of 58.7 million tonnes in 2020 (FAOSTAT, 2022). Whereas, in India in 2020-2021 sorghum production was 4.78 million tonnes from 4.24 million hectares with an average yield of 1128 kg/ha (DES, DA&FW, GoI, 2022). While in Odisha the average yield was too low (633 kg/ha) with the total production of 3.68 thousand tonnes from an area of 5.81 thousand ha in 2019-20 (DA&FP, Odisha, 2020). The main sorghum production countries are the USA and India. Sorghum is grown under recommended dose of 80kg N, 40kg P<sub>2</sub>O<sub>5</sub> and 40kg K<sub>2</sub>O per hectare. Integrated use of different sources of plant nutrients such as organic manure and bioinoculants in combination with chemical fertilizers improves soil fertility and

crop yield (Ghosh *et al.*, 2003). Judicious use of FYM with chemical fertilizers improves soil's physical, chemical, and biological properties and improves crop productivity (Sharma *et al.* 2007). Conventionally, nutrient management in sorghum production has relied heavily on use of chemical fertilizers and often applied in excessive amounts without considering the crop's nutrient needs or the characteristics of the soil. This unbalanced approach has led to nutrient imbalances, reduced soil fertility, increased fertilizer costs, and environmental pollution through nutrient runoff and leaching. Application of FYM and inorganic nitrogen levels recorded higher grain and fodder yield in sorghum fields (Shinde *et al.*, 2001). The basic concept behind integrated nutrient management INM is to keep soil fertility high by using all accessible sources of fertilizer. The amount of organic matter in soil affects the nutrients that are available to plants for nutrient uptake (Sarkar, 2005). The present experiment was conducted to study the impact of integrated nutrient management on growth, yield and economics of sorghum.

## 2. Materials and method

A field experiment was conducted at Post Graduate research farm of MSSoA, Centurion University of Technology and Management, Paralakhemundi (23<sup>0</sup>39 N latitude and 87<sup>0</sup>42 E Longitude) to study the effect of integrated nutrient management on the growth and yield of rabi sorghum. Meteorological data recorded during the crop period has been mentioned in Fig 1. Initial soil analysis had been done before sowing. The soil of the experimental site was sandy loam, acidic in reaction with pH 6.4, medium in organic carbon 0.3%, deficient in nitrogen (228kg ha<sup>-1</sup>) low in phosphorus (14 kg ha<sup>-1</sup>), and medium in potassium (120 kg ha<sup>-1</sup>). The experiment was laid out in randomized block design (RBD) with 10 treatments and each treatment was replicated thrice. The treatments are T<sub>1</sub>: Control, T<sub>2</sub>: 50% RDF, T<sub>3</sub>: 75%RDF, T<sub>4</sub>: 100%RDF, T<sub>5</sub>: 50%RDF+5t FYM ha<sup>-1</sup>, T<sub>6</sub>: 50% RDF + 10t FYM ha<sup>-1</sup>, T<sub>7</sub>: 75% RDF+5t FYM ha<sup>-1</sup>, T<sub>8</sub>: 75%RDF+10t FYM ha<sup>-1</sup>, T<sub>9</sub>: 100%RDF + 5t FYM ha<sup>-1</sup>, T<sub>10</sub>: 100%RDF+10t FYM ha<sup>-1</sup>. The recommended doses of fertilizer were applied through urea, SSP, and MOP. Urea application was done at two splits, one as basal and another as top dressing at 30 DAS. Sorghum (SSG 59-3) was sown in rabi with the spacing 45 cm x 15 cm and seed rate 8 kg ha<sup>-1</sup>. The growth attributes and yield attributes as per the schedule period were recorded. After harvesting the crop yield and economics was calculated. The data were analysed statistically by the analysis of variance (ANOVA) in MS excel 2010 and the significance of different sources of variation were tested by EMS and F-test at 5% probability.



**Fig. 1: Graph on meteorological data recorded during the crop period**

### 3. Results and discussion

#### 3.1 Growth attributes

Growth attributes of sorghum such as plant height, no. of leaves plant<sup>-1</sup>, plant dry matter accumulation, leaf area index and stem girth at 90 DAS were mentioned in Table 1. The treatment T<sub>10</sub>: 100%RDF+10t FYM ha<sup>-1</sup> recorded significantly highest plant height (250.8cm), no. of leaves plant<sup>-1</sup> (9.0), plant dry matter accumulation (564 g m<sup>-2</sup>), leaf area index (2.88) and stem girth (4.1cm) which was also showed significantly at par with the treatments viz. T<sub>9</sub>: 100%RDF+ 5t FYM ha<sup>-1</sup>, T<sub>8</sub>: 75%RDF+ 10t FYM ha<sup>-1</sup> and T<sub>4</sub>:100% RDF for all the above parameters. Whereas, in case of plant height and no. of leaves plant<sup>-1</sup> T<sub>10</sub>: 100%RDF+10t FYM ha<sup>-1</sup> also remained on par with T<sub>7</sub>: 75%RDF+5t FYM ha<sup>-1</sup>. But the data showed in leaf area index indicated that T<sub>10</sub> also statistically at par with T<sub>7</sub>: 75%RDF+5t FYM ha<sup>-1</sup>, T<sub>6</sub>: 50%RDF+10t FYM ha<sup>-1</sup> and T<sub>3</sub>: 75% RDF. However, significantly lowest growth attributes recorded from T<sub>1</sub>: Control (where no nutrient application was done). Higher plant height, no. of leaves plant<sup>-1</sup>, stem girth and LAI in T<sub>10</sub> also resulted higher plant dry matter accumulation in T<sub>10</sub>. Sorghum is a very susceptible and nitro-positive crop. So, the increase of fertilizer level and FYM level increases the height of the plant. Similar results showed that application of 100% recommended dose of fertilizer (RDF) significantly increased plant height than by using 75% or 50% RDF and applying 10 tons of FYM ha<sup>-1</sup> (Kushwaha *et al.*, 2014). The combination of organics and inorganics will boost the leaf number of plants, this concept was mentioned by Sher *et al.* (2022), Sabhad *et al.* (2020) and Afzal *et al.* (2011). Combination of fertilizer and FYM might be provided optimum nutrient to plants by modifying soil and plant environment suitable for better performance of plant morphology as well as physiology, which resulted overall crop growth (Priya *et al.*, 2014).

**Table 1. Effect of integrated nutrient management on growth attributes of rabi sorghum**

Treatments	Plant height (cm)	No. of leaves plant <sup>-1</sup>	Dry matter accumulation (g m <sup>-2</sup> )	LAI	Stem girth(cm)
T <sub>1</sub> : Control	140.5	7.0	382	2.13	2.7
T <sub>2</sub> : 50% RDF	194.9	7.3	352	2.26	2.9
T <sub>3</sub> : 75% RDF	207.2	7.2	445	2.56	3.2
T <sub>4</sub> :100% RDF	245.2	8.5	508	2.69	3.6
T <sub>5</sub> : 50%RDF+ 5t FYM ha <sup>-1</sup>	198.6	7.4	435	2.48	3.0
T <sub>6</sub> : 50%RDF+10t FYM ha <sup>-1</sup>	200.7	7.9	470	2.62	3.3
T <sub>7</sub> : 75%RDF+5t FYM ha <sup>-1</sup>	243.2	8.4	504	2.63	3.4
T <sub>8</sub> : 75%RDF+ 10t FYM ha <sup>-1</sup>	246.5	8.7	516	2.76	3.6
T <sub>9</sub> : 100%RDF+ 5t FYM ha <sup>-1</sup>	248.6	8.8	544	2.83	3.9
T <sub>10</sub> : 100%RDF+10t FYM ha <sup>-1</sup>	250.8	9.0	564	2.88	4.1
S. Em. (±)	14.2	0.3	19.27	0.14	0.17
C. D. (P=0.05)	42.8	0.9	57	0.4	0.5

**at 90 DAS****3.2 Yield attributes**

After the sorghum crop was harvested, yield parameters viz., number of ear head plant<sup>-1</sup>, length of ear head plant<sup>-1</sup>(cm), dry weight of ear head(g), grain weight earhead<sup>-1</sup>(g), and test weight(g) were recorded and presented in Table 2. Effect of integrated nutrient management (INM) significantly influenced yield attributes of rabi sorghum such as no. of ear head plant<sup>-1</sup>, length of ear head, dry weight of ear head and grain weight ear head<sup>-1</sup>. Whereas, weight of 1000 seeds (test weight) were not influenced by treatments of INM. Among all the treatments significantly highest no. of ear head plant<sup>-1</sup> (1.67), length of ear head (35.7cm), dry weight of ear head (40.4g) and grain weight ear head<sup>-1</sup> (33.6g) were recorded from T<sub>10</sub> (100%RDF+10t FYM ha<sup>-1</sup>). Whereas T<sub>1</sub> (control) resulted lowest value of all the parameters. In case of length of ear head plant<sup>-1</sup> and grain weight ear head<sup>-1</sup> the treatment T<sub>10</sub> was statistically at par with T<sub>4</sub>, T<sub>6</sub>, T<sub>7</sub>, T<sub>8</sub> and T<sub>9</sub>. The data on no. of ear head plant<sup>-1</sup> showed that T<sub>10</sub> was statistically on par with T<sub>4</sub>, T<sub>8</sub> and T<sub>9</sub> but the data on dry weight of ear head showed that T<sub>10</sub> was statistically at par with only T<sub>9</sub>. Additional supply of FYM with 75% RDF and 100% RDF recorded higher yield attributes, which might be due to better nutrient supply through organic and inorganic sources for long period that affects crop growth and photosynthetic activity (Tomar *et al.*, (2017).

**Table 2. Effect of integrated nutrient management on yield attributes of rabi sorghum**

Treatments	Yield Attributes				
	Number of ear head plant <sup>-1</sup>	Length of ear head plant <sup>-1</sup> (cm)	Dry weight of ear head (g)	Grain weight ear head <sup>-1</sup> (g)	Test weight (g)
T <sub>1</sub> : Control	1.04	27.9	21.0	17.5	9.9
T <sub>2</sub> : 50% RDF	1.18	31.0	30.2	27.1	10.5
T <sub>3</sub> : 75% RDF	1.24	31.3	32.2	27.9	10.4
T <sub>4</sub> : 100% RDF	1.47	34.5	34.5	30.9	10.5
T <sub>5</sub> : 50%RDF+ 5t FYM ha <sup>-1</sup>	1.22	31.2	32.3	28.0	10.5
T <sub>6</sub> : 50%RDF+10t FYM ha <sup>-1</sup>	1.31	31.5	32.4	28.4	10.5
T <sub>7</sub> : 75%RDF+5t FYM ha <sup>-1</sup>	1.44	34.2	33.4	30.7	10.5
T <sub>8</sub> : 75%RDF+ 10t FYM ha <sup>-1</sup>	1.48	34.7	34.8	31.3	10.5
T <sub>9</sub> : 100%RDF+ 5t FYM ha <sup>-1</sup>	1.51	35.2	36.7	31.9	10.5
T <sub>10</sub> : 100%RDF+10t FYM ha <sup>-1</sup>	1.67	35.7	40.4	33.6	11.2
S. Em. (±)	0.09	1.4	1.3	1.7	0.23
C. D. (P=0.05)	0.2	4.2	3.8	5.1	NS

### 3.3. Yield


The grain yield, stover yield and harvest index are presented in Table 3. The highest grain yield was recorded from 100%RDF+10t FYM ha<sup>-1</sup>(T<sub>10</sub>) which was statistically at par with 100%RDF+5t FYM ha<sup>-1</sup> (T<sub>9</sub>), 75%RDF+10t FYM ha<sup>-1</sup> (T<sub>8</sub>), 75%RDF+5t FYM ha<sup>-1</sup> (T<sub>7</sub>) and 100%RDF ha<sup>-1</sup> (T<sub>4</sub>). With the application of extra 10 t FYM in a hectare there was increase in 7.2% grain yield in the treatment 100%RDF+10t FYM ha<sup>-1</sup>(T<sub>10</sub>) over 100% RDF. This shows that addition of FYM with RDF has effect on improving sorghum yield. Similar findings are also recorded by Raja *et al.*, (2010). The lowest grain yield was recorded from the control (T<sub>1</sub>) where no nutrient application was done. The highest stover yield was found from the T<sub>10</sub> and T<sub>10</sub> was statistically remained on par with the treatments viz., T<sub>3</sub>, T<sub>4</sub>, T<sub>7</sub>, T<sub>8</sub> and T<sub>9</sub>. The lowest stover yield was obtained from the control treatment (T<sub>1</sub>) where no nutrient application has done. Harvest index of sorghum was not influenced significantly by the different treatments of integrated nutrient management. The beneficial effect of 100%RDF along with FYM might be associated with sufficient availability of nutrients throughout the growth period especially during critical growth stages, which resulted vigorous growth and yield attributes that ultimately resulted better yield (Jat *et al.*, 2014).

**Table 3. Effect of integrated nutrient management on yield of rabi sorghum.**

Treatments	Yield (kg ha <sup>-1</sup> )		
	Grain	Stover	Harvest index (HI)
T <sub>1</sub> : Control	978	2449	28
T <sub>2</sub> : 50% RDF	1950	4313	31
T <sub>3</sub> : 75% RDF	2103	4952	30
T <sub>4</sub> : 100% RDF	2242	5065	31
T <sub>5</sub> : 50% RDF+ 5t FYM ha <sup>-1</sup>	2086	4760	30
T <sub>6</sub> : 50% RDF+10t FYM ha <sup>-1</sup>	2172	4879	31
T <sub>7</sub> : 75% RDF+5t FYM ha <sup>-1</sup>	2240	4993	31
T <sub>8</sub> : 75% RDF+ 10t FYM ha <sup>-1</sup>	2252	5198	30
T <sub>9</sub> : 100% RDF+ 5t FYM ha <sup>-1</sup>	2358	5469	30
T <sub>10</sub> : 100% RDF+10t FYM ha <sup>-1</sup>	2416	5493	31
S. Em. (±)	82.96	190.37	1
C. D. (P=0.05)	242	545	NS

### 3.4 Nutrient uptake

The data on total nutrient uptake (N, P and K uptake) of sorghum is presented in Fig. 2. Total N (168 kg ha<sup>-1</sup>), P (17.5 kg ha<sup>-1</sup>) and K (98 kg ha<sup>-1</sup>) uptake of sorghum were recorded significantly highest from the treatment 100% RDF+10t FYM ha<sup>-1</sup> (T<sub>10</sub>) and all the values are lowest in T<sub>1</sub> (Control).



**Fig. 2: Effect of integrated nutrient management on total nutrient uptake of rabi sorghum.**

Whereas, total N uptake showed that T<sub>10</sub> is statistically at par with T<sub>9</sub>. But total P uptake showed that T<sub>9</sub>, T<sub>8</sub> and T<sub>4</sub> were statistically significant with highest recorded treatment T<sub>10</sub>. Though fertilizer supplies plant nutrient quickly, but judicious and combined application of both fertilizer and FYM might be improved soil microbial activity, which helped in better root proliferation as

well as increased different plant nutrient availability. This might be the reason for higher biomass production and nutrient uptake in 100%RDF+10t FYM ha<sup>-1</sup>. Similar results also found by Tomar *et al.*, (2017).

### 3.5 Economics

Cost of cultivation, gross return, net return and benefit: cost ratio of sorghum for different treatments has been calculated and presented in Table 4. The cost of cultivation (Rs. 37665/- ha<sup>-1</sup>) and gross return (Rs. 85108/- ha<sup>-1</sup>) of sorghum were highest in the treatment 100%RDF+10t FYM ha<sup>-1</sup> (T<sub>10</sub>) due to higher application of fertilizer and FYM. Moreover, in terms of benefit-cost ratio, the highest (1.53) results were obtained with the application of 100% RDF and in terms of net returns the highest (Rs. 48010/- ha<sup>-1</sup>) value was obtained with the application of 100%RDF+ 5t FYM ha<sup>-1</sup>. Application of higher quantity of FYM increased cost of cultivation and decreased benefit: cost ratio in treatments where RDF and FYM applied combinedly. But due to high yield and 5t less FYM application resulted maximum net return from the treatment 100%RDF+ 5t FYM ha<sup>-1</sup> (T<sub>9</sub>). In line with the above result Tomar *et al.* (2017) reported that application of 75%NPK + 5t FYM ha<sup>-1</sup> in maize recorded higher net return (Rs. 29634/- ha<sup>-1</sup>) and lower B:C ratio (2.57) as compared to the treatment 75%NPK, Rs. 27789/- ha<sup>-1</sup> and 2.62 respectively in 2011.

**Table 4. Effect of integrated nutrient management on economics of *rabi* sorghum.**

Treatments	Economics			
	Cost of cultivation (Rs. ha <sup>-1</sup> )	Gross return (Rs. ha <sup>-1</sup> )	Net return (Rs. ha <sup>-1</sup> )	B:C ratio
T <sub>1</sub> : Control	23755	35711	11956	0.50
T <sub>2</sub> : 50% RDF	28560	68653	40093	1.40
T <sub>3</sub> : 75% RDF	29915	74364	44449	1.49
T <sub>4</sub> : 100% RDF	31265	78991	47726	1.53
T <sub>5</sub> : 50%RDF+ 5t FYM ha <sup>-1</sup>	32460	73587	41127	1.27
T <sub>6</sub> : 50%RDF+10t FYM ha <sup>-1</sup>	34960	76654	41694	1.19
T <sub>7</sub> : 75%RDF+5t FYM ha <sup>-1</sup>	33815	79392	45577	1.35
T <sub>8</sub> : 75%RDF+ 10t FYM ha <sup>-1</sup>	36315	79728	43413	1.20
T <sub>9</sub> : 100%RDF+ 5t FYM ha <sup>-1</sup>	35165	83175	48010	1.37
T <sub>10</sub> : 100%RDF+10t FYM ha <sup>-1</sup>	37665	85108	47443	1.26

### 4. Conclusion

Integrated Nutrient Management plays a crucial role in well growth and development of sorghum plants. By incorporating the recommended doses of fertilizers and farm yard manure, farmers can promote balanced nutrition, enhance soil fertility, and improve overall

plant growth. However, it is important to consider other agronomic practices and environmental factors that influence plant growth for successful sorghum production. It should be emphasised that although utilising simply 100% RDF of inorganic fertilisers would improve benefit-cost ratio, but considering the net return, soil health, environmental well-being and sustainability taking into account the integrated strategy of applying 100% RDF+5t FYM ha<sup>-1</sup> would be advantageous.

## References

1. Afzal M, Ahamad A, Ahamad AH (2011). Effect of nitrogen on growth and yield of Sorghum forage under three cutting system. *Cercetari Agronomice in Moldova*. **45**(4):152.
2. Agricultural statistics at a glance. 2021. Department of Agriculture, Cooperation & Farmers Welfare. Directorate of Economics & Statistics, Government of India. [https://eands.dacnet.nic.in/PDF/Agricultural%20Statistics%20at%20a%20Glance%20-%202020%20\(English%20version\).pdf](https://eands.dacnet.nic.in/PDF/Agricultural%20Statistics%20at%20a%20Glance%20-%202020%20(English%20version).pdf) (Accessed on 2<sup>nd</sup> November 2022).
3. FAOSTAT. 2022. Crop Statistics. <http://www.fao.org/faostat/en/#data/QCL> (Accessed on 2nd November 2022).
4. Five Decades of Odisha Agriculture Statistics. 2020. Directorate of agriculture and food production, Odisha. <http://agriodisha.nic.in/content/pdf/Five%20Decades%20of%20Odisha%20Agriculture%20Statistics.pdf> (Accessed on 2nd September 2021).
5. Ghosh PK, Bandyopadhyay, Tripathi AK, Hati KM, Mandal KG, Mishra AK (2003). Effect of integrated management of farmyard manure, phosphocompost, poultry manure, and inorganic fertilizer for rain-fed sorghum (*sorghum bicolor*) in vertisol of central India. *Indian J. Agron.* **48**: 48-52.
6. Jat MK, Purohit HS, Singh B, Garhwal RS, Choudhary M (2013). Effect of integrated nutrient management on yield and nutrient uptake in sorghum (*Sorghum bicolor*). *Indian J. Agron.* **58**(4): 543-547.
7. Priya S, Kaushik MK, Sharma SK, Priyanka K (2014). Impact of integrated nutrient management on growth and productivity of hybrid maize (*Zea mays* L.). *Ann Biol.* **30**(1): 106-108.

8. Raja W, Khanday BA, Haq A (2010). Integrated nutrient management in rice (*Oryza sativa* L.) *Res. J. Agric. Sci.* **1**:150-152
9. Sabhad HV, Sakarvadia HL, Hirpara DV, Jadeja AS, Vekaria LC (2020). Influence of cadmium and FYM on growth, yield and quality of fodder sorghum. *Int. J. Conserv. Sci.* **8**(6): 546-549.
10. Sarkar AK (2005). Managing natural resources for increasing agricultural production in eastern India. *J. Indian Soc. soil Sci.* **53**: 435-437
11. Sharma A, Singh H, Nanwal RK (2007). Effect (*Triticum aestivum*) under limited and adequate irrigation supplies. *Indian J. Agron.* **52**: 120-123.
12. Sher A, Adnan M, Sattar A, Ul-Allah S, Ijaz M, Hassan MU, El Askary A (2022). Combined application of organic and inorganic amendments improved the yield and nutritional quality of forage sorghum. *Agron.* **12**(4): 896.
13. Shinde GG, Deshpande PM, Lomte MH, Deshmukh AS, Algave BN (2001). Yield maximization of irrigated rabi sorghum. *International Sorghum and Millets Newsletter.* **42**: 31-34.
14. Tomar SS, Singh A, Dwivedi A, Sharma R, Naresh RK, Kumar V, Singh BP (2017). Effect of integrated nutrient management for sustainable production system of maize (*Zea mays* L.) in indo-gangetic plain zone of India. *Int. J. Chem. Stud.* **5**(2), 310-316.