

Effect of Sulphur and Vermicompost on Growth Parameters, Yield Attributes and Yield of Chickpea

Abstract

A field experiment was conducted during *Rabi* season of 2022-2023 at Rajoula Agriculture farm, of Mahatma Gandhi Chitrakoot Gramodaya Vishwavidyalaya Chitrakoot, Satna (M.P.). The present experiment having 4 levels of sulphur and 3 levels of vermicompost of 9 treatment combination replicated thrice in factorial randomized block design. Chickpea variety GNG-1958 (Marudhar) was grown with recommended agronomic practices. On the basis of the results emanated from present investigation, it could be concluded that application of graded dose of sulphur significantly increases growth parameters (plant height, number of branches and number of nodule plant⁻¹), yield attributes (number of pod plant⁻¹, number of seed plant⁻¹ and seed index) and seed yield. All the growth and yield parameters were significantly higher S @25 kg ha⁻¹ as compared to S @20 kg ha⁻¹ and S @15 kg ha⁻¹. Similarly different levels of vermicompost significantly increases growth parameters (plant height, number of branches and number of nodule plant⁻¹), yield attributes (number of pod plant⁻¹, number of seed plant⁻¹ and seed index) and seed yield. All the growth and yield parameters were significantly higher V @5t ha⁻¹ as compared to V @2.5 t ha⁻¹ and V @0t ha⁻¹. The combination of S @25 kg ha⁻¹ and V @5t ha⁻¹ were found best in terms of growth and yield.

Comment [U1]: Rephrase

Keywords: Chickpea, Growth Parameters, Vermicompost and Yield

1. Introduction

Pulses play a pivotal role and occupy a unique position in Indian agriculture by virtue of their inherent capacity to grow on marginal lands. It is an easily available source of protein in the rural heart of India. Pulses provide significant nutritional and health benefits and are known to reduce several non-communicable diseases such as colon cancer and cardio-vascular diseases (Jukanti *et al.* 2012). India is the largest producer and consumer of pulses in the world. Major pulses grown in India include chickpea, pigeon pea, lentil, urd bean, mung bean, pea, lablab bean, moth bean, horse bean. It provides protein rich diet to the vegetarian of the Indian and complements the staple cereals in the diets with proteins, essential amino acids, vitamins and minerals (Pingoliya *et al.* 2013).

In India pulses are grown in 30.37 Mha of area with an annual production of 26.96 MT and productivity 888 kg ha⁻¹ and Gram production 136.13 lakh tonnes (record). (**According to PIB 2023**). In M.P. pulses are grown in 21.60 lakh ha of area with an annual production of 32.14 lakh tons and productivity 1488.0 kg ha⁻¹. (**Ministry of Agri. & FW 2021**).

They contain 22- 24 % protein, which is almost twice the protein in wheat & thrice that of rice (**Shukla et al. 2013**) and carbohydrate (61.51 %), fat (4.5 %) and relatively free from anti nutritional factors(**Saxena, 1990**). Chickpea is rich in protein content (20.47 g/100g), carbohydrate (62.95 g/100g), fibre (12.2 g/100g), phosphorous (252 mg/100g), high amount of minerals such as calcium (57 mg/100g), magnesium (79 mg/100g), iron (4.31 mg/100g) and zinc (15 mg/100g), low in fat content and most of it is polyunsaturated (**Wallace et al. 2016**).It is originated in south eastern turkey (**Redden et al. 2007**). Chickpea as a legume crop plays a significant role in improving soil fertility by fixing the atmospheric nitrogen (**Balai et al. 2017**).

Sulphur is a secondary nutrient and plays a vital role in plant metabolism as the main constituent of the sulphur containing amino acids (methionine and cysteine), Vitamin C (Glutathione, biotine and thiamine), lipoic acid and acetyl CO-A. In addition to these functions, ferro-sulphur proteins play an important role in nitrogen fixation and electron movement in photosynthesis (**Kadioglu, 2004**). Sulphur has positive effect on root growth in plants and this elements also help in the nodule formation in legumes crops. It is also associated with the aromatic compounds. Sulphur, in chickpea, mainly influences the protein content. Sulphur helps towards conversion of nitrogen into protein in pulse crops. Sulphur also improves the S containing amino acid in crop and thus enhances the protein content (**Das et al., 2016**).

Regular application of organics in amounts sufficient to meet the requirements of crops not only results in increasing crop yield but also improve the soil fertility and organic matter content (**Ramesh et al., 2008**).Continuous use of inorganic fertilizers has brought loss of vital soil fauna and flora. Organic production systems maintained and improved the soil health through stimulating the activity of soil organisms and organic manures are also helpful in alleviating the increasing incidence or deficiency of secondary and micronutrients and is capable of sustaining crop productivity. Organic manures modify the soil physical behavior and increases the efficiency of applied nutrients (**Pandey et al.,2007**). Organic manures not only supply a higher amount of different nutrient elements but also contains beneficial microbes like nitrogen

Comment [U2]: Add references
https://ijrbat.in/upload_papers/01102017084900Ddocumnt99.pdf

fixing bacteria, mycorrhizae and growth promoting substances for betterment of crops (**Barik et al., 2006**).

Vermicompost is usually a finely divided peat-like material with excellent structure, porosity, aeration, drainage and moisture-holding capacity (**Ismail 2005; Edwards et al., 2011**). It plays a vital role in dictating the biochemical cycles as it supports the growth and activities of soil micro flora. It enhances the colonization of *Mycorrhizae*, *Rhizobium*, *Azotobacter* and *Azospirillum* which in turn improve the nitrogen (N) as well as phosphorus (P_2O_5) supply and other micronutrients (Zn, Fe, Cu, Mn) besides imparting the resistance to plant against various soil borne diseases and insect pest attack. It enhances the root growth due to better soil physico-chemical properties (soil structure, porosity, less bulk density, organic matter, water holding capacity and cation exchange capacity (CEC)).

Comment [U3]: Add Reference
https://ijrbat.in/upload_papers/0206201701103029.pdf

2. Materials and Methods

2.1 Experimental Site

The experiment was carried out at Rajaula Agriculture farm, Mahatma Gandhi Chitrakoot Gramodaya Vishwavidyalaya Chitrakoot, Satna (M.P.) which lies in the semi-arid and sub-tropical region of Madhya Pradesh between 25.148° North latitude and 80.855° East longitude. The altitude of town is about 190-210 meter above mean sea level.

2.2 Edaphic Condition

The soil was moist, well drained with uniform plane topography. The soil of the experimental field was alluvial in origin, sandy loam in texture and slightly alkaline in reaction having pH 7.28 (1:2.5 soil: water suspension method given by **Jackson, 1973**), low in organic carbon percentage in soil is 0.24 per cent (Walkley and Black's rapid titration method given by **Walkley and Black, 1934**), low in available nitrogen 98.00 kg ha⁻¹ (Alkaline permanganate method given by **Subbiah and Asija, 1956**), medium in available phosphorus as sodium bicarbonate-extractable P was 17.31 kg ha⁻¹ (Olsen's calorimetrically method, **Olsen et al., 1954**), high in available potassium was 219.98 kg ha⁻¹ (Flame photometer method given by **Hanwey and Heidel, 1952**) and low in available sulphur 25.41 kg ha⁻¹ (Turbidimetric method given by **Chesnin and Yien, 1950**).

2.3 Experimental details

Chart 1. The experiment was conducted with 4 levels of sulphur and 3 levels of vermicompost of 9 treatment combination as indicated below.

Phosphorous Levels	Symbol	Vermicompost Levels	Symbol
0 kg ha ⁻¹	S ₀	0 ton ha ⁻¹	V ₀
15 kg ha ⁻¹	S ₁	2.5 ton ha ⁻¹	V ₁
20 kg ha ⁻¹	S ₂	5.0ton ha ⁻¹	V ₂
25 kg ha ⁻¹	S ₃		

2.4 Fertilizer and manure application

FYM was applied @ 10qha⁻¹ as basal dose. After the layout of experimental plot, the fertilizers were weighed and applied in the plots and thoroughly mixed with soil. As per the experimental recommended doses of Nitrogen, Phosphorus and Potassium were applied to all the plots. Recommended dose of Nitrogen, Phosphorus and Potassium were applied through Urea, DAP and MOP (20:40:20 kg ha⁻¹). Sulphur was applied at the time of sowing as per treatment.

2.5 Seed and sowing:

The seed sowing was done on 20th Nov. 2022. The seed was sown in line after making a narrow furrow with the help of pointed wooden stick at different row spacing. The seeds were dropped in the furrow after mixture with fine dust of soil and then after seeds were covered with thin soil layer. The total quantity of seed was required @ 3.2 kgha⁻¹. The chickpea variety was “GNG-1958 (Marudhar)”.

2.6 Harvesting and Threshing

The crop was harvested on 16th march, 2023 when it reached to its physiological maturity i.e. when the leaves were turned yellow and more than 70% pod were full matured. Threshing of 17th March, 2023 plot wise produce was done manually. The seed weight was recorded after sun drying the seed for three days. The seed weight thus obtained were converted into quintals per hectare on the basis of net plot size.

2.9 Statistical Analysis: The data on various characters studied during the course of investigation were statistically analyzed for factorial randomized block design. Wherever treatment differences were significant (“F” test), critical differences were worked out at five per cent probability level. The data obtained during the study were analyzed statistically using the methods advocated by **Gomez and Gomez (1984)**.

3. Result and Discussion

3.1. Growth Parameters

Growth parameters of chickpea were significantly influenced by the use of different sulphur levels and vermicompost levels. Plant height (cm) was significantly higher S @25 kg ha⁻¹ as compared to S @20 kg ha⁻¹ and S @15 kg ha⁻¹. The mean plant height (cm) at S₀, S₁₅, S₂₀ and S₂₅ were 35.26, 37.26, 38.90 and 41.80 cm. The mean plant height (cm) at V₁, V₂ and V₃ were 37.55, 38.37 and 39.00 cm. The mean no. of branches at S₀, S₁₅, S₂₀ and S₂₅ were 22.86, 24.33, 25.66 and 26.93. No. of branches was significantly higher in V₃ @5 t ha⁻¹ as compared to V₂ @2.5 t ha⁻¹ and V₁ @0 t ha⁻¹. The mean no. of branches at V₁, V₂ and V₃ were 24.47, 24.95 and 25.42. No. of nodule per plant was significantly higher S @25 kg ha⁻¹ as compared to S @20 kg ha⁻¹ and S @15 kg ha⁻¹. The mean no. of nodule per plant at S₀, S₁₅, S₂₀ and S₂₅ were 10.56, 11.33, 12.23 and 13.23 respectively. No. of nodule per plant was significantly higher in V₃ @5 t ha⁻¹ as compared to V₂ @2.5 t ha⁻¹ and V₁ @0 t ha⁻¹. The mean no. of nodule per plant at V₁, V₂ and V₃ were 11.52, 11.85 and 12.15 respectively. These findings are further supported by **Mir et al. (2013)**, **Joshi et al. (2016)**, **Kumaret al. (2018)** and **Kumar et al. (2022)**.

Table no.-1: Effect of different treatment combinations on growth parameters of barley

Treatment	Plant height (cm)	No. of branches	No. of nodules plant ⁻¹
Sulphur levels			
S ₀	35.26	22.86	10.56
S ₁₅	37.26	24.33	11.33
S ₂₀	38.90	25.66	12.23
S ₂₅	41.80	26.93	13.23
S.E.m±	0.23	0.23	0.07
C.D. (P= 0.05)	0.70	0.67	0.22

Vermicompost levels			
V ₁	37.55	24.47	11.52
V ₂	38.37	24.95	11.85
V ₃	39.00	25.42	12.15
S.E.m±	0.20	0.19	0.06
C.D. (P= 0.05)	0.60	0.58	0.19

Comment [U4]: Add

3.2. Yield Components

Yield attributing characters such as number of pod plant⁻¹, number of seed plant⁻¹ and seed index (g) were significantly influenced by the use of different sulphur levels and vermicompost levels. Number of pod plant⁻¹, number of seed plant⁻¹ and seed index were significantly higher S @25 kg ha⁻¹ as compared to S @20 kg ha⁻¹ and S @15 kg ha⁻¹. The mean number of pod plant⁻¹ at S₀, S₁₅, S₂₀ and S₂₅ were 34.03, 41.43, 50.40 and 58.56 respectively. The mean number of seed plant⁻¹ at S₀, S₁₅, S₂₀ and S₂₅ were 33.43, 43.26, 52.66 and 61.76. The mean seed index at S₀, S₁₅, S₂₀ and S₂₅ were 20.58, 23.36, 24.97 and 26.77 g respectively.

Number of pod plant⁻¹, number of seed plant⁻¹ and seed index (g) were significantly higher in V₃@5.0 t ha⁻¹ as compared to V₂@2.5 t ha⁻¹ and V₁@0 kg ha⁻¹. The mean number of pod plant⁻¹ at V₁, V₂ and V₃ is 43.35, 46.22 and 48.75 respectively. The mean number of seed plant⁻¹ at V₁, V₂ and V₃ were 44.65, 47.55 and 51.15 respectively. The mean seed index at V₁, V₂ and V₃ were 23.06, 23.89 and 24.81 gm respectively. These findings are further supported by Islam *et al.* (2011), and RamandKatiyar (2013)

Table no.-2: Effect of different treatment combinations on yield attributes of barley

Treatments	No. of pod plant ⁻¹	No. of seeds plant ⁻¹	Seed index
Sulphur levels			
S ₀	34.03	33.43	20.58
S ₁₅	41.43	43.26	23.36
S ₂₀	50.40	52.66	24.97
S ₂₅	58.56	61.76	26.77

S.E.m±	0.43	0.39	0.15
C.D. (P= 0.05)	1.28	1.15	0.46
Vermicompost levels			
V₁	43.35	44.65	23.06
V₂	46.22	47.55	23.89
V₃	48.75	51.15	24.81
S.E.m±	0.37	0.34	0.13
C.D. (P= 0.05)	1.12	1.00	0.40

Comment [U5]: Add Two annova

3.3 Seed yield

Seed yield ($q\ ha^{-1}$) was significantly influenced by the use of different sulphur levels and vermicompost levels at harvest stage. Seed yield ($q\ ha^{-1}$) at harvest stage was significantly higher at S_{25} as compared to S_{20} and S_{15} . The mean seed yield ($q\ ha^{-1}$) at S_0 , S_{15} , S_{20} and S_{25} were 13.33, 14.80, 15.80 and 16.96 $q\ ha^{-1}$ respectively at harvest stage. Seed yield ($q\ ha^{-1}$) at harvest stage was significantly higher in $V_3@5.0\ t\ ha^{-1}$ as compared to $V_2@2.5\ t\ ha^{-1}$ and $V_1@0\ t\ ha^{-1}$. The mean seed yield ($q\ ha^{-1}$) at V_1 , V_2 and V_3 were 14.75, 15.27 and 15.65 $q\ ha^{-1}$ respectively at harvest stage. These findings are further supported by the findings of Srinivasulu *et al.* (2015), Singh *et al.* (2015), Bera and Ghosh (2017) and Makolet *et al.* (2020).

Table no.-3: Effect of different treatment combinations on yields of barley

Treatments	Seed yield ($q\ ha^{-1}$)
Sulphur levels	
S₀	13.33
S₁₅	14.80
S₂₀	15.80
S₂₅	16.96
S.E.m±	0.13
C.D. (P= 0.05)	0.39

Vermicompost levels	
V ₁	14.75
V ₂	15.27
V ₃	15.65
S.E.m±	0.11
C.D. (P= 0.05)	0.33

Comment [U6]: Add Two annova

5. Conclusion

The experimental results indicated that superiority in regard to growth parameters, yield components and productivity parameters viz, grain yield ($q\ ha^{-1}$) with the use of treatment combination S @25 kg ha^{-1} and V @5t ha^{-1} gave in soil ensure highest growth parameters, yield components and productivity, of barley crop as comparison to all the treatments.

6. References

- Balai, K., Jajoria, M., Verma, R., Deewan, P. and Bairwa, S.K. (2017). Nutrient content, uptake, quality of chickpea and fertility status of soil as influenced by fertilization of Phosphorus and Zinc. *Journal of Pharmacognosy and Phytochemistry*.6 (1):392- 398.
- Barik, A. K.; Arindam, Das; Giri, A. K. and Chattopadhyaya G. N., (2006). Effect of integrated plant nutrient management on growth, yield and production economics of wet season rice. *Indian J. Agric. Sci.*, 76(1): 657-660.
- Bera, M and G.K. Ghosh, (2015). Efficacy of sulphur sources on green gram(*Vigna radiata* L.) in red and lateritic soil of west Bengal. *Intel. J. of Plant, Animal and Environ. Sci.*,5(2): 2231-4490.
- Edwards, C.A.; Subler, S. and Arancon, N. (2011). Quality criteria for vermicompost. In: Edwards CA, Arancon NQ, Sherman RL (eds.) Vermiculture technology: earthworms, organic waste and environmental management. *CRC Press*, Boca Raton, pp 287–301
- Gomez, K. A., & Gomez, A. A. (1984). Statistical procedures for agricultural research. *John wiley & sons*.

- Hanway, J.J; and Heidel, H. (1952).** Soil analysis methods as used in Iowa State College, Soil Testing Laboratory. *Iowa Agriculture* **54**: 1-31.
- Islam, M., S. Mohsan, S. Ali, R. Khalid, F.U. Hassan, A. Mahmood and A.Subhani, (2011).** Growth, nitrogen fixation and nutrient uptake by chickpea (*Cicer arietinum*) in response to phosphorus and sulfur application under rainfed conditions in Pakistan. *Intl. J. Agric. Biol.*, **13**:725-730.
- Ismail, S.A. (2005).** The earthworm book. Other India Press, Mapusa, pp 101.
- Jackson, M.L. (1973).** Soil chemical analysis. Prentice Hall of India Pvt. Ltd, New Delhi.
- Joshi, D., Gediya, K. M., Patel, J. S., Birari, M. M and S. Gupta (2016).** Effect of organic manures on growth and yield of summer cowpea (*Vigna unguiculata* (L.) Walp.) under middle Gujrat conditions. *Agricultural Science Digest*, **36**(2): 134-137.
- Jukanti, A.K., Gaur, P.M., Gowda, C.L.L. and Chibbar, R.N. (2012).** Nutritional quality and health benefits of chickpea (*Cicer arietinum* L.). *A Review British J. Nutr.* **108**: S11-S26.
- Kadioglu, A., (2004).** *Plant Physiology, Practice Guide*. Trabzon, Turkey.
- Kumar H., R. Singh, D. D. Yadav, M. Saquib, V. P. Chahal, R. Yadav and O. S. Yadav (2018).** Effect of integrated nutrient management (INM) on productivity and profitability of chickpea (*Cicer arietinum* L.). *International Journal of Chemical Studies*, **6**(6): 1672-1674.
- Kumar, B. J., & Mehera, B. (2022).** Effect of potassium and sulphur on growth and yield of chickpea (*Cicer arietinum* L.). *The Pharma Innovation Journal* 2022; **11**(4): 2139-2143.
- Makol, F. H., Gandahi, A. W., Memon, A. H., Jatoi, S. H., Abbasi, J. A., & Buriro, I. A. (2020).** Effect of Sulphur Application on Growth and Yield of Chickpea (*Cicer arietinum* L.) Under Rice Chickpea Cropping System. *Journal of Applied Research in Plant Sciences*, **1**(1), 9-12.

- Mir, A.H., S.B. Lal, M. Salmani, M. Abid and I. Khan, (2013).** Growth, yield and nutrient content of blackgram (*Vignamungo*) as influenced by levels of phosphorus, sulphur and phosphorus solubilizing bacteria. *SAARC J. Agric.*, 11 (1):1-6.
- Olsen, S.R., Cole, C.V., Watanable, F. S. and Dean, L. A. (1954).** Estimation of available phosphorus in soil by extraction with sodium bicarbonate. *USDA, Cric.* 930:19- 23
- Pandey, N.; Verma, A. K.; Anurag and Tripathi, R. S. (2007).** Integrated nutrient management in transplanted hybrid rice (*Oryza sativa* L.). *Indian Journal of Agronomy* 52(1): 40–2.
- Pingoliya, K, K., Dotaniya, M, L., Mathur, A, K. (2013).** Role of phosphorus and iron in chickpea (*Cicer arietinum* L.). *Lap Lambert Academic Publisher, Germany.*
- Ram, S. and T.P.S. Katiyar, (2018).** Response of sulphur and zinc on yield, quality and nutrient uptake of summer mung bean (*Vigna radiata* L. Wilczek). *J. of Pharmacy and Phytochemistry*, SP1:3243-3245.
- Ramesh, P.; Panwar, N. R.; Singh, A. B. and Ramana, S. (2008).** Effect of organic manures on productivity, nutrient uptake and soil fertility of maize–linseed cropping system. *Indian Journal of Agricultural Sciences* 78 (4): 351–4.
- Redden, B., Furman, B, J., Upadhyaya, H, D., Pundir R, P, S., Gowda, C, L, L., Coyne, C., Enne King, D., (2007).** Biodiversity Management in Chickpea. In: Yadav, S, S., Redden R., Chen, W., Sharma, B., editors. *Chickpea Breeding & Management*. CABI, Wallingford, UK. Pp 355-368.
- Saxena, M, C. (1990).** Problems and potential of chickpea production in the nineties. In: *Chickpea in the nineties. Proc Second Int. Workshop on chickpea Imp*, 4-8th December, 1989, ICRISAT Centre, Patancheru, Andhra Pradesh, India, pp. 13-25.

- Shukla, M., Patel, R. H., Verma, G., Deewan, P., Dotaniya, M, L. (2013).** Effect of bio-organics and chemical fertilizers on growth and yield of chickpea (*Cicer arietinum* L.) under middle Gujrat condition. *Vegetos.* **26**(1): 183-187.
- Singh, S., Bawa, S. S., Singh, S., Sharma, S. C., & Sheoran, P. (2015).** Productivity, profitability and sustainability of rain-fed chickpea under inorganic and biofertilization in foothills of north-west Himalayas. *Archives of Agronomy and Soil Science*, **61**(8), 1151-1163.
- Srinivasulu, D.V., R.M. Solanki, C.R. Kumari, and M.V. Babu, (2015).** Nutrient uptake, yield and protein content of chickpea (*Cicer arietinum* L.) as influenced by irrigation and sulphur levels in medium black soils. *Inter. J. of Agri. Sci.*, **11**:54-58.
- Subbiah, B.V. and Asija, C.L. (1956).** A rapid procedure for the estimation of available N in Soil. *Curr. Sci.* **25**:259-260.
- Walkley, A. and Black, C. S.A. (1934).** Old piper, S.S. soil and plant analysis. *Soil Sci.* **37**:29-38.
- Wallace, T. C., Murray, R., Kathleen, M. and Zelman, K. (2016).** The nutritional value and health benefits of chickpeas and humus. *Nutrients.* **8**(12): 766.