

Top Edible Wild Plants of Eastern Mediterranean Region. Part II: Anti-inflammatory Activity

Abstract:

In the first part of this series of review articles, we presented the anticancer activities of the most important wild edible plants of eastern Mediterranean region, which we named as the “Deca-plants” (D-P) Why it denomination?. In this review article, we will present the anti-inflammatory activities of these very important plants. After a brief introduction, anti-inflammatory activities of the D-P will be introduced in both traditional medicine of the peoples of this region, and in published scientific literature. Methods of use will be presented in the first part, and methods of testing and results, for modern research. In both cases, some of the D-Phave notable anti-inflammatory activities. In the extensive discussion section of this article, a major focus will be presented for anti-inflammatory activities of natural products contained in these plants, and some comparisons will be made with other plants. At this point, it is worth mentioning that the D-P have notably different number of anti-inflammatory activity reports, where on the top of them *Foeniculum vulgare* can be found. Contrary to that and to the best of our knowledge, *Cyclamen persicum* was not published for anti-inflammatory activity, even though some of its secondary metabolites are very well known for having this activity. Finally, the conclusions of this article will be accompanied with recommendations for future research. It was missing to refer to the method and tools used for the selection, organization, systematization and analysis of the bibliographic references.

Keywords: anti-inflammatory, medicinal plants, traditional medicine, plant extracts, essential oils, natural products, norisoprenoids, isothiocyanates

1. Introduction

Inflammation is a natural response of the immune system to damage triggered by various factors such as pathogens, cell damage and toxic materials [1]. This response, inflammation, can occur with many mechanisms in all body parts, especially major organs: heart, liver, pancreas, lungs, kidney, intestinal track, reproductive system and the brain.

Inflammation involved in the vast majority of sicknesses [2,3], and the treatments of these health disorders start with the treatment of inflammations that resulted them. **Figure 1** summarizes the inflammation-disease relationships for major chronic diseases.

Figure 1. Inflammation, which? and major chronic diseases. The caption of this figure is reduced only to diseases.

Until today (August 2023), 22 synthetic, nonsteroidal anti-inflammatory drugs (NSAID's) were approved by the USA Food and Drug Agency (FDA) [4]. Even though most of them have relatively high efficiency, their side and adverse effects of most of them are also well known, and some are relatively severe [5].

Since ancient times, humans tried to use their food not only as nutritional source, but also to treat health disorder, and in this sense, medicinal plants have very important role. This approach is increasingly adopted by modern medicine for treatment of almost all health ailments, and inflammation is among the top of these [6]. So, since the Mediterranean diet is one of the healthiest on Earth [7], recent modern studies are approving the capacity of this diet in combating chronic diseases and inflammation [8].

2. Ethnomedicinal Anti-inflammatory Activities of the D-P

In the first part of this article series which presented the anticancer activity of the D-P, we extensively presented their ethnobotanical uses, especially as nutritional source [9]. So, in this article only the ethnomedicinal anti-inflammatory activity.

Interestingly, the D-P were rarely reported to have notable anti-inflammatory activity in the ethnomedicine of the reviewed region. O. Said and his colleagues reported traditional medicine use of *Cyclamen persicum* to treat skin inflammations; *Majorana syriaca*, *Micromeria fruticosa* and *Salvia fruticosa* against intestinal inflammation [10].

In a follow-up review article, H. Azaizeh and his colleagues reported ethnomedicinal treatment of bacterial infections with *Cichorium pumilum* and *Salvia fruticosa* for intestinal inflammation [11]. In the same year of 2006, E. Lev reported in a review article that *Foeniculum vulgare* and *Salvia fruticosa* are used to treat intestinal diseases, but he did not

clearly indicate inflammations [12]. He also mentioned *Sinapis alba* for treatment of infections.

Even though *Gundeliatournefortii* was not mentioned by the previous groups for ethnomedicinal use as anti-inflammatory treatment, M. Asadi-Samani and his colleagues indicated that this plant is used for this purpose, without specifying the type of inflammation or the treatment method [13]. Finally, S. Oran and D. Al-Eisawi reported the use of *Cichorium pumilum* to treat eczema [14].

3. Anti-inflammatory Activities of the Deca-plants and Their Natural Products

The results of modern research of the anti-inflammatory activities of the D-P were published in a few dozens of articles. As we mentioned earlier, the number of these published studies notably differ from one plant to another. But these differences relate also to the natural products responsible for these activities, where in some of the D-P they were isolated and characterized and in others they were not. Anti-inflammatory activity of the D-P is presented in **Table 1**.

Table 1. Published Anti-inflammatory Activities of the D-P in Eastern Mediterranean region.

Testing Method, Results and Reference/s
<p style="text-align: center;"><i>Arum palaestinum</i></p> <p>Leaves were extracted with 96% aqueous ethanol, and extract was used to treat carrageenan-induced paw edema in albino rats. Extract was given orally (200-300 mg/kg b.wt.) and the effect was very weak. [15]</p> <p>80% Aqueous methanolic leaves extract was prepared and used to treat UVB-irradiated (2 J/cm²) human cultured skin. Concentration of IL-1α was measured and the results show high activity. [16]</p> <p>Inflammation was induced by LPS in human monocytic cell line (THP-1)-derived macrophages, then they were treated with leaves aqueous ethanolic (96% v/v) extract. The treatment significantly suppressed the production of pro-inflammatory biomarkers and increased the production of anti-inflammatory biomarkers. [17]</p>
<p style="text-align: center;"><i>Cichorium pumilum</i></p> <p>Roots were extracted with 95% aqueous ethanol, and extract was fractionized with ethyl acetate (EA). LPS-induced inflammation in rats was treated with the EA fraction (100 mg/kg per day, oral gavage, 7 days), showing high activity. Authors proposed MAPK signaling as mechanism of action, and lactucin (Figure 2), isolated from EA fraction, as active natural product. [18]</p> <p>Whole plant extracted with 70% aqueous ethanol, and extract was fractionized to obtain phenolic acids-rich fraction and three pure acids (Figure 3). Injury in mice was induced by injection of methotrexate (Figure 3), and animals were treated with the crude extract (100 mg/kg per day, 10 days) and the acids-rich fraction (10 mg/kg per day, 10 days). All five extracts had significant activity, but ferulic acid was most active. [19]</p> <p>Leaves were extracted with 80% aqueous ethanol, and extract was found active against LPS-induced inflammation in THP-1-derived macrophage M0 cells. [20]</p>
<p style="text-align: center;"><i>Cyclamen persicum</i></p> <p>No published studies (see Discussion).</p>
<p style="text-align: center;"><i>Foeniculum vulgare</i></p> <p>Dry fruits were extracted with 80% aqueous methanol. Inflammation was induced in mice and rats by carrageenan (paw edema), arachidonic acid (ear edema) and formaldehyde (arthritis). Extract was orally administered (200 mg/kg per day, 10 days), with 0.5%</p>

sodium carboxymethyl cellulose. In the three tests extract was notably active, and best results were recorded after 3 h. [21]

Rats and mice with carrageenan-induced paw edema were treated with fruits essential oil (EO), with doses of 0.05, 0.1 and 0.2 mL/kg, intraperitoneal injection. Comparing with control (Etodolac), EO was most efficient in 0.05 and 0.2 mL/kg. [22]

Ear edema was induced in mice by arachidonic acid. Seeds were separately extracted with 80% aqueous ethanol and water, and both extracts were analyzed for active compounds. Activity was measured by 5-lipoxygenase release inhibition. Both extracts and some components (γ -terpinene, fenchone, *trans*-anethole, **Figure 4**) were notably active. [23]

Inflammation in rats was induced by injection of cyclosporine. Then, animals were fed with seeds (17 g/kg per day, 45 days), and various inflammatory biomarkers were tested. Treatment resulted significant effect compared with control. [24]

Rats and mice were subjected to inflammation induction by carrageenan (paw edema), acetic acid (writhing) and formaldehyde (paw licking). The three cases were treated with seeds ethanolic extract, resulting positive effect in all of them. [25]

LPS-stimulated inflammation in RAW 264.7 macrophage cells, and they were treated with 80% aqueous methanol fruits extract (100 μ g/mL). Several inflammation biomarkers were tested resulting significant improvement. [26]

5-Lipoxygenase (5-LOX) assay was used to test the activity of fruits commercial EO, resulting significant inhibition. [27]

Fruits were extracted with 70% aqueous ethanol and fractionized with several solvents. Dichloromethane fraction treated LPS-induced inflammation in RAW 264.7 macrophage cells. Effect was observed by measuring several inflammation biomarkers. [28]

Inflammation in mice was induced by LPS and the animals were treated with fruits EO (250 μ L/kg), resulting improvement of several inflammation biomarkers. Authors propose ERK signaling (extracellular signal-regulated kinase) as a possible mechanism of action. [29]

Aqueous extract of seeds was prepared, and it was used in ointment (2% and 7%) to treat formaldehyde-induced inflammation in rats. After 16 days, the 7% ointment reduced inflamed area by 91%, while 2% preparation effect was 81% and standard (mupirocin) 92%. [30]

Fresh leaves 50% aqueous methanolic extract was active against inflammation in THP-1-derived macrophage cells. [31]

A solution (100 g in 400 mL of water) of dry aqueous seed extract, was administered (0.8 mL/kg per day, 4 days) to rat puppies with necrotizing enterocolitis (CO_2 (g) inhalation). Testing several biomarkers showed positive effect of the treatment (puppies were scarified). [32]

Dry fruits and leaves were extracted with 96% aqueous ethanol, and extract was orally administered to rats with carrageenan-induced inflammation: 87.5, 175 and 350 mg/kg. All doses showed positive effect. [33]

95% Aqueous ethanol seeds extract had significant effect against LPS-stimulated inflammation in RAW 264.7 macrophage cells. [34]

Fruits EO was prepared by using *n*-hexane as collecting solvent, and GC-MS analysis found estragole as major component (**Figure 5**). Inflammation in neutrophils that were isolated from human blood, was induced using the fMLF/CB superoxide generation (N-Formylmethionyl-leucyl-phenylalanine, **Figure 5**), and it was treated with EO. Positive effect was measured by several biomarkers, and authors propose blocking of Ca^{+2} from Gi-protein to MAPKs (mitogen-activated protein kinases) by estragole, as mechanism of action. [35]

Seeds were extracted with ethanol/ CO_2 and extract was active against carrageenan-

induced paw edema in rats. [36]

Seeds EO was active in 5-LOX inhibition assay. [37]

Plant waste was analyzed for active phytochemicals. Six phenolic acids were identified, and the active anti-inflammatory (inhibition of COX-1 and COX-2) compounds were kaempferol, isorhamnetin, and quercetin glucuronide (**Figure 6**). [38]

Aerial parts, including fruits, were separately extracted with *n*-hexane, methylene chloride, ethyl acetate and methanol. All extract showed positive effect on carrageenan-induced paw edema in rats, and methanol extract was most potent. In this study, comprehensive chemical compositions of extracts were determined. [39]

R-(+)-Limonene (**Figure 7**), a major ingredient of seeds EO, and fenchone (**Figure 4**), were commercially purchased and applied for wound healing in rats. Both compounds had anti-inflammatory activity by increase of collagen synthesis. [40]

Pure commercial *trans*-anethole (**Figure 4**) had anti-inflammatory activity against LPS-induced lung injury, by regulation of Th17/Treg (T cell helper 17, T cell regulatory) function. [41]

Pure coumarins (**Figure 7**) isolated from fruits, were active against LPS-induced inflammation in RAW 264.7 macrophages and in a TPA-stimulated mouse model of skin inflammation (TPA, 12-O-tetradecanoylphorbol-13-acetate). [42]

Gundeliatournefortii

Aerial parts were extracted with 95% aqueous ethanol, and extract was active against edema induced in mice by formaldehyde (paw) or xylene (ear). [43]

Inflammations was separately induced in rats by acetic acid, formaldehyde, and hot plate. These were treated with aerial parts EO, resulting positive effect in the three cases. Chemical composition of EO was determined, and major components are shown in **Figure 8**. [44]

Aerial parts were burned and mixed with milk to prepare an ointment. This was applied on second degree burns in rats, resulting significant decrease of burns. [45]

Aqueous extract of flowering buds was prepared and used to treat inflammation caused by implanted hepatocarcinoma cells in mice (reduction of F4/80 count). [46]

Majorana syriaca

Aerial parts EO was prepared and analyzed with GC-MS for chemical composition. Major compounds are shown in **Figure 9**. EO had activity against LPS-induced inflammation in RAW 364.7 macrophages, tested by inhibition of NO production. [47]

Three acids (**Figure 10**) known for their anti-inflammatory activity (LPS-induced inflammation in RAW 364.7 macrophages) were isolated and quantified by LC-MS, from the aerial parts ethyl acetate extract. [48]

Aerial parts 90% aqueous ethanol extract was prepared and found active against carrageenan-induced paw edema in rats. [49]

Methanolic leaves extract was suppressed the release of cytokine IL-6 but not cytokine IL-10 in human peripheral blood mononuclear cells. [50]

Aqueous and *n*-hexane leaves extracts were separately prepared. Extracts were used to treat (50 mg/kg per day, 10 days, intraperitoneal administration) mice were infected with *Schistosoma mansoni*. Both extracts had positive effect. [51]

Follow-up (ethanolic extract) of previous study using scanning electron microscopy. This study focused on *M. syriaca*. [52]

Leaves 70% aqueous ethanolic extract inhibited 5LOX, COX-1 and sPLA₂-Venzymes. [53]

Malva sylvestris

80% Aqueous methanolic leaves extract was prepared and used to treat UVB-irradiated (2 J/cm²) human cultured skin. Concentration of IL-1 α was measured and the results show

low activity. [16]

Flowers were extracted with diethyl ether, and 200 mg/kg of it were used to treat wounds in alloxan-induced diabetic rats. Treatment reduced wound area by collagen production and inhibition of inflammation. [54]

Follow up of previous study: flowers chloroform extract, same dose, healthy rats. Wound healing and anti-inflammatory activities were higher. [55]

Leaves were extracted with 96% aqueous ethanol, and extract was analyzed with LC-MS to obtain scopoletin, quercetin and malvidin-3,5-glucoside (**Figure 11**). The crude extract and the three compound were used to treat TPA-induced ear edema in mice. Extract (0.2 mg/ear) and malvidin-3,5-glucoside (0.05 μ mol/ear) were active. [56]

70% Aqueous ethanolic leaves extract was mixed (20%) with Orabase, and the resulting ointment treated wounds in rats. Wound area was decreased, and inflammation progress was inhibited, compared with control. [57]

Leaves were extracted with ethanol and the extract was fractionized with chloroform. Both extract and fraction had significant anti-inflammatory activity against *Aggregatibacter actinomycetemcomitans*-infected human oral cells. Activity was expressed by down regulation of inflammatory genes. [58]

A cream contains 1% flowers aqueous extract, had wound healing and inflammation preventive activities in mice. [59]

Aerial parts were separately extracted with water, ethanol, and *n*-hexane, and from the aqueous extract, a polysaccharide was isolated and characterized. Aqueous extract and polysaccharide had significant activity of preventing inflammation in acetic acid-induced ulcerative colitis in rats. [60]

Leaves ethanolic extract was prepared and fractionized with *n*-hexane, chloroform, ethyl acetate and water. Using three inflammation models (*in vitro* and *in vivo*, mice), it was found that only the aqueous fraction had notable activity. [61]

Leaves and flowers ethanolic extract was prepared and fractionized with *n*-hexane, chloroform, ethyl acetate and water. LPS-induced *in vitro* inflammation model was used to test activity of extract and fractions. Release of inflammatory mediators PGE₂, PGD₂, TXB₂, and PGF_{2 α} was measured, revealed activity mainly for extract, ethyl acetate and aqueous fractions. Authors propose inhibition of prostanoid production as the mechanism of action. [62]

Three extracts were prepared from leaves: ethanolic, 70% aqueous ethanolic and aqueous. Extracts were tested in postoperative peritoneal adhesion in rats, showing different, but positive effects for all of them. There was low influence on inflammatory biomarkers. [63]

Flowers were extracted with 70% aqueous ethanol, and the same solvent was used to separately extract flowers of *Carum carvi* and *Medicago sativa*. Each extract was separately tested against formaldehyde-induced paw edema in rats. All extracts were active (600 mg/kg) and the combination of the extracts was most potent. [64]

The 95% aqueous methanol leaves extract was prepared and used to treat ischemia-reperfusion injury in rats (*in vivo*) and in H9c2 cells (*in vitro*). [65]

Aerial parts were extracted with ethanol and fractionized with water. This fraction decreased the release of inflammatory biomarkers, especially IL-6 in HIV infected TZM-b1 cells. [66]

Aerial parts ethanolic extract was used to prepare a topical cream that treated atopic dermatitis in children. [67]

Fruits powder, aqueous and ethanolic extracts were used to treat carrageenan-induced paw edema in rats. All products were active, and the aqueous extracts (100 mg/kg) was most potent. [68]

Leaves ethanolic extract and flowers aqueous extract, were prepared; and ferulic acid (**Figure 3**) was isolated from extracts. The three products were tested against inflammation of wounds in mice, as well as in HaCaT cells. All products were active decreasing release of inflammatory biomarkers (IL-6, IL-1 β). [69]

Aerial parts ethanolic extract was used (10%) to prepare a cream for wound healing in humans. There was no significant activity. [70]

Aqueous and acetone extracts of aerial parts were prepared and tested for inhibition of bovine serum albumin denaturation. Aqueous extract was more active than acetone extract, but less than control, diclofenac sodium. [71]

Micromeria fruticosa

Aerial parts were extracted with 80% aqueous methanol, at 80 °C (boiling point 82.5 °C). The extract was tested in three concentrations (20, 40, 60 $\mu\text{g/mL}$) for inhibition of *in vitro* oxidation of 3,3',5,5'-tetramethylbenzidine by MPO/H₂O₂/Cl⁻ system (MPO, myeloperoxidase). The concentration of 60 $\mu\text{g/mL}$ was most active with 86% inhibition. [72]

Aqueous extract of aerial parts was prepared and tested for anti-inflammatory activity in four models: MPO inhibition, carrageenan-induced paw edema in mice, acetic acid-induced vascular permeability in mice, and indomethacin-induced ulcer in mice. Among three concentrations, 200 mg/kg was most potent. [73]

Salvia fruticosa

Aerial parts were separately extracted with ethanol, chloroform, 1-butanol, and water. Chloroform extract had the highest activity against carrageenan-induced paw edema in rats (25 mg/kg, orally). [74]

Methanolic extract of aerial parts was prepared and tested against LPS-induced inflammation, *in vitro* (RAW 264.7 cells) and *in vivo* (mice). In both cases, inhibition proinflammatory biomarkers production was measured: TNF- α , IL-6 and IL-1 β . [75]

Aerial parts were separately extracted with water, methanol, *n*-butanol, acetone, and chloroform. All extracts (including of two other species of *Salvia*) were tested against carrageenan-induced paw edema in rats. *n*-Butanol extract of *S. fruticosa* was most active. [76]

Methanolic extract of aerial parts was found active (500 mg/kg, intraperitoneal injection) against carrageenan-induced edema in mice. [77]

Aerial parts were extracted with 70% aqueous methanol, and the extract was used to treat AlCl₃ – induced Alzheimer's disease neuroinflammation in rats. Using 375 or 750 mg/kg, daily for three months, had notable effect of reducing the release of inflammatory biomarkers, including acetylcholinesterase (AChE). [78]

Roots and aerial parts were separately and successively extracted with chloroform, ethyl acetate, methanol, and 1-butanol. The methanolic extracts had the same effect against carrageenan-induced paw edema in mice. [79]

Nanovesicles were loaded (17-66 mg/mL) with leaves EO and were found active in soybean lipoxygenase inhibition assay. Major compounds in this EO are shown in **Figure 12**. [80]

Leaves and flowers were separately extracted with methanol (mercerization) and ultrasound-assisted extraction. Extract was found active against LPS-induced inflammation in RAW 264.7 cells. [81]

Sinapis alba

Mice diet that included 5% seeds, imiquimod-induced psoriasisiform inflammation on shaved back skin of the animals. [82]

Seeds aqueous extract was active against LPS-induced inflammation in RAW 264.7 cells. [83]

Seeds were extracted with 50% aqueous ethanol. Extract had positive effect on TPA or arachidonic acid or croton oil-induced ear edema in mice. [84]

Figure 2. Lactucin isolated from roots of *Cichorium pumilum* [18]



Figure 3. Active acids from *C.pumilum* against methotrexate injury [19]

Figure 4. Anti-inflammatory natural products from *Foeniculum vulgare* [23]

Figure 5. Estragole from *F. vulgare*, active against fMLF-induced inflammation[36]

Kaempferol, O₄

Figure 6. Anti-inflammatory natural products isolated from *F. vulgare* waste[38]

R-(+)-Limon



Figure 7. Anti-inflammatory pure natural products found in *F. vulgare*[40,42]

Figure 8. Major constituents of aerial parts essential oil of *Gundelia tournefortii*[44]



Figure 9. Major constituents of aerial parts essential oil of *Majorana syriaca*[47]

Figure 10. Anti-inflammatory acids from aerial parts ethyl acetate extract of *M.syriaca*[48]




Figure 11. Active compounds from *Malva sylvestris*[56]

Figure 12. Major constituents of leaves essential oil of *Salvia fruticosa* [80]

4. Discussion

The presentation of published modern research results of anti-inflammatory activity of the Deca-plants (D-P) revealed very interesting fact: while some plants were extensively published, others were limitedly published or not published at all (*Cyclamen persicum*). Why? Another interesting and contradictory reporting was presented about the anti-inflammatory activity of *Arum palaestinum*. Despite being aware of the different testing methods, the reported results of anti-inflammatory activity of the leaves extracts of *Arum palaestinum* according to references 15 (very weak) and 16 (very strong) are easily notable. The different extracting solvents: 96% aqueous ethanol and 80% aqueous methanol, respectively, does not provide possible explanation since literature reports concerning the efficiency of these solvents are quite contradictory [85-88].

The importance of anti-inflammation research and its findings consistency, arises from the understanding that we already mentioned in the **Introduction**, that inflammation is involved in almost all diseases. To make this statement even more valid, we will present some of the very important recent publications. I.G. Onyango and his colleagues reviewed the relationship between neuroinflammation and Alzheimer's disease: they present a comprehensive view of the causes of neuroinflammation and major known interventions [89]. A. Kurowska and her colleagues, focused on the importance of healthy diet in prevention of inflammatory processes that lead to neurological diseases [90]. This presentation connects directly to our interest in this article about D-P, and it has another advantage: it reviews unhealthy, pro-inflammatory foods and not only anti-inflammatory nutrition.

The recent research of A. García-Sánchez and his colleagues proved a direct relationship between inflammation and obesity and hypertension [91]. For example, in normal-weight human subjects of the research, the concentrations of pro-inflammatory cytokines were 206.4 pg/mL for TNF- α , and 63.9 pg/mL for IL-6. For overweight and obese subjects, concentrations were 195.6, 66.5 and 496.2, 217.0 pg/mL, respectively. Following the previous findings, D. Moriki and her colleagues found clear negative linkage between adherence of anti-inflammatory diet and atopic diseases prevalence, as shown in **Table 2** [92].

Table 2. Adherence to an anti-inflammatory diet and atopic diseases prevalence among Greek youngsters [92]

Adherence to an Anti-Inflammatory diet	Low	Moderate	High
Asthma symptoms in the past 12 months (%)	10.8	8.3	4.6
Allergic rhinitis symptoms in the past 12 months (%)	29.2	24.7	24.5
Allergic rash symptoms in the past 12 months (%)	8.9	9.5	8.5
Overweight/Obese (%)	33.0	31.6	32.6

To conclude this part, we will cite the comprehensive review article of F. Stumpf and her colleagues [93]. They presented the benefits of healthy diet as strong anti-inflammatory factor, and the unhealthy and pro-inflammatory nutrition, like in the review article of A. Kurowska and her colleagues [90]. But the article of F. Stumpf *et al.* also reviewed the role of malnutrition in inflammation.

Three studies have reported anti-inflammatory activity of *A. palaestinum* [15-17], but none of them linked this activity to specific natural products (NPs) contained in this plant. In a very detailed study, I.M. Abu-Reidah and his colleagues, have published the chemical composition of *A. palaestinum* [94]. Carefully studying the list of compounds that they presented, shows that there are at least two natural products with known and published anti-inflammatory activity: laurocapram (azone) and 7-methoxy coumarin [95,96], that their structures are shown in **Figure 13**.



Figure 13. NPs with reported anti-inflammatory activity contained in *A. palaestinum* [92,93]

In section 3 (Anti-inflammatory Activities of the Deca-plants and Their Natural Products), we have cited three studies of anti-inflammatory activity of *Cichorium pumilum* [18-20]. One of them however (19, E.E. Eltamany *et al.*), linked this activity with three phenolic acids (Figure 3). These acids are very widespread in the plant kingdom. But studying the chemical composition of *C. pumilum*, reveals the fact that it contains NPs family, found in some other plants: norisoprenoids [97]. Later studies have shown that two of these NP's, 4 and 5 in **Figure 14**, have notable anti-inflammatory activity [98,99].

Figure 14. Norisoprenoids contained in *C. pumilum* [97-99]

To the best of our knowledge, *Cyclamen persicum* was not published by modern research as having anti-inflammatory activity. Even though the D-P have very few mentions as having such activity in traditional medicine of the reviewed region, *C. persicum* is used to treat skin inflammations [10]. Moreover, and more important, analyzing the chemical composition of

this plant shows that it contains several compounds, in significant concentrations, with clear and notable anti-inflammatory activity. For example, M. Cornea-Cipcigan and her colleagues, analyzed leaves of flowers of a few *Cyclamen* species, including *C. persicum*. They published their findings in a comprehensive research article showing a high content of carotenoids [100]. These compounds are widely reported and reviewed for having anti-inflammatory activity [101].

Another specific and clearer example was the work of H. Ishizaka and his colleagues: they discovered that among the volatile compounds of *C. persicum*, the isomers *E,E*- and *Z,E*-farsenols are present [102], **Figure 15**. These compounds, especially the *E,E*-isomer, have well known anti-inflammatory activity [103,104].

Figure 15. Farsenols, anti-inflammatory NPs, found in *C. persicum* [102-104]

Inhibition of nitric oxide (NO) production is one of the major anti-inflammatory activities [105], and in this context, it was found that 95% aqueous ethanolic tubers extract inhibited the production of NO in LPS-induced inflammation in cancer cells [106]. Interestingly, the authors of this report do not indicate that this activity is anti-inflammatory, and this is the reason we did not mention it section 3.

As we have presented in Table 1 above, the anti-inflammatory activity of *Foeniculum vulgare* is well known and extensively published [21-42]. This clear and notable property is utilized in some indirect and advanced methods. A. Ammara and his colleagues prepared selenium nanoparticles (SeNP) using *F. vulgare* aqueous seeds extract that reduced sodium selenite [107]:



These SeNP had activity against collagenase type II-induced paw arthritis in mice.

Z.T. Gias and her colleagues tested the biological activities of a commercial formulation (PanchPhoron) that is composed of: *Foeniculum vulgare* (fennel), *Trigonella foenum-graecum* Linn (fenugreek), *Nigella sativa* (black cumin), *Cuminum cyminum* (cumin) and *Brassica nigra* (black mustard) [108]. They found that this formulation has three inflammation-related activities in mice: antinociceptive, anti-inflammatory and analgesic.

M.R. Loizzo and her colleagues reported that EO of the leaves of *Majorana syriaca* had activity against LPS-induced inflammation in RAW 364.7 macrophages, tested by inhibition of NO production [47]. They analyzed this EO by GC-MS and found that Carvacrol, thymol, p-cymene and 2-isopropyl-1-methoxy-4-methylbenzene (Figure 9) were the major constituents, but they did not link the anti-inflammatory activity to all or some of these compounds.

In terms of chemical composition of the EO of *M. syriaca* leaves, the report of M.R. Loizzo *et al.* is consistent with previous reports such as of S. Abu-Lafi and his colleagues [109], where carvacrol and thymol were major ingredients. Several publications have reported the notable anti-inflammatory activity of these compounds [110,111].

Malva sylvestris is one of the most studied, published, and reviewed plants [for example, 112-114]. In our presentation, we have cited 19 publications of the anti-inflammatory activity of this plant [16,54-71]. Phenolic compounds are among the top anti-inflammatory NPs contained in *M. sylvestris*, and malvidin-3,5-glucoside is the most active (Figure 11), [56]. These results are consistent with other published studies about the anti-inflammatory properties of malvidin and its derivatives [115]. But none of the 19 studies that we cited, indicated delphinidin, a phenolic compound contained in *M. sylvestris* [116], and possesses high anti-inflammatory activity [117].

In section three, we cited 8 publications about the anti-inflammatory activity of *Salvia fruticosa* [74-81], and in Figure 12 we presented the major components of *S. fruticosae* leaves EO, that has anti-inflammatory activity [80]. But this plant contains high concentrations of rosmarinic acid [118] (Figure 10), which also has strong anti-inflammatory activity [48,119,120].

Another phenolic acid contained in *S. fruticosae* leaves [118], with proven anti-inflammatory activity is salvianolic acid B [121], which also was not mentioned in all the articles we cited about this plant. The structure of this acid is shown in **Figure 16**.

Figure 16. Salvianolic acid B, anti-inflammatory agent found in *S. fruticosa* [118,121]

None of the three articles that we cited about the anti-inflammatory activity of *Sinapis alba* [82-84], mentioned, not to say presented, active anti-inflammatory NPs that this plant contains. The chemical composition of this plant was studied and published: seeds EO [122-124], seeds *n*-hexane extract [125], and a very recent, **comprehensive, and detailed** composition that was achieved by several extractions and fractionations [126].

Carefully studying the lists of compounds in the last four cited publications [122-126], reveals the fact that many of these NPs have known and published anti-inflammatory activity. Here we will present the most studied and published, and they are presented in **Table 3** and **Figure 17**.

Table 3. Selected Anti-Inflammatory Compounds in *Sinapis alba*

Compound Name	Testing Method/s [Reference]
Allyl isothiocyanate	LPS-induced neuroinflammation [127]
Benzyl isothiocyanate	LPS-induced inflammation in RAW 264.7 macrophages and TPA-induced ear enema in mice [128] Indomethacin-induced gastric injury [129]
Genipin	Croton oil-induced ear edema in mice [130]
Geniposide	Wound healing in diabetic rats [131]
Glucoraphanin	LPS-induced inflammation in THP-1 cells [132]
Sinapic acid	Carrageenan-induced paw edema in rats [133]
Sinapine	Fatty liver inflammation in mice [134]
Sulphoraphane	LPS-induced inflammation in RAW 264.7 cells and in mice [135] LPS-induced inflammation in human dendritic cells [136]




Figure 17. Selected anti-inflammatory NPs found in *S. alba* [127-136]

5. Selected Wild Edible Non-Deca-Plants with Anti-inflammatory activity

The plants, subject of this review article that we named as Deca-plants (D-P) are the most consumed edible wild plants in Eastern Mediterranean region. This last statement is not based on scientific reports, but rather personal knowledge of the author, and it is highly correct for the region between Eastern Mediterranean shores and the Jordan river.

But peoples of the smaller or the greater reviewed regions, eat dozens of other wild plants, obviously, as well as dozens of cultivated plants. Among the edible wild plants that are not included in the D-P (N-D-P), some compete with the D-P plants in the extent of consumption, such as *Malva nicaeensis*, that is eaten like *Malva sylvestris*, and in hilly areas even more. Some of these N-D-P, possess notable anti-inflammatory activity. In **Table 4** we present eight of these plants, the most common ways of their consumption, and selected reports of their anti-inflammatory activity.

Table 4. Selected Wild Edible Non-Deca-Plants with Reported Anti-Inflammatory Activity

Plant Name	Main Method/s of Consumption [Anti-inflammatory Report]
<i>Capparis spinosa</i>	Pickled buds [137]
<i>Malva nicaeensis</i>	Cooked or fresh in salads (like <i>M. sylvestris</i>) [138]
<i>Matricaria aurea</i>	Herbal drinks [139]
<i>Notobasis syriaca</i>	Peeled young stems eaten fresh, seeds as coffee substitute [140]
<i>Portulaca oleracea</i>	Fresh in salads, cooked or in omelets [141]
<i>Rumex acetosa</i>	Pastry filling (like <i>Spinacia oleracea</i> , Spinach) or fresh [142]
<i>Salvia hierosolymitana</i>	Leaves stuffed with rice (like stuffed grape leaves) [143]
<i>Urtica urens</i>	Young stems eaten fresh, in salads or cooked [144]

6. Conclusions

- 1) The top ten most edible wild plants (Deca-plants, D-P) of Eastern Mediterranean region have notable anti-inflammatory activities.
- 2) Some of the D-P were limitedly studied and/or published, and there is a need to investigate the anti-inflammatory activities of these plants.
- 3) It is highly important to find the natural products contained in the D-P and are responsible for the anti-inflammatory activities.
- 4) When these natural products are known, it is important to study the mechanisms of action of their anti-inflammatory activities.
- 5) It is important to study synergistic effects of active anti-inflammatory D-P products.

7. References

- 1) Chen, L., Deng, H., Cui, H., Fang, J., Zuo, Z., Deng, J., Li, Y., Wang, X., Zhao, L., Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget*.**2017**, *9*, 7204-7218. DOI: 10.18632/oncotarget.23208

- 2) Dhingra, A.K., Chopra, B., Inflammation as a Therapeutic Target for Various Deadly Disorders: A Review. *Curr. Drug Targets*. **2020**, *21*, 582-588. DOI:10.2174/1389450120666191204154115
- 3) Laveti, D., Kumar, M., Hemalatha, R., Sistla, R., Naidu, V.G., Talla, V., Verma, V., Kaur, N., Nagpal, R., Anti-inflammatory treatments for chronic diseases: a review. *Inflamm. Allergy Drug Targets*. **2013**, *12*, 349-361. DOI: 10.2174/18715281113129990053
- 4) Ghlichloo, I., Gerriets, V., Nonsteroidal Anti-inflammatory Drugs (NSAIDs). **2023**, May 1. In: StatPearls [Internet]. [[NIH National Library of Medicine](#)]
- 5) Jahnavi, K., Pavani Reddy, P., Vasudha, B., Narender, B., Non-steroidal anti-inflammatory drugs: an overview, *J. Drug Deliv. Ther.* **2019**, *9*, 442-448. DOI: 10.22270/jddt.v9i1-s.2287
- 6) Ricker, M.A., Haas, W.C., Anti-Inflammatory Diet in Clinical Practice: A Review. *Nutr. Clin. Pract.* **2017**, *32*, 318-325. DOI: 10.1177/0884533617700353
- 7) Guasch-Ferré, M., Willett, W.C., The Mediterranean diet and health: a comprehensive overview. *J. Intern. Med.* **2021**, *290*, 549-566. DOI: 10.1111/joim.13333
- 8) Tsigalou, C., Konstantinidis, T., Paraschaki, A., Stavropoulou, E., Voidarou, C., Bezirtzoglou, E., Mediterranean Diet as a Tool to Combat Inflammation and Chronic Diseases. An Overview. *Biomedicines*. **2020**, *8*, 201. DOI: 10.3390/biomedicines8070201
- 9) Azab, A., Top Edible Wild Plants of Eastern Mediterranean Region. Part I: Anticancer Activity. *Eur. J. Med. Plants*. **2023**, *34*, 29-51. DOI: 10.9734/EJMP/2023/v34i61143
- 10) Said, O., Khalil, K., Fulder, S., Azaizeh, H., Ethnopharmacological survey of medicinal herbs in Israel, the Golan Heights and the West Bank region. *J. Ethnopharmacol.* **2002**, *83*, 251-265. DOI: 10.1016/s0378-8741(02)00253-2
- 11) Azaizeh, H., Saad, B., Khalil, K., Said, O., The state of the art of traditional Arab herbal medicine in the eastern region of the Mediterranean: a review. *Evid. Based Complement. Alternat. Med.* **2006**, *3*, 229-235. DOI: 10.1093/ecam/nel034
- 12) Lev, E., Ethno-diversity within current ethno-pharmacology as part of Israeli traditional medicine--a review. *J. Ethnobiol. Ethnomed.* **2006**, *2*, 4. DOI: 10.1186/1746-4269-2-4
- 13) Asadi-Samani, M., Rafieian-Kopaei, M., Azimi, N., Gundelia: a systematic review of medicinal and molecular perspective. *Pak. J. Biol. Sci.* **2013**, *16*, 1238-1247. DOI: 10.3923/pjbs.2013.1238.1247
- 14) Oran, S.A., Al-Eisawi, D., Ethnobotanical survey of the medicinal plants in the central mountains (North-South) in Jordan. *J. Biodiver. Environ. Sci.* **2015**, *6*, 381-400. [[ResearchGate](#)]
- 15) Khalaf, N.A., Naik, R.R., Shakya, A.K., Shalan, N., Al-Othman, A., Antioxidant, anti-inflammatory and anti-diarrheal activity of ethanolic extract of medicinal plants grown in Jordan and Palestine. *Orient. J. Chem.* **2015**, *31*, 1923-1928. DOI:10.13005/ojc/310408
- 16) Abu-Qatouseh, L., Mallah, E., Mansour, K., Evaluation of anti-Propionibacterium acnes and anti-inflammatory effects of polyphenolic extracts of medicinal herbs in Jordan. *Biomed. Pharmacol. J.* **2019**, *12*, 211-217. DOI: 10.13005/bpj/1629

- 17) Kmail, A., Jaradat, N., Mansour, B., Abu-Labdeh, R., Zakarneh, S., Abu-Farha, S., Hussein, F., Issa, L., Saad, B.. Phytochemical analysis, cytostatic, cytotoxic, and anti-inflammatory effects of *Arum palaestinum*, *Ocimum basilicum*, and *Trigonella foenum-graecum* in human monocytic cell line (THP-1)-derived macrophages. *Eur. J. Integr. Med.* **2022**, *54*, 102159. DOI: 10.1016/j.eujim.2022.102159
- 18) Han, C., Wu, X., Zou, N., Zhang, Y., Yuan, J., Gao, Y., Chen, W., Yao, J., Li, C., Hou, J., Qin, D., (2021) *Cichorium pumilum* Jacq Extract Inhibits LPS-Induced Inflammation via MAPK Signaling Pathway and Protects Rats From Hepatic Fibrosis Caused by Abnormalities in the Gut-Liver Axis. *Front. Pharmacol.* **2021**, *12*, 683613. DOI: 10.3389/fphar.2021.683613
- 19) Eltamany, E.E., Mosalam, E.M., Mehanna, E.T., Awad, B.M., Mosaad, S.M., Abdel-Kader, M.S., Ibrahim, A.K., Badr, J.M., Goda, M.S., Potential Gonado-Protective Effect of *Cichorium endivia* and Its Major Phenolic Acids against Methotrexate-Induced Testicular Injury in Mice. *Biomedicines*. **2022**, *10*, 1986. DOI: 10.3390/biomedicines10081986
- 20) Donadio, G., Bellone, M.L., Mensitieri, F.; Parisi, V.; Santoro, V., Vitiello, M.; Dal Piaz, F.; De Tommasi, N., Characterization of Health Beneficial Components in Discarded Leaves of Three Escarole (*Cichorium endivia* L.) Cultivar and Study of Their Antioxidant and Anti-Inflammatory Activities. *Antioxidants*. **2023**, *12*, 1402. DOI: 10.3390/antiox12071402
- 21) Choi, E.M., Hwang, J.K., Anti-inflammatory, analgesic and antioxidant activities of the fruit of *Foeniculum vulgare*. *Fitoterapia*. **2004**, *75*, 557-565. DOI: 10.1016/j.fitote.2004.05.005
- 22) Özbek, H., The Anti-inflammatory Activity of the *Foeniculum vulgare* L. Essential Oil and Investigation. *Int. J. Pharmacol.* **2005**, *50*, 329-331. DOI: 10.3923/ijp.2005.329.331
- 23) Lee, J.H., Lee, D.U., Kim, Y.S., Kim, H.P., 5-Lipoxygenase Inhibition of the Fructus of *Foeniculum vulgare* and Its Constituents. *Biomol. Ther. (Seoul)*. **2012**, *20*, 113-117. DOI: 10.4062/biomolther.2012.20.1.113
- 24) Shalby, A.B., Hamza, A.H., Ahmed, H.H., New insights on the anti-inflammatory effect of some Egyptian plants against renal dysfunction induced by cyclosporine. *Eur. Rev. Med. Pharmacol. Sci.* **2012**, *16*, 455-461. [[ResearchGate](#)]
- 25) Elizabeth, A.A., Josephine, G., Muthiah, N.S., Muniappan, M., Evaluation of analgesic and anti-inflammatory effect of *Foeniculum vulgare*. *Res. J. Pharm. Biol. Chem. Sci.* **2014**, *5*, 658-668. [[ResearchGate](#)]
- 26) Yang, I.J., Yu, H.Y., Lee, D.U., Shin, H.M., Anti-inflammatory Effects of the Fruits of *Foeniculum vulgare* in Lipopolysaccharide Stimulated Macrophages. *Kor. J. Life Sci.*, **2014**, *24*, 981-987. DOI: 10.5352/JLS.2014.24.9.981
- 27) Aazza, S., Lyoussi, B., Megías, C., Cortés-Giraldo, I., Vioque, J., Figueiredo, A.C., Miguel, M.G., Anti-oxidant, anti-inflammatory and anti-proliferative activities of Moroccan commercial essential oils. *Nat. Prod. Commun.* **2014**, *9*, 587-594. [[ResearchGate](#)]
- 28) Tettey, C.O., Yang, I., Ocloo, A., Shin, H.M., Vasorelaxant and Anti-Inflammatory Activities of the Methylene Chloride Fraction of *Foeniculum vulgare* Fruit Extract. *J. Food Biochem.* **2015**, *39*, 55-63. [[ResearchGate](#)]

- 29) Lee, H.S., Kang, P., Kim, K.Y., Seol, G.H., *Foeniculum vulgare* Mill. Protects against Lipopolysaccharide-induced Acute Lung Injury in Mice through ERK-dependent NF- κ B Activation. *Kor. J. Physiol. Pharmacol.* **2015**, *19*, 183-189. DOI:10.4196/kjpp.2015.19.2.183
- 30) Sahane, R., Gokhale, M., Nalawade, P., Wound healing activity of *Foeniculum vulgare* in Sprague Dawley rats. *World J. Pharm. Res.* **2015**, *4*, 1802-1807. [[Google Scholar](#)]
- 31) Pacifico, S., Galasso, S., Piccolella, S., Kretschmer, N., Pan, S-P., Nocera, P., Lettieri, A., Bauer, R., Monaco, P., Winter wild fennel leaves as a source of anti-inflammatory and antioxidant polyphenols. *Arab. J. Chem.* **2018**, *11*, 513-524. DOI:10.1016/j.arabjc.2015.06.026
- 32) Yakut, H.I., Koyuncu, E., Cakir, U., Tayman, C., Koyuncu, İ., Taskin Turkmenoglu, T., Cakir, E., Ozyazici, A., Aydogan, S., Zenciroglu, A., Preventative and therapeutic effects of fennel (*Foeniculum vulgare*) seed extracts against necrotizing enterocolitis. *J. Food Biochem.* **2020**, *44*, e13284. DOI: 10.1111/jfbc.13284
- 33) Setyawati, P., Abdul, A., Reni Ariastuti, D., Anti Inflammatory Activity Test of Ethanol Extract of Fennel Leaves and Fruits (*Foeniculum Vulgare* Mill.) In Wistar Rats. *J. Nutraceut. Herb. Med.* **2021**, *3*, 1-9. DOI: 10.23917/jnhm.v3i2.12847
- 34) Panthong, S., Itharat, A., Aukkanibut, S., Intarawan, T., Anti-inflammatory Effect and Total Flavonoid Content of The Ethanolic Seed Extracts of Three Umbelliferae Species. *Sci. Technol. Asia.* **2020**, 174-181. DOI:10.14456/scitechasia.2020.58
- 35) Korinek, M., Handoussa, H., Tsai, Y-H., Chen, Y-Y., Chen, M-H., Chiou, Z-W., Fang, Y., Chang, F-R., Yen, C-H., Hsieh, C-F., Chen, B-H., El-Shazly, M., Hwang, T-L., Anti-Inflammatory and Antimicrobial Volatile Oils: Fennel and Cumin Inhibit Neutrophilic Inflammation via Regulating Calcium and MAPKs. *Front. Pharmacol.* **2021**, *12*, 674095. DOI: 10.3389/fphar.2021.674095
- 36) Tanveer, M., Shehzad, A., Butt, M.S., Shahid, M., Pharmacodynamics of *foeniculum vulgare* against ulcer, inflammation, hepatotoxicity and nephrotoxicity. *J. Anim. Plant Sci.* **2021**, *31*, 841-853. DOI: 10.36899/JAPS.2021.3.0274
- 37) Servi, H., Şen, A., Yildirim Servi, E., Doğan, A., Chemical composition and biological activities of essential oils of *Foeniculum vulgare* Mill. and *Daucus carota* L. growing wild in Turkey. *J. Res. Pharm.* **2021**, *25*, 142-152. [[Google Scholar](#)]
- 38) Crescenzi, M.A., D'Urso, G., Piacente, S., Montoro, P., UPLC-ESI-QTRAP-MS/MS Analysis to Quantify Bioactive Compounds in Fennel (*Foeniculum vulgare* Mill.) Waste with Potential Anti-Inflammatory Activity. *Metabolites.* **2022**, *12*, 701. DOI:10.3390/metabo12080701
- 39) Nassar, M.I., Aboutabl, E. S., Makled, Y.A., El-Khrisy, E.D., Osman, A.F., Secondary metabolites and pharmacology of *Foeniculum vulgare* Mill. Subsp. Piperitum. *Rev. Latinoam. de Química.* **2010**, *38*, 103-112. [[ResearchGate](#)]
- 40) Keskin, I., Gunal, Y., Ayla, S., Kolbasi, B., Sakul, A., Kilic, U., Gok, O., Koroglu, K., Ozbek, H., Effects of *Foeniculum vulgare* essential oil compounds, fenchone and limonene, on experimental wound healing. *Biotech. Histochem.* **2017**, *92*, 274-282. DOI:10.1080/10520295.2017.1306882

- 41) Zhang, S., Chen, X., Devshilt, I., Yun, Q., Huang, C., An, L., Dorjbat, S., He, X., Fennel main constituent, trans-anethole treatment against LPS-induced acute lung injury by regulation of Th17/Treg function. *Mol. Med. Rep.* **2018**, *18*,1369-1376. DOI:10.3892/mmr.2018.9149
- 42) Yang, I.J., Lee, D.U., Shin, H.M., Anti-inflammatory and antioxidant effects of coumarins isolated from *Foeniculum vulgare* in lipopolysaccharide-stimulated macrophages and 12-O-tetradecanoylphorbol-13-acetate-stimulated mice. *Immunopharmacol.Immunotoxicol.* **2015**,*37*, 308-317. DOI: 10.3109/08923973.2015.1038751
- 43) Oryan, S., Nasri, S., Amin, G R., Kazemi, M.S., Anti nociceptive and anti-inflammatory effects of aerial parts of *Gundeliatournefortii* L. on NMRI male mice.*J.Shahrekord Univ. Med. Sci.***2011**, *12*, 8-15. [[Google Scholar](#)]
- 44) Qnais, E., Bseiso, Y., Wedyan, M., Al-Omari, M., Alkhateeb, H., Chemical composition and antinociceptive effects of essential oil from aerial parts of *Gundeliatournefortii* L Asteraceae (Compositae) in rats. *Trop. J. Pharm. Res.* **2016**, *15*, 2183-2190.DOI:10.4314/tjpr.v15i10.17
- 45) Javanmardi, S., Safari, I., Aghaz, F., Khazaei, M., Wound Healing Activities of *Gundeliatournefortii* L Extract and Milk-Cream Ointment on Second-Degree Burns of Rat Skin. *Int. J. Low Extrem. Wounds.* **2021**,*20*, 272-281. DOI: 10.1177/1534734620921589
- 46) Amer, J., Salhab, A., Jaradat, N., Abdallah, S., Aburas, H., Hattab, S., Ghanim, M., Alqub, M.,*Gundeliatournefortii* inhibits hepatocellular carcinoma progression by lowering gene expression of the cell cycle and hepatocyte proliferation in immunodeficient mice.*Biomed.Pharmacother.* **2022**,*156*, 113885. DOI: 10.1016/j.biopha.2022.113885
- 47) Loizzo, M.R., Menichini, F., Conforti, F., Tundis, R., Bonesi, M., Saab, A.M., Statti, G.A., de Cindio, B., Houghton, P.J., Menichini, F., Frega, N.G., Chemical analysis, antioxidant, antiinflammatory and anticholinesterase activities of *Origanum ehrenbergii*Boiss and *Origanum syriacum* L. essential oils. *Food Chemi.* **2009**, *117*,174-180. DOI: 10.1016/j.foodchem.2009.03.095
- 48) Shen, D., Pan, M.H., Wu, Q.L., Park, C.H., Juliani, H.R., Ho, C.T., Simon. J.E., LC-MS method for the simultaneous quantitation of the anti-inflammatory constituents in oregano (*Origanum* species). *J. Agric. Food Chem.* **2010**,*58*, 7119-7125. DOI: 10.1021/jf100636h
- 49) Awaad, A.S., El-Meligy, R.M., Qenawy, S.A., Atta, A.H., Soliman, G.A., Anti-inflammatory, antinociceptive and antipyretic effects of some desert plants. *J. Saudi Chem. Soc.* **2011**, *15*, 367-373. DOI: 10.1016/j.jscs.2011.02.004
- 50) Assaf, A.M., Amro, B.I., Mashallah, S., Haddadin, R.N., Antimicrobial and anti-inflammatory potential therapy for opportunistic microorganisms. *J. Infect. Dev. Ctries.***2016**,*10*, 494-505. DOI:10.3855/jidc.7610
- 51)Tag, H.M., Henedak, M.M., Abouelhasan, E.M., Hassan, E.A., Zayed, R.A., Mohammed, S.T., Antioxidative and hepatoprotective effects of *Artemisia santonica* and *Origanum syriacum* against *Schistosoma mansoni* infection. *J. Appl. Biol. Sci.* **2020**, *14*, 86-97. [[ResearchGate](#)]

- 52) Tag, H.M., Abouelhassan, E.M., Henedak, M.M., Hassan, E.A., (2021). Effectiveness of *origanum syriacum* on *schistosomamansonii* infected mice: parasitological, histopathological and scanning electron microscopical studies. *Adv. Anim. Vet. Sci.* **2021**, *9*, 221-229. DOI: 10.17582/journal.aavs/2021/9.2.221.229
- 53) Alonazi, M.A., Jemel, I., Moubayed, N., Alwhibi, M., El-Sayed, N.N., Ben Bacha, A., Evaluation of the in vitro anti-inflammatory and cytotoxic potential of ethanolic and aqueous extracts of *Origanum syriacum* and *Salvia lanigera* leaves. *Environ. Sci. Pollut. Res.* **2021**, *28*, 19890-19900. DOI: 10.1007/s11356-020-11961-z
- 54) Pirbalouti, A.G., Azizi, S., Koohpayeh, A., Hamed, B., Wound healing activity of *Malva sylvestris* and *Punica granatum* in alloxan-induced diabetic rats. *Acta Pol. Pharm.* **2010**, *67*, 511-516. [[ResearchGate](#)]
- 55) Pirbalouti, A.G., Koohpayeh, A., Wound Healing Activity of Extracts of *Malva sylvestris* and *Stachys Lavandulifolia*. *Int. J. Biol.* **2011**, *3*, 174-179. DOI: 10.5539/ijb.v3n1p174
- 56) Prudente, A.S., Loddi, A.M., Duarte, M.R., Santos, A.R., Pochapski, M.T., Pizzolatti, M.G., Hayashi, S.S., Campos, F.R., Pontarolo, R., Santos, F.A., Cabrini, D.A., Otuki, M.F., Pre-clinical anti-inflammatory aspects of a cuisine and medicinal millennial herb: *Malva sylvestris* L. *Food Chem. Toxicol.* **2013**, *58*, 324-331. DOI: 10.1016/j.fct.2013.04.042
- 57) Kovalik, A.C., Bisetto, P., Pochapski, M.T., Campagnoli, E.B., Pilatti, G.L., Santos, F.A., Effects of an orabase formulation with ethanolic extract of *Malva sylvestris* L. in oral wound healing in rats. *J. Med. Food.* **2014**, *17*, 618-624. DOI: 10.1089/jmf.2013.0001
- 58) Benso, B., Rosalen, P.L., Alencar, S.M., Murata, R.M., *Malva sylvestris* Inhibits Inflammatory Response in Oral Human Cells. An In Vitro Infection Model. *PLoS ONE*, **2015**, *10*, e0140331. DOI: 10.1371/journal.pone.0140331
- 59) Afshar, M., Ravarian, B., Zardast, M., Moallem, S.A., Fard, M.H., Valavi, M., Evaluation of cutaneous wound healing activity of *Malva sylvestris* aqueous extract in BALB/c mice. *Iran. J. Basic Med. Sci.* **2015**, *18*, 616-622. [[ResearchGate](#)]
- 60) Hamed, A., Rezaei, H., Azarpira, N., Jafarpour, M., Ahmadi, F., Effects of *Malva sylvestris* and its isolated polysaccharide on experimental ulcerative colitis in rats. *J. Evid. Based Complement. Alternat. Med.* **2016**, *21*, 14-22. DOI: 10.1177/2156587215589184
- 61) Benso, B., Franchin, M., Massarioli, A.P., Paschoal, J.A., Alencar, S.M., Franco, G.C., Rosalen, P.L., Anti-Inflammatory, Anti-Osteoclastogenic and Antioxidant Effects of *Malva sylvestris* Extract and Fractions: *In Vitro* and *In Vivo* Studies. *PLoS ONE*, **2016**, *11*, e0162728. DOI: 10.1371/journal.pone.0162728
- 62) Martins, C.A., Campos, M.L., Irioda, A.C., Stremel, D.P., Trindade, A.C., Pontarolo, R., Anti-Inflammatory Effect of *Malva sylvestris*, *Sida cordifolia*, and *Pelargonium graveolens* Is Related to Inhibition of Prostanoid Production. *Molecules*. **2017**, *22*, 1883. DOI: 10.3390/molecules22111883
- 63) Jomezadeh, V., Sheibani, S., Tavassoli, A., Karimani, A., Zirak, M.R., Mohammadpour, A. H., Khakzad, M.R., Afshar, M., Tavassoli, F., Effect of *Malva sylvestris* extract on postoperative peritoneal adhesion in rats. *Jundishapur J. Nat. Pharm. Prod.* **2019**, *14*, e12069. DOI: doi: 10.5812/jjnpp.12069

- 64) Seddighfar, M., Mirghazanfari, S.M., Dadpay, M., Analgesic and anti-inflammatory properties of hydroalcoholic extracts of *Malva sylvestris*, *Carum carvi* or *Medicago sativa*, and their combination in a rat model. *J.Integr. Med.* **2020**,18, 181-188. DOI:10.1016/j.joim.2020.02.003
- 65) Xiao, Y., Oumarou, D.B., Wang, S., Liu, Y., Circular RNA Involved in the Protective Effect of *Malva sylvestris* L. on Myocardial Ischemic/Re-Perfused Injury. *Front. Pharmacol.* **2020**,11, 520486. DOI: 10.3389/fphar.2020.520486
- 66) Benso, B., Rosalen, P.L., Pasetto, S., Marquezin, M.C., Freitas-Blanco, V., Murata, R.M., *Malva sylvestris* derivatives as inhibitors of HIV-1 BaL infection. *Nat. Prod. Res.* **2021**,35, 1064-1069. DOI: 10.1080/14786419.2019.1619720
- 67) Meysami, M., Hashempur, M.H., Kamalinejad, M., Emtiazy, M., Efficacy of Short Term Topical *Malva Sylvestris* L. Cream in Pediatric Patients with Atopic Dermatitis: A Randomized Double-Blind Placebo-Controlled Clinical Trial. *Endocr. Metab. Immun. Disord. Drug Targets.* **2021**, 21(, 1673-1678. DOI: 10.2174/1871530320666201023125411
- 68) Mustafa, A., Ali, M., Comparative Phyto and Physico-chemical Standardization of Fresh and Different market samples with the Anti-inflammatory Studies of Fruit Parts of *Malva sylvestris* L. *Res. J.Pharmacol.Pharmacodynam.* **2022**, 14, 219-224. DOI: 10.52711/2321-5836.2022.00038
- 69) Contardi, M., Ayyoub, A.M., Summa, M., Kossyvaki, D., Fadda, M., Liessi, N., Armirotti, A., Fragouli, D., Bertorelli, R., Athanassiou, A., Self-Adhesive and Antioxidant Poly(vinylpyrrolidone)/Alginate-Based Bilayer Films Loaded with *Malva sylvestris* Extracts as Potential Skin Dressings. *ACS Appl. Bio. Mater.* **2022**,5, 2880-2893. DOI:10.1021/acsabm.2c00254
- 70) Mohaghegh, Z., Golfakhrabadi, F., Faal Siyahkal, S., Dastoorpoor, M., Zaheri Abdevand, Z., Montazeri, S., Sharifipour, F., Bahmaei, H., Javadnoori, M., The Effect of *Malva Sylvestris* Cream on Episiotomy Pain and Healing: A Randomized Controlled Clinical Trial. *Int. J. Community Based Nurs. Midwifery.* **2022**, 10, 248-258. DOI:10.30476/IJCBNM.2022.95772.2082
- 71) Belkhodja, H., Bouhadi, D., Sedjrari, K., Sehanine, S. (2023). Evaluation of the Anti-inflammatory and Anti-hemolytic Potential of Polyphenolic Components of Common Mallow (*Malva sylvestris*). *Asian J. Dairy and Food Res.* **2023**, DRF-321. DOI:10.18805/ajdfr.DRF-321
- 72) Abu-Gharbieh, E.F., Bustanji, Y.K., Mohammad, M.K., In vitro effects of *Micromeria fruticosa* on human leukocyte myeloperoxidase activity. *J. Pharm. Res.* **2010**, 3, 2492-2493. [[ResearchGate](#)]
- 73) Abu-Gharbieh, E., Shehab, N.G., Khan, S. A. Anti-inflammatory and gastroprotective activities of the aqueous extract of *Micromeria fruticosa* (L.) Druce ssp *Serpyllifolia* in mice. *Pak. J. Pharm. Sci.* **2013**, 26, 799-803. [[ResearchGate](#)]
- 74) El-Sayed, N.H., El-Eraky, W., Ibrahim, M.T., Mabry, T.J., Anti-inflammatory and ulcerogenic activities of *Salvia triloba* extracts. *Fitoterapia.* **2006**,77, 333-335. DOI: 10.1016/j.fitote.2006.04.002

- 75) Bzour, J.F., Oran, S.A., Khaleel, M., Mashallah, S., Bustanji, Y., Inhibitory effects of methanolic extract of *Salvia fruticosa* Mill. on pro-inflammatory cytokines production in RAW 264.7 in vitro cellular model and in Balb/c mice in vivo animal model. *J. Med. Plants Res.* **2011**, 5, 6662-6671. DOI: 10.5897/JMPR11.586
- 76) Çadirci, E., Süleyman, H., Gürbüz, P., UzKuruüzüm, A., Güvenalp, Z., DemirezerÖmür, L., Anti-inflammatory effects of different extracts from three *Salvia* species. *Turk. J. Biol.* **2012**, 36, 59-64. DOI: 10.3906/biy-1001-17
- 77) Elmezogi, J., Zetrini, A., Ben-Hussein, G., Anwair, M., Gbaj, A., El-Ashheb, M., Nahar, L. and Sarker, S.D., Evaluation of anti-inflammatory activity of some Libyan medicinal plants in experimental animals. *Arch. Biol. Sci.* **2012**, 64,1059-1063. DOI:10.2298/ABS1203059E
- 78) Ahmed, H.H., Salem, A.M., Sabry, G.M., Husein, A.A., Kotob, S.E., Possible therapeutic uses of *Salvia triloba* and *Piper nigrum* in Alzheimer's disease-induced rats. *J. Med. Food.* **2013**, 16, 437-446. DOI: 10.1089/jmf.2012.0165
- 79) Boukhary, R., Raafat, K., Ghoneim, A.I., Aboul-Ela, M., El-Lakany, A., Anti-Inflammatory and Antioxidant Activities of *Salvia fruticosa*: An HPLC Determination of Phenolic Contents. *Evid. Based Complement. Alternat. Med.* **2016**, 7178105. DOI:10.1155/2016/7178105
- 80) Risaliti, L., Kehagia, A., Daoulzi, E., Lazari, D., Bergonzi, M.C., Vergkizi-Nikolakaki, S.,Hadjipavlou-Litina, D.,Bilia, A.R., Liposomes loaded with *Salvia triloba* and *Rosmarinus officinalis* essential oils: In vitro assessment of antioxidant, antiinflammatory and antibacterial activities. *J. Drug Deliv. Sci. Technol.* **2019**, 51, 493-498. DOI:10.1016/j.jddst.2019.03.034
- 81) Brindisi, M., Bouzidi, C., Frattaruolo, L., Loizzo, M.R., Cappello, M.S., Dugay, A.,Deguin, B., Lauria, G., Cappello, A.R.,Tundis, R., New Insights into the Antioxidant andAnti-Inflammatory Effects of Italian*Salvia officinalis* Leaf and FlowerExtracts in Lipopolysaccharide andTumor-Mediated InflammationModels. *Antioxidants.***2021**, 10, 311. DOI: 10.3390/antiox10020311
- 82) Yang, R., Zhou, Q., Wen, C., Hu, J., Li, H., Zhao, M., Zhao, H., Mustard seed (*Sinapis Alba* Linn) attenuates imiquimod-induced psoriasiform inflammation of BALB/c mice. *J. Dermatol.* **2013**,40, 543-552. DOI: 10.1111/1346-8138.12119
- 83) Lee, K-Y.,Hong, C-H., Inhibitory Effect of Semen *Sinapis Alba* onImmediate Hypersensitivity Reaction. *J. Korean Med.Ophthalmol.Otolaryngol. Dermatol.***2014**, 27, 177-188. DOI: 10.6114/jkood.2014.27.4.177
- 84) Xian, Y.F., Hu, Z., Ip, S.P., Chen, J.N., Su, Z.R., Lai, X.P., Lin, Z.X., Comparison of the anti-inflammatory effects of *Sinapis alba* and *Brassica juncea* in mouse models of inflammation. *Phytomedicine.* **2018**,50, 196-204. DOI: 10.1016/j.phymed.2018.05.010
- 85) Safdar, M.N., Baig, U.Y., Riaz, M.M., Mumtaz, A., Jabbar, S., Ur-Rehman, N., Ahmad, Z., Malik, H., Yousaf, S., Extraction of polyphenols from different herbs for the development of functional date bars. *Food Sci. Technol.* **2022**, 42,e43521. DOI: 10.1590/fst.43521

- 86) Mikucka, W., Zielinska, M., Bulkowska, K., Witonska, I., Recovery of polyphenols from distillery stillage by microwave-assisted, ultrasound-assisted and conventional solid-liquid extraction. *Sci. Rep.* **2022**,*12*, 3232. DOI: 10.1038/s41598-022-07322-0
- 87) Boeing, J.S., Barizão, É.O., E Silva, B.C., Montanher, P.F., de Cinque Almeida, V., Visentainer, J.V., Evaluation of solvent effect on the extraction of phenolic compounds and antioxidant capacities from the berries: application of principal component analysis. *Chem. Cent. J.* **2014**,*8*, 48. DOI: 10.1186/s13065-014-0048-1
- 88) Nguyen, N.Q., Nguyen, M.T., Nguyen, V.T., Le, V.M., Trieu, L.H., Le, X.T., Khang, T.V., Giang, N.T., Thach, N.Q., Hung, T.T., The effects of different extraction conditions on the polyphenol, flavonoids components and antioxidant activity of *Polysciasfruticosa* roots. *IOP Conf.Ser. Mater.Sci.Eng.* **2020**,*736*, 022067. DOI: 10.1088/1757-899X/736/2/022067
- 89) Onyango, I.G., Jauregui, G.V., Cárna, M., Bennett, J.P., Stokin, G.B., Neuroinflammation in Alzheimer's Disease. *Biomedicines.* **2021**, *9*, 524. DOI:10.3390/biomedicines9050524
- 90) Kurowska, A., Ziemichód, W., Herbet, M., Piatkowska-Chmiel, I., The Role of Diet as a Modulator of the Inflammatory Process in the Neurological Diseases. *Nutrients.* **2023**, *15*, 1436. DOI: 10.3390/nu15061436
- 91) García-Sánchez, A., Gómez-Hermosillo, L., Casillas-Moreno, J., Pacheco-Moisés, F., Campos-Bayardo, T.I., Román-Rojas, D., Miranda-Díaz, A.G., Prevalence of Hypertension and Obesity: Profile of Mitochondrial Function and Markers of Inflammation and Oxidative Stress. *Antioxidants.* **2023**, *12*, 165. DOI: 10.3390/antiox12010165
- 92) Moriki, D., Antonogeorgos, G., Koumpagioti, D., Chaloutsi, D., Ellwood, P., García-Marcos, L., Priftis, K.N., Douros, K., Panagiotakos, D., Adherence to an Anti-Inflammatory Diet and Atopic Diseases' Prevalence in Adolescence: The Greek Global Asthma Network Study. *Nutrients.* **2023**, *15*, 3191. DOI: 10.3390/nu15143191
- 93) Stumpf, F., Keller, B., Gressies, C., Schuetz, P., Inflammation and Nutrition: Friend or Foe? *Nutrients.* **2023**, *15*, 1159. DOI: 10.3390/nu15051159
- 94) Abu-Reidah, I.M., Ali-Shtayeh, M.S., Jamous, R.M., Arráez-Román, D., Segura-Carretero, A., Comprehensive metabolite profiling of *Arum palaestinum* (Araceae) leaves by using liquid chromatography–tandem mass spectrometry. *Food Res. Int.* **2015**, *70*, 74–86. DOI: 10.1016/j.foodres.2015.01.023
- 95) Degim, I.T., Uslu, A., Hadgraft, J., Atay, T., Akay, C., Cevheroglu, S. The effects of Azone and capsaicin on the permeation of naproxen through human skin. *Int. J. Pharm.* **1999**,*179*, 21–25. DOI: 10.1016/s0378-5173(98)00353-6
- 96) Cheriyan, B.V., Kadhirvelu, P., Nadipelly, J., Shanmugasundaram, J., Sayeli, V., Subramanian, V., Anti-nociceptive Effect of 7-methoxy Coumarin from *Eupatorium Triplinerveahl* (Asteraceae). *Pharmacogn. Mag.* **2017**, *13*, 81–84. DOI: 10.4103/0973-1296.197650
- 97) Kisiel, W., Michalska, K., & Szneler, E., Norisoprenoids from aerial parts of *Cichorium pumilum*. *Biochem.Syst.Ecol.* **2004**, *3*, 343–346. DOI: 10.1016/j.bse.2003.08.005

- 98) Yang, E.J., Ham, Y.M., Lee, W.J., Lee, N.H., Hyun, C.G., Anti-inflammatory effects of apo-9'-fucoxanthinone from the brown alga, *Sargassum muticum*. *Daru*. **2013**,*21*,62. DOI: 10.1186/2008-2231-21-62
- 99) Kim, H-S.,Shanura Fernando, I.P., Lee, S-H., Ko,S-C., Kang, M.C., Ahn, G., Je, J-G., Sanjeeva, K.K., Rho, J-R., Shin, H-J., Lee, W-W., Lee, D-S., Jeon, Y-J., Isolation and characterization of anti-inflammatory compounds from *Sargassum hornerivia* high-performance centrifugal partition chromatography and high-performance liquid chromatography. *Algal Res*. **2021**,*54*, 102209. DOI: 10.1016/j.algal.2021.102209
- 100) Cornea-Cipcigan, M., Bunea, A.,Bouari, C.M.,Pamfil, D., Páll, E.,Urcan, A.C.,MARGAOAN, R., Anthocyanins and Carotenoids Characterization in Flowers and Leaves of Cyclamen Genotypes Linked with Bioactivities Using Multivariate Analysis Techniques. *Antioxidants*.**2022**, *11*, 1126. DOI: 10.3390/antiox11061126
- 101) Rasmus, P., Kozłowska, E., Antioxidant and Anti-Inflammatory Effects of Carotenoids in Mood Disorders: An Overview. *Antioxidants*. **2023**, *12*, 676. DOI:10.3390/antiox12030676
- 102) Ishizaka, H., Yamada, H., Sasaki, K., Volatile compounds in the flowers of *Cyclamen persicum*, *C. purpurascens* and their hybrids. *Sci.Hortic*. **2022**, *94*, 125-135. DOI:10.1016/S0304-4238(01)00362-4
- 103) Jung, Y.Y., Hwang, S.T., Sethi, G., Fan, L., Arfuso, F., Ahn, K.S., Potential Anti-Inflammatory and Anti-Cancer Properties of Farnesol. *Molecules*. **2018**,*23*, 2827. DOI: 10.3390/molecules23112827
- 104) Alkhanjaf, A.A., Athar, M.T., Ullah, Z., Alsayhab, A.M., Umar, A., Shaikh, I.A., Farnesol Protects against Cardiotoxicity Caused by Doxorubicin-Induced Stress, Inflammation, and Cell Death: An In Vivo Study in Wistar Rats. *Molecules*. **2022**,*27*, 8589. DOI: 10.3390/molecules27238589
- 105) Lin, C-Y., Kao, S-H., Hung, L-C., Chien, H-J., Wang, W-H., Chang, Y-W., Chen, Y-H., Lipopolysaccharide-Induced Nitric Oxide and Prostaglandin E2 Production Is Inhibited by Tellimagrandin II in Mouse and Human Macrophages. *Life*,**2021**, *11*,411. DOI: 10.3390/life11050411
- 106) Özyay, C., Karagür, E. R., Akça, H., Mammadov, R., *Cyclamen* L. Inhibits nitric oxide production in LPS-stimulated NSCLC cells. *Jordan J. Pharm. Sci.* **2020**, *13*, 257-264. [[ResearchGate](#)]
- 107) Arif, A., Bhatti, A., John, P., Therapeutic Potential Of *Foeniculum vulgare* Mill. Derived Selenium Nanoparticles In Arthritic Balb/c Mice. *Int. J.Nanomed*. **2019**,*14*, 8561-8572. DOI: 10.2147/IJN.S226674
- 108) Gias, Z.T., Afsana, F., Debnath, P., Alam, M.S., Ena, T.N., Hossain, M.H., Jain, P., Reza, H.M., A mechanistic approach to HPLC analysis, antinociceptive, anti-inflammatory and postoperative analgesic activities of panchphoron in mice. *BMC Complement. Med. Ther.* **2020**,*20*, 102. DOI: 10.1186/s12906-020-02891-x
- 109) Abu-Lafi, S., Odeh, I., Dewik, H., Qabajah, M., Hanus, L.O., Dembitsky, V.M., Thymol and carvacrol production from leaves of wild Palestinian *Majorana syriaca*. *Bioresour. Technol.* **2008**,*99*, 3914-3918. DOI: 10.1016/j.biortech.2007.07.042

- 110) Lima, M., Quintans-Júnior, L.J., de Santana, W.A., Martins Kaneto, C., Pereira Soares, M.B., Villarreal, C.F., Anti-inflammatory effects of carvacrol: evidence for a key role of interleukin-10. *Eur. J.Pharmacol.* **2013,699**, 112-117. DOI: 10.1016/j.ejphar.2012.11.040
- 111) Riella, K.R., Marinho, R.R., Santos, J.S., Pereira-Filho, R.N., Cardoso, J.C., Albuquerque-Junior, R.L., Thomazzi, S.M., Anti-inflammatory and cicatrizing activities of thymol, a monoterpene of the essential oil from *Lippiagracilis*, in rodents. *J.Ethnopharmacol.* **2012,143**, 656-663. DOI: 10.1016/j.jep.2012.07.028
- 112) Gasparetto, J.C., Martins, C.A., Hayashi, S.S., Otuky, M.F., Pontarolo, R., Ethnobotanical and scientific aspects of *Malva sylvestris* L.: a millennial herbal medicine. *J. Pharm.Pharmacol.* **2012,64**, 172-189. DOI: 10.1111/j.2042-7158.2011.01383.x
- 113) Paul, D., A review on biological activities of common Mallow (*Malva sylvestris* L.). *J. Life Sci.* **2016**, 4, 1-5.[[Google Scholar](#)]
- 114) Mousavi, S.M., Hashemi, S.A., Behbudi, G., Mazraedoost, S., Omidifar, N., Gholami, A., Chiang, W.H., Babapoor, A., PynadathuRumjit, N., A Review on Health Benefits of *Malva sylvestris* L. Nutritional Compounds for Metabolites, Antioxidants, and Anti-Inflammatory, Anticancer, and Antimicrobial Applications. *Evid. Based Complement. Alternat. Med.* **2021,2021**, 5548404. DOI: 10.1155/2021/5548404
- 115) Merez-Sadowska, A., Sitarek, P., Kowalczyk, T., Zajdel, K., Jecek, M., Nowak, P., Zajdel, R., FoodAnthocyanins: Malvidin and ItsGlycosides as Promising Antioxidantand Anti-Inflammatory Agents withPotential Health Benefits. *Nutrients.* **2023**, 15, 3016. DOI: 10.3390/nu15133016
- 116) Batiha, G.E., Tene, S.T., Teibo, J.O., Shaheen, H.M., Oluwatoba, O.S., Teibo, T.K., Al-Kuraishy, H.M., Al-Garbee, A.I., Alexiou, A., Papadakis, M., The phytochemical profiling, pharmacological activities, and safety of *Malva sylvestris*: a review. *NaunynSchmiedebergs Arch.Pharmacol.* **2023,396**, 421-440. DOI: 10.1007/s00210-022-02329-w
- 117) Wang, C.H., Zhu, L.L., Ju, K.F., Liu, J.L., Li, K.P., Anti-inflammatory effect of delphinidin on intramedullary spinal pressure in a spinal cord injury rat model. *Exp. Ther. Med.* **2017,14**, 5583-5588. DOI: 10.3892/etm.2017.5206
- 118) Gürbüz, P., Dokumaci, A.H., Gündüz, M.G., Perez, C., Göger, F., Paksoy, M.Y., Yerer, M.B., ÖmürDemirezer, L., *In vitro* biological activity of *Salvia fruticosa* Mill. infusion against amyloid β -peptide-induced toxicity and inhibition of GSK-3 β , CK-1 δ , and BACE-1 enzymes relevant to Alzheimer's disease. *Saudi Pharm. J.* **2021,29**, 236-243. DOI: 10.1016/j.jsps.2021.01.007
- 119) Rocha, J., Eduardo-Figueira, M., Barateiro, A., Fernandes, A., Brites, D., Bronze, R., Duarte, C.M., Serra, A.T., Pinto, R., Freitas, M., Fernandes, E., Silva-Lima, B., Mota-Filipe, H., Sepodes, B., Anti-inflammatory effect of rosmarinic acid and an extract of *Rosmarinus officinalis* in rat models of local and systemic inflammation. *Basic Clin.Pharmacol. Toxicol.* **2015,116**, 398-413. DOI: 10.1111/bcpt.12335
- 120) Bao, L., Ji, R., Jin, Y., Sun, S., AReview of the Anti-InflammatoryEffects ofRosmarinic Acid onInflammatory Diseases.*Front. Pharmacol.***2020,11**, 153. DOI: 10.3389/fphar.2020.00153

- 121) Wang, S.X., Hu, L.M., Gao, X.M., Guo, H., Fan, G.W., Anti-inflammatory activity of salvianolic acid B in microglia contributes to its neuroprotective effect. *Neurochem. Res.* **2010**,35, 1029-1037. DOI: 10.1007/s11064-010-0151-1
- 122) Al-Qudah, M.A., Al-Jaber, H.I., Muhaidat, R., Hussein, E.I., Abdel, A.A., Hamid, A.S., Chemical composition and antimicrobial activity of the essential oil from *Sinapis alba* L. and *Sinapis arvensis* L.(Brassicaceae) growing wild in Jordan. *Res. J. Pharm. Biol. Chem. Sci.* **2011**, 2, 1136-1144. [[ResearchGate](#)]
- 123) Peng, C., Zhao, S.Q., Zhang, J., Huang, G.Y., Chen, L.Y., Zhao, F.Y., Chemical composition, antimicrobial property and microencapsulation of Mustard (*Sinapis alba*) seed essential oil by complex coacervation. *Food Chem.* **2014**,165, 560-568. DOI:10.1016/j.foodchem.2014.05.126
- 124) Singh, S., Das, S.S., Singh, G., Perroti, M., Schuff, C., Catalán, C.A., Comparison of chemical composition, antioxidant and antimicrobial potentials of essential oils and oleoresins obtained from seeds of *Brassica juncea* and *Sinapis alba*. *MOJ Food Process. Technol.* **2017**, 4, 113-120. [[ResearchGate](#)]
- 125) Sujatha, M., Karthika, K., Sivakamasungari, S., Mariajancyrani, J., Chandramohan, G., GC-MS analysis of phytochemicals and total antioxidant activity of hexane extract of *Sinapis alba*. *Int. J. Pharm. Chem. Biol. Sci.* **2014**, 4, 112-117. [[Google Scholar](#)]
- 126) Torrijos, R., Righetti, L., Cirlini, M., Calani, L., Mañes, J., Meca, G., Dall'Asta, C., Phytochemical profiling of volatile and bioactive compounds in yellow mustard (*Sinapis alba*) and oriental mustard (*Brassica juncea*) seed flour and bran. *LWT Food Sci. Technol.* **2023**, 173, 114221. DOI: 10.1016/j.lwt.2022.114221
- 127) Subedi, L., Venkatesan, R., Kim, S.Y., Neuroprotective and Anti-Inflammatory Activities of Allyl Isothiocyanate through Attenuation of JNK/NF- κ B/TNF- α Signaling. *Int. J. Mol. Sci.* **2017**,18, 1423. DOI: 10.3390/ijms18071423
- 128) Lee, Y.M., Seon, M.R., Cho, H.J., Kim, J.S., Park, J.H., Benzyl isothiocyanate exhibits anti-inflammatory effects in murine macrophages and in mouse skin. *J. Mol. Med. (Berl).* **2009**,87, 1251-1261. DOI: 10.1007/s00109-009-0532-6
- 129) El Badawy, S.A., Ogaly, H.A., Abd-Elsalam, R.M., Azouz, A.A., Benzyl isothiocyanates modulate inflammation, oxidative stress, and apoptosis via Nrf2/HO-1 and NF- κ B signaling pathways on indomethacin-induced gastric injury in rats. *Food Funct.* **2021**, 12, 6001-6013. DOI: 10.1039/d1fo00645b
- 130) Koo, H.J., Song, Y.S., Kim, H.J., Lee, Y.H., Hong, S.M., Kim, S.J., Kim, B.C., Jin, C., Lim, C.J., Park, E.H., Antiinflammatory effects of genipin, an active principle of gardenia. *Eur. J. Pharmacol.* **2004**,495, 201-208. DOI: 10.1016/j.ejphar.2004.05.031
- 131) Chen, X.Y., Jiang, W.W., Liu, Y.L., Ma, Z.X., Dai, J.Q., Anti-inflammatory action of geniposide promotes wound healing in diabetic rats. *Pharm. Biol.* **2022**,60, 294-299. DOI: 10.1080/13880209.2022.2030760
- 132) Vo, Q.V., Nam, P.C., Dinh, T.N., Mechler, A., Van Tran, T.T., Anti-inflammatory activity of synthetic and natural glucoraphanin. *J. Serbian Chem. Soc.* **2019**, 84, 445-453. [[Google Scholar](#)]

- 133) Theodosios-Nobelos, P., Papagiouvannis, G., Rekka, E.A., Ferulic, Sinapic, 3,4-Dimethoxycinnamic Acid and Indomethacin Derivatives with Antioxidant, Anti-Inflammatory and Hypolipidemic Functionality. *Antioxidants*. **2023**, *12*, 1436. DOI: 10.3390/antiox12071436
- 134) Li, Y., Li, J., Su, Q., Yuanfa Liu, Y., Sinapine reduces non-alcoholic fatty liver disease in mice by modulating the composition of the gut microbiota. *Food Funct*. **2019**, *10*, 3637–3649. DOI: 10.1039/c9fo00195f
- 135) Ranaweera, S.S., Dissanayake, C.Y., Natraj, P., Lee, Y.J., Han, C.H., Anti-inflammatory effect of sulforaphane on LPS-stimulated RAW 264.7 cells and ob/ob mice. *J. Vet. Sci.* **2020**, *21*, e91. DOI: 10.4142/jvs.2020.21.e91
- 136) Fernandez-Prades, L., Brasal-Prieto, M., Alba, G., Martin, V., Montserrat-de la Paz, S., Cejudo-Guillen, M., Santa-Maria, C., Dakhaoui, H., Granados, B., Sobrino, F., Palomares, F., Lopez-Enriquez, S., Sulforaphane Reduces the Chronic Inflammatory Immune Response of Human Dendritic Cells. *Nutrients*. **2023**, *15*, 3405. DOI: 10.3390/nu15153405
- 137) El Azhary, K., Tahiri Jouti, N., El Khachibi, M., Moutia, M., Tabyaoui, I., El Hou, A., Achtak, H., Nadifi, S., Habti, N., Badou, A., Anti-inflammatory potential of *Capparis spinosa* L. in vivo in mice through inhibition of cell infiltration and cytokine gene expression. *BMC Complement. Altern. Med.* **2017**, *17*, 81. DOI: 10.1186/s12906-017-1569-7
- 138) Küpeli Akkol, E., Türkcanoğlu, G., Taştan, H., Sobarzo-Sánchez, E., Prevention of Inflammation Initiation on Acetic Acid-Induced Ulcerative Colitis in Rats by *Malva nicaeensis* All. *Front. Biosci. (Landmark Ed)*. **2023**, *28*, 142. doi: 10.31083/j.fb12807142
- 139) Minaiyan, M., Ghassemi, D.N., Mahzouni, P., Ansari, R.M., Effect of *Matricaria aurea* (Loefl.) Shultz-Bip. hydroalcoholic extract on acetic acid-induced acute colitis in rats. *Iran. J. Basic Med. Sci.* **2011**, *14*, 67-74. [[Google Scholar](#)]
- 140) Azab, A., Nassar, A., Kaplanski, J., Mahajneh, R., Agam, G., Azab, A.N., Effects of aqueous extract of *Notobasis sylvatica* on lipopolysaccharide-induced inflammation in rats. *Asian Pac. J. Trop. Med.* **2018**, *11*, 48-52. DOI: 10.4103/1995-7645.223533
- 141) Miao, L., Tao, H., Peng, Y., Wang, S., Zhong, Z., El-Seedi, H., Dragan, S., Zengin, G., Cheang, W.S., Wang, Y., Xiao, J., The anti-inflammatory potential of *Portulaca oleracea* L. (purslane) extract by partial suppression on NF-κB and MAPK activation. *Food Chem.* **2019**, *290*, 239-245. DOI: 10.1016/j.foodchem.2019.04.005
- 142) Bae, J-Y., Lee, Y-S., Han, S.Y., Jeong, E.J., Lee, M.K., Kong, J.Y., Lee, D.H., Cho, K.J., Lee, H-S., Ahn, M-J., A Comparison between Water and Ethanol Extracts of *Rumex acetosa* for Protective Effects on Gastric Ulcers in Mice. *Biomol. Ther. (Seoul)*. **2012**, *20*, 425-430. DOI: 10.4062/biomolther.2012.20.4.425
- 143) De Felice, A., Bader, A., Leone, A., Sosa, S., Loggia, R.D., Tubaro, A., De Tommasi, N., New polyhydroxylated triterpenes and anti-inflammatory activity of *Salvia hierosolymitana*. *Planta Med.* **2006**, *72*, 643-649. DOI: 10.1055/s-2006-931573
- 144) Mzid, M., Ben Khedir, S., Bardaa, S., Sahnoun, Z., Rebai, T., Chemical composition, phytochemical constituents, antioxidant and anti-inflammatory activities of *Urtica urens* L. leaves. *Arch. Physiol. Biochem.* **2017**, *123*, 93-104. DOI: 10.1080/13813455.2016.1255899