

INTERACTION OF RED CELL INDICES AND BLOOD GROUP IN DIABETIC PATIENTS ABOVE 50 YEARS OF AGE

ABSTRACT

A group of metabolic diseases with a hyperglycemic phenotype known as diabetes mellitus impact both industrialized and developing nations. It is brought on by deviations in insulin synthesis, action, or both. This study seeks to associate the interaction of red cell indices and blood group in diabetic patients above 50 years of age. In this cross-sectional study, 100 individuals were recruited from a consenting population at a private hospital in Port Harcourt, Rivers State, 50 of whom had diabetes and the other 50 of whom had not. The subjects were people older than 50. To ascertain the red cell indices, the whole blood count 3-part analyzer was used to analyze the blood samples for FBC and grouping of cells using the tile approach to determine the ABO Blood Groups. Results revealed that blood group O, with a total of 25 (50%), had the highest prevalence of diabetes mellitus. Blood group AB came in last with a total of 5 participants (10%), followed by blood group A with a total of 12 subjects (24%) and blood group B with a total of 8 subjects (16%) respectively. In this study, blood group O predominated over blood groups A and B (47% against 25% versus 17%, respectively). The least common blood type in this study was AB (11%). When the PCV, HB, RBC, MCV, MCH, MCHC, and RDW were examined between diabetic and non-diabetic participants, no statistically significant ($p < 0.05$) difference was found. The relative P-values for these variables were 0.5736, 0.6005, 0.9605, 0.6005, 0.9225, 0.4454, and 0.2682 respectively. According to the results of our study, diabetics are more likely to have the blood types O and AB. We can infer that blood type might be a risk factor and might be helpful in determining the condition.

Key Words. Diabetes, Red cell indices, ABO Blood Groups

Comment [A1]: Its only plain summary, shall be better to write an introduction, research objective, method, result, conclusion and recommendation separately and precisely
Abbreviations and acronyms are not also recommended in the abstract

1. Introduction

Both industrialized and developing nations experience diabetes mellitus (WHO, 2014), a hyperglycemic phenotypic group of metabolic diseases, brought on by abnormalities in insulin synthesis, action, or both. The complex interaction between inherited and environmental factors causes these errors (Sarwar, 2014). In 2013, there were 382 million cases of diabetes globally, up from an estimated 30 million cases in 1985, according to the World Health Organization (WHO).

The World Health Organization (WHO) describes diabetes mellitus as a chronic metabolic disorder characterized by elevated blood glucose levels that, over time, have an impact on the heart, blood vessels, kidneys, eyes, and nerves (WHO, 2014). 1.7% of the people in Nigeria and the rest of the world suffer from diabetes mellitus, which is a serious concern (Wokoma *et al.*, 2002). According to estimates made by Wild *et al.*, (2004), there would be 366,212 million diabetics worldwide by the year 2030. In Rivers State, the prevalence of diabetes mellitus has climbed to roughly 42.7%. However, according to the International Diabetes Federation (IDF), by 2045, adults aged 40 and older are expected to have a diabetes rate of about 10.9%, up roughly 9.3% from 2019. (Saeed *et al.*, 2019).

Families pass down blood type antigens, which are crucial for comprehending genetics, inheritance patterns, disease susceptibility, and the security of blood transfusions. Blood group antigens have been connected to a number of disorders, both in their absence and presence. Due to the fact that diabetes mellitus and blood type share the same genetic and immunological foundations (Desisto *et al.*, 2014). The relationship between the ABO blood group and disease risk has recently become the subject of extensive research.

Thankfully, new research from authors like Anifowoshe *et al.* (2017) and Oladele *et al.* (2020) indicates that ABO blood types affect disease susceptibility, which can be used as a management strategy. Studies by Qureshi and Bhatti (2003) and Bener and Yousafzai (2014), among others, also demonstrate that ABO blood groups affect diabetes susceptibility. ABO blood groups are hereditary and are controlled by a gene on chromosome nine known as the ABO gene or immunoglobulin (Zahid *et al.*, 2016). Thus, some cases of diabetes may be genetic (Abdel-Gaffar *et al.*, 2020; Yahaya *et al.*, 2020).

Additionally, there is evidence supporting the notion that demographic and environmental factors have an impact on illness susceptibility (Yahaya *et al.*, 2017). Shittu *et al.* (2017) advised that high prevalence of diabetes and pre-diabetes among women and those younger than 61 in Oyo State, Nigeria, is noteworthy. In all, these revealed that the association of ABO blood groups and red cell indices can offer basic information for controlling a person's health as well as figure out how diseases are distributed among diabetics.

Consequently, it is necessary to ascertain the distribution of the ABO blood group and among diabetics in a given population so as to effectively employ ABO blood types as well as red cell indices to anticipate and manage a disease. In Port Harcourt, Rivers State, Nigeria, as in other underdeveloped countries, there are few such studies among the diabetic community. Hence, this study aims at ascertaining the relationship between ABO blood groups and red cell indices in diabetics visiting a private hospital in Port Harcourt Rivers State, Nigeria.

RBC parameters are part of the Complete Blood Cell Count (CBC), a panel of analytical assays typically used to distinguish between various kinds of anemia (Doig and Zhang, 2017). Red Blood Cell Count (RBC), Hematocrit (HCT), Hemoglobin (HGB), Mean Corpuscular Volume (MCV), Mean Corpuscular Hemoglobin (MCH), Mean Corpuscular Hemoglobin Concentration (MCHC), and Red Cell Distribution Width (RDW) are RBC parameters (Vinupritha *et al.*, 2017). Continuously exposing RBCs to hyperglycemia causes hemoglobin protein to be permanently glycated, which alters the hemoglobin molecule's structural and functional properties (Turpin *et al.*, 2020). In addition to changing the mechanical properties and internal viscosity of RBCs, hyperglycemia also promotes aggregation and osmotic fragility. It also causes proteins to become glycated. The structure and hemodynamic properties of the erythrocyte are affected by these processes (Juhi, 2015; Li and Yang, 2018). These changes could be reflected in any or all of the red blood cell analytical measures, such as RBC count, HGB, HCT, MCV, MCH, and RDW.

A measurement of the size diversity of circulating erythrocytes is the red cell distribution width (RDW). In the differential diagnosis of macrocytic anemia, it is mostly utilized as an auxiliary index. A high RDW is linked to anemia caused by iron deficiency. According to a study, whether their RDW is higher or even within the normal reference range, middle-aged and older persons have a noticeably elevated risk of cardiovascular disease (CVD) events. (Anderson *et al.*, 2007; Ani and Ovbiagele, 2009; Patel *et al.*, 2009; Perlstein *et al.*, 2007)

Materials and methods

2.1 Study Design

Patients who sought treatment at a private hospital participated in this cross-sectional study.

Comment [A2]: Well written but the last paragraphs that states about red blood cell indices are miss-placed. I suggest if written before the problem statement/justification

Comment [A3]: ✓ The method is too shallow and not informative as a body/blueprint of a research article.
✓ The study design is not clear. You said it is cross-sectional/prevalence study but you tried to compare interaction among diabetic patients (cases) and diabetes-free (controls) which seems like a case control study design. Matched with age.
✓ Nothing is written or some process of sample size determination is not stated. How do you find a sample size of 100 and how do you categorize as 50:50 (DM vs Non-DM)?
✓ Study population is still unclear.
✓ Sampling technique is not stated and there is no single sentence stated, how did you select the study subjects and how do you assure data quality?
✓ Method of analysis and data processing are not clearly stated or not stated at all.
✓ Which statistical software is used to run the analysis, both descriptive and analytical part. In addition, what statistical model was used to identify the interaction of RBC indices and Diabetes as well as to identify some basic contributing factors?

2.2 Study Area

At Port Harcourt, the state capital of Rivers State, a private hospital hosted this facility-based cross-sectional study. With a population of 1,865,000 and a location along the Bonny River, Port Harcourt is a metropolis in Nigeria's South-South geopolitical zone that is sometimes frequently referred to as the Niger Delta.

2.3 Study Population

In total, 100 male and female participants who were 50 years of age or older were included in this study. The other 50 test subjects, who functioned as controls and did not have diabetes mellitus, whereas the 50 others were diabetic patients.

2.4 Collection of Blood Samples, Storage and Transportation

Following the directions of Chesbrough's (Cheesbrough, 2010) blood samples were drawn from the veins using a vacutainer tube and needle and placed in EDTA anticoagulated bottles with clear labels. Following their immediate transfer to the lab, these samples were kept at 4°C until testing using a triple packing method.

2.5 Methodology

2.5.1 ABO Blood Grouping

Method: Grouping Cells Using Tiles

A white, flawless tile was divided into sections. In compartments 1, 2, and 3, one drop of anti-A serum, one drop each of anti-B and anti-AB serums, and one drop of the patient's serum was added (auto control). A single drop of 20% patient red blood cells that had been properly blended with an applicator stick was injected into each of the 4 compartments. After gently moving the tile, it was permitted to stand for three to five minutes. Within five minutes, the agglutination on the tile was studied macroscopically and microscopically, with the results being read and recorded.

3 RESULTS

Comment [A4]: Only frequencies of some variables are mentioned and are not in line with the title and problem statement and the objective. Your objective is not met per the result written. It shall be well narrated in text form but not.

3.1 Demographic Details of Study Participants

The study included 100 individuals in total. Those with diabetes were 50, subjects without diabetes were 50. Males without diabetes were 25, while those with diabetes were 27. Females without diabetes were 28, while those with diabetes were 23. Details are shown in Table.1.

3.2 Frequency Distribution of on ABO Blood Group in Diabetic and Non Diabetic Subjects.

Table.2 revealed that blood group O had the greatest incidence of diabetes mellitus 25(50), followed in descending order by blood groups A 12(24), B 8(16), and AB 5 (10).

Table.1 Demographic Details of Study Participants

Parameters	Number
Total Number of Subjects	100
Total Number of Diabetic Subjects	50
Total Number of Non-Diabetic Subjects	50
Total Number of Diabetic Males	27
Total Number of Diabetic Females	23
Total Number of Diabetics with Blood Group A	25
Total Number of Diabetics with Blood Group B	17
Total Number of Diabetics with Blood Group AB	11
Total Number of Diabetics with Blood Group O	47

Blood Group	Diabetic		Non-Diabetic		Total	Total
	Frequency	Percentage (%)	Frequency	Percentage (%)	Frequency	Percent (%)

A	12	12	13	13	25	25
B	8	8	9	9	17	17
AB	5	5	6	6	11	11
O	25	25	22	22	47	47

Table.2: Frequency Distribution of on ABO Blood Group in Diabetic and Non Diabetic Subjects.

DISCUSSION

Numerous researchers have attempted to find evidence of a connection between diabetics and specific ABO groupings. Variable, inconsistent, and regionally distinct outcomes have been obtained. Although some researchers have shown a link between blood types and diabetes, other studies have not been able to confirm this. (Sukalingam and Ganesan, 2015; Bener and Yousafzai, 2014; Dali and Aour, 2014; Moinzadeh *et al.*, 2014).

This study's findings showed that blood group O, of which there was a total of 25 (50%), had the highest prevalence of diabetes mellitus. Blood group AB came in last with a total of 5 participants (10%), followed by blood group A with a total of 12 subjects (24%) and blood group B with a total of 8 subjects (16%) respectively. In this study, blood group O predominated over blood groups A and B (47% against 25% versus 17%, respectively). The least common blood type in this study was AB (11%). The findings of this study are at odds with a study by Waseem AG *et al.* (2012) that discovered diabetics had a higher likelihood of having blood type AB than non-diabetics. Blood types B (32%) and O (18%) showed the strongest and weakest links, respectively, with diabetes mellitus, according to a study by Zabida *et al.* (2021). There is a lot of evidence connecting certain blood types, particularly the A, AB, and Rh-positive blood groups, to the condition of diabetes mellitus, according to Sidhu *et al.*'s inquiry on the subject in 2018.

In their investigation on the relationship between ABO blood groups and diabetes mellitus, Muhammad *et al.* (2010) discovered a negative correlation between ABO blood groups A

Comment [A5]: Result is narrated in the discussion section but findings shall be discussed well in comparison with existed researches with scientific justification about the similarities and disparities.

and O and diabetes, with A and O group having decreased risks of diabetes.

When ABO blood groups are compared to diabetic populations around the world, different blood groups are distributed in different ways.

This study found that blood types O and A were the most prevalent among diabetics, and these results were in line with those of studies conducted in Japan and other Sub-Saharan African nations. Other nations including Nigeria, Algeria, Qatar, and Iraq have the highest prevalence of blood group O. While the B blood group was the most prevalent blood group in nations like Malaysia, Italy, and Trinidad (Jayawardena et al., 2012).

Racial and geographic differences may have an impact on how the disease is expressed genetically could be one explanation for the contradictory findings about the link between ABO blood types and diabetes.

5. CONCLUSION

According to the results of our study, diabetics are more likely to have the blood types O and AB. We can infer that blood type might be a risk factor and might be helpful in determining the condition.

CONSENT AND ETHICAL APPROVAL

After receiving approval from the Department of Medical Laboratory Science at Rivers State University, Port Harcourt, informed consent was obtained from patients who appeared to be in good health prior to enrollment.

REFERENCES

Doig, K. and Zhang, B. (2017). A Methodical Approach to Interpreting the Red Blood Cell Parameters of the Complete Blood Count. *Am Soc Clin Lab Sci.*, 30(3), 173–85.

- Vinupritha, P., Hariharan, M., Kathirvelu, D. and Chinnadurai, S. (2017). Estimation of HbA1C using CBC in the diagnosis of diabetes. *Asian J Pharm Clin Res.*, 10(9),214–8.
- Turpin, C., Catan, A., Guerin-Dubourg, A., Debussche, X., Bravo, S. and Álvarez, E.I. (2020) Enhanced oxidative stress and damage in glycosylated erythrocytes. *PLoS One*.
- Juhi, A. (2015). Comparative study of erythrocyte fragility in diabetes mellitus and nondiabetes Mellitus. *Int J Med Res Heal Sci.*, 4(1), 183–5.
- Li, Q. and Yang, L.Z. (2018). Hemoglobin A1c level higher than 9.05% causes a significant impairment of erythrocyte deformability in DM. *Acta Endocrinol.*,14(1), 66–75.
- Anderson, J.L., Ronnow, B.S., horne, B.D., Carlquist, J.F. and May, H.T. (2007). Usefulness of a complete blood count-derived risk score to predict incident mortality in patients with suspected cardiovascular disease. *Am J Cardiol.*, 99, 169-174.
- Ani, C. and Ovbiagele, B. (2009). elevated red blood cell distribution width predicts mortality in persons with known stroke. *J Neurol Sci.*, 277,103-108.
- Patel, K.V., Ferrucci, L., Ershler, W.B., Longo, D.L. and Guralnik, J.M. (2009). Red blood cell distribution width and the risk of death in middle-aged and older adults. *Arch Intern Med.*,169, 515-523.
- Perlstein, T.S., Weuve, J., Pfeffer, M.A. and Beckman, J.A. (2007). Red blood cell distribution width and mortality risk in a community-based prospective cohort. *Arch Intern Med.*, 169: 588-594.
- Sukalingam, K. and Ganesan, K. (2015). “Rh blood groups associated with risk to obesity and diabetes mellitus: a report on Punjabi population in Selangor, Malaysia,” *International Journal of Integrative Medical Sciences*, vol. 2, no. 4, pp. 105–109
- Bener, A. and Yousafzai, M. (2014). “The distribution of the ABO blood groups among the diabetes mellitus patients,” *Nigerian journal of clinical practice*, vol. 17, no. 5, pp. 565–568.

- Dali, S. M., Aour, M., Belmokhtar, A. F., Belmokhtar, R. and Boazza, F. (2014). "The relationship between ABO/Rh blood groups and type 2 diabetes mellitus in Maghnia, western Algeria," *South African Family Practice*, vol. 53, no. 6, pp. 568–572.
- Moinzadeh, F., Najafabady, G. M. and Toghiani, A. (2014). "Type 2 diabetes mellitus and ABO/Rh blood groups," *Journal of research in medical sciences: the official journal of Isfahan University of Medical Sciences*, vol. 19, no. 4, p. 382.
- Saeedi, P., Petersohn, I., Salpea, P., Malanda, B., Karuranga, S., Unwin, N., Colagiuri, S., Guariguata, L., Motala, A. A., Ogurtsova, K., Shaw, J. E., Bright, D., Williams, R., & IDF Diabetes Atlas Committee (2019). Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9th edition. *Diabetes research and clinical practice*, 157, 107843.
- Wokoma, F. S. (2002). Diabetes and hypertension in Africa, an overview. *Diabetes international*, 12(12), 36-40.
- Desisto, C. L., Kim, S. Y. and Sharma, A. J. (2014). Peer reviewed: Prevalence estimates of gestational diabetes mellitus in the United States, pregnancy risk assessment monitoring system (prams), 2007–2010. *The Lancet*, 11(2), 541 – 549.
- Anifowoshe AT, Owolodun OB, Akinseye KM, Iyioloa OA, Oyeyemi BF (2017) Gene frequencies of ABO and Rh blood groups in Nigeria: a review. *Egypt J Med Human Genetics* 18(3):205–210. <https://doi.org/10.1016/j.ejmhg.2016.10.004>
- Oladele EO, Yahaya TO, Adewunmi OO, David B, Oladipo AJ (2020) Distribution of ABO/rhesus blood groups among hepatitis B virus (HBV) positive patients in Lagos, South-Western, Nigeria. *FUDMA J Sci* 4(3):196–200
- Qureshi MA, Bhatti R (2003) Frequency of ABO blood groups among the diabetes mellitus type 2 patients. *J Coll Phys Surg Pak* 13(8):453–455

- Zahid H, Yahyaoui A, Uwingabiye J, El Khazraji A, El Hadeif R, Messaoudi N (2016) Phenotypic and allelic distribution of ABO and Rh-D blood Groups in Blood Transfusion Center of Avicenna Military Hospital, Marrakech, Morocco. *Int J Med Health Res* 2(1):1–9
- Abdel Ghafar MT, Shalaby KH, Okda HI, Rizk and F.H. (2020) Association of ABCA1 (C69T) gene polymorphism with dyslipidemia and type 2 diabetes among the Egyptian population. *Meta Gene* 25:100714. <https://doi.org/10.1016/j.mgene.2020.100714>
- Yahaya T, Oladele E, Fatodu I, Abdulazeez A, Yeldu Y (2020) Concentrations and health risk evaluation of heavy metals and microorganisms in groundwater in Lagos, Southwest Nigeria. *Journal of Advances in Environmental Health Research* 8(3):234–242. <https://doi.org/10.22102/jaehr.2020.245629.1183>
- Yahaya T, Obaroh IO, Oladele EO (2017) The roles of environmental pollutants in the pathogenesis and prevalence of diabetes: a review. *J Appl Sci Environ Manag* 21(1):5–8. <https://doi.org/10.4314/jasem.v21i1.1>
- Shittu RO, Kasali FO, Biliaminu SA, Odeigah LO, Sule AG, Musah Y (2017) Prevalence of diabetes and pre-diabetes in Oke-Ogun Region of Oyo State, Nigeria. *J Med Rese Health Educ* 1:1