

Original Research Article

Robust NIRS models for non-destructive prediction of physicochemical properties and ageing of basmati rice

Abstract:

Aim: To determine physicochemical properties and age of rice by non-destructive technique.

Place and duration of study: Study was conducted at Division of Food Science and Postharvest Technology, Indian Agricultural Research Institute, New Delhi during 2020 to 2021.

Methodology: Rice were kept for accelerated aging at 42.6°C temperature & 71% RH for a duration of 30 days. Changes in four physicochemical properties namely amylose content, volume expansion ratio (VER), water absorption ratio (WAR), and kernel elongation ratio (KER) were evaluated destructively (by spectrophotometer and cooking method) and non-destructively (by spectroradiometer) at every alternate day, during 30 days storage.

Results: The physicochemical parameters of rice showed a good correlation with spectral signatures. Subsequently, Principal component Analysis (PCA), Partial Least Square Regression (PLSR), and Multiple Linear Regression (MLR) were used to model the physicochemical changes occurring during the process of accelerated aging using spectral reflectance values. Based on values of Coefficient of determination (R^2) and Root mean square error (RMSE) accuracy of models was determined. Predictions with the MLR model resulted in a coefficient of determination (R^2) of 0.82, 0.87, 0.9, 0.7, 0.83 and 0.82 with root mean square error (RMSE) of 0.18, 0.13, 0.21, 0.124 and 4.2 for amylose content, VER, WAR, KER, and ageing process respectively for calibration.

Conclusion: The study demonstrated the potential of NIRS in non-destructively predicting the physicochemical parameters of rice.

Keywords: Accelerated Ageing; Basmati rice; Non-destructive; Reflectance; Robust; Spectroradiometer.

Abbreviations:

MLR	Multiple Linear Regression
PLSR	Partial Least Square Regression
WAR	Water Absorption Ratio
VAR	Volume Expansion Ratio
KER	Kernel Elongation Ratio
PB	Pusa Basmati
PCA	Principal Component Analysis

1. INTRODUCTION

Rice (*Oryza sativa* L.) is a global staple food, especially in South Asia, with India as a major producer after China. It is primarily grown during the kharif season (June to September) but produced year-round in select regions. Rice consumption is significant in Arabian and South Asian countries, comprising carbohydrates, starch, and various nutrients. Basmati, a fragrant rice variety, is crucial in India, leading in production and export, serving as a significant crop for the country's economy (Kaur & Chand, 2011).

Rice undergoes beneficial aging after harvesting, resulting in changes in physicochemical properties. Aging criteria are established based on physical and chemical properties, but the exact mechanism remains partly understood. Natural aging involves storing rice at room temperature for 3-6 months, a time-consuming process. Accelerated aging, where rice is stored at elevated temperatures for minutes to days, offers a more efficient alternative. Temperature, humidity, and storage duration are key factors impacting rice quality during aging (Hussain et al., 2021).

Aged rice is preferred by Indian and Arabian consumers for its improved texture and premium qualities, especially in the case of basmati rice known for its aroma, long grains, and excellent cooking properties. Accurate assessment of rice aging is crucial to maintain quality and prevent fraudulent practices. Traditional methods for assessing rice aging are complex, time-consuming, and labor-intensive, requiring expensive equipment. Modern food industries demand quick, portable, and non-destructive techniques for quality determination (El-Mesery et al., 2019).

Visible near-infrared (VNIR) spectroscopy is a valuable non-destructive technique widely used across industries like agriculture, cosmetics, food, polymers, pharmaceuticals, and textiles due to its speed, cost-efficiency, and minimal labor requirements (Yang et al., 2005). VNIR spectroscopy can assess commodity quality in the field and during processing, relying on the scattering and absorption of light. Scattering reflects external properties like particle size and density, while absorption reveals the chemical composition (Birth et al., 1985).

VNIR spectroscopy is most successfully used for variety discrimination, determination of moisture content, pH, acidity, Various sugars, various diseases/rots, and evaluation of internal quality (Qin & Lu, 2007). VNIR spectroscopy works in the wavelength range between 350- 2500 nm. VNIR spectrometer takes signature in terms of reflectance, absorbance, and transmittance (Jha & Garg, 2010). VNIR spectroscopy is used for non-destructive evaluation of rice i.e amylose content of milled rice (Villareal et al., 1994; Delwiche et al., 1995), quality of rice starch (Bao et al., 2001), textural properties of rice (Windham et al., 1997), quality characteristics of rice (Delwiche et al., 1996), evaluation of rice wine in terms of soluble solid and pH (Liu et al., 2007).

This experiment utilized VNIR spectroscopy in reflectance mode to assess the aging of various basmati rice varieties. The spectroradiometer's advantages include easy monitoring of the aging process and improved rice grading based on age, potentially increasing market value. The study aims to non-destructively analyze rice age using a VNIR spectroradiometer, with objectives to characterize artificially aged rice grains through spectroscopy and physical parameters and to develop models for discriminating and estimating aging in different rice grains.

2. MATERIAL AND METHODS

2.1. Procurement of paddy

Freshly harvested four varieties of basmati rice [Pusa Basmati (PB)- 1121, PB-1509, PB- 1637, PB-1718] and one non-basmati variety [*Pusa Sugandh* (PS)] were procured from Seed Production Unit, ICAR- Indian Agricultural Research Institute, New Delhi. Which was grown in the *kharif* season (June to October).

2.2. Sample preparation

All five varieties of paddy namely PB- 1121, PB-1509, PB- 1637, PB- 1718, and PS were hulled in Satake rice huller (Satake Japan). After hulling brown rice fed to a Satake rice polisher for polishing up to 6%. Broken rice was separated and only head rice was used for further work.

2.3. Accelerated ageing of rice

For accelerated ageing freshly harvested milled rice of varieties PB 1121, PB 1509, PB 1718, PB 1637, and Pusa Sugandh were kept in a controlled chamber maintained at Relative humidity (RH) of 71 %, Temperature of 42.6 °C for a period of 30 days as suggested by Rayguru et. al. (2011) at National Phytotron Facility, IARI New Delhi.

2.4. Visible Near-infrared spectroscopy analysis

Spectral signatures of fresh and accelerated aged samples (kept in a petri dish up to 1 cm thickness) were acquired before the destructive analysis in the wavelength range between 350- 2500 nm at 1 nm

intervals using a handheld spectroradiometer [Analytical Spectral Device (ASD) Fieldspec® Spectroradiometer (350 to 2500 nm), Boulder, USA] at Hyperspectral Remote Sensing Laboratory, IARI, New Delhi. Care was taken to calibrate the device with a standard white plate before acquiring the spectral signatures. The precaution was also taken to ensure no gaps between grains to avoid signal losses. For each variety and accelerated storage period, six spectra were acquired to ensure the repeatability of the spectral signatures. A device was calibrated after every six spectra.

2.5. Determination of physicochemical properties of rice

2.5.1. Amylose content

Amylose content in fresh and accelerated aged rice was determined using the spectrophotometric method suggested by Juliano (1971). 100 mg rice grains were grounded using mortar and pestle and transferred into the 100 ml volumetric flask. To it, 1 ml of ethanol and 10 ml of NaOH were added. The mixture was heated in a boiling water bath for 10 min. Distilled water was subsequently added to make the volume up to 100 ml. Out of 100 ml only 2.5 ml of the solution was then transferred into the 100 ml flask and 20 ml distilled water was added to it along with 3-4 drops of phenolphthalein indicator till it turns pink. Then 0.1 N HCl was added drop by drop until the pink color disappears. At last, 1 ml of iodine was added to it and the volume was made up to 50 ml using distilled water. The absorbance of the solution was measured at 510 nm and the amylose content was determined using a standard curve.

2.5.2. Volume expansion ratio

Volume expansion ratio (VER) was measured by using the toluene displacement method (Sindhu et al., 1975). A specific volume of toluene was added to 250 ml of measuring cylinder then put 10 grains of uncooked rice and the initial reading was noted. Similarly, a change in the volume of 10 grains after cooking was also recorded.

$$VER = \frac{V_c}{V_{uc}}$$

Where,

V_c = Volume cooked rice

V_{uc} = Volume of uncooked rice

2.5.3. Water absorption ratio

Take 10 g of milled head rice and put it into 30 g of distilled water. Then beaker was kept in a boiling water bath (97 ± 2 °C) for cooking. After cooking take the final weight of the rice (Nayak & Mohapatra, 2019).

$$WAR = \frac{W_c - W_{uc}}{W_{uc}}$$

Where,

W_c = Weight of cooked rice

W_{uc} = Weight of uncooked rice

2.5.4. Kernel elongation ratio

It is the ratio of the average length of grain after cooking to them before cooking. 10 grains of head rice were randomly selected from the sample and measured the length using a micrometer. Then grains were kept for cooking and measured for the final length (Azeez and Shafi, 1966).

$$KER = \frac{X_c}{X_{uc}}$$

Where,

X_c = Length of cooked rice

Xuc= Length of uncooked rice

2.6. Chemometric analysis and selection of wavelength

For performing chemometric analysis different mathematical calculations and statistical methods were used. In chemometric analysis operations such as correlation, regression, calculation of the first derivative, differentiation of wavelength, development of model, and evaluation of model were done as described in the research work of various researchers (Nordey et al., 2017).

The data available after the chemometrics was too large. The next step is to reduce the huge data in a particular manner and process only into selected data. Visible near-infrared (VNIR) spectroscopy wavelength ranges from 350 to 2500 nm were used in this study. The Visible Near-infrared reflectance spectra of rice samples shown in **Fig. 1**. In this huge range of wavelengths for each parameter, it is essential to select particular ranges of wavelengths because there are close relationships that can occur between wavelength and parameters. For the selection of wavelengths, correlation of physicochemical parameters namely amylose content, volume expansion ratio, water absorption ratio, and kernel elongation ratio with raw spectra as well as with the first derivative of reflectance were performed. These correlations were done by correlating four parameters with spectral reflectance data ranging from (351 to 2500 nm). As the correlation value ranges between -1 to 1, wavelengths showing high values of correlation were used for further analysis.




Fig. 1. Spectral signature of Pusa Basmati 1121 (PB 1121) after different periods of accelerated storage

2.7. Multivariate analysis

Multivariate analysis of the data was conducted to develop robust models for predicting the physicochemical parameters at different levels of accelerated aging and to predict the level of accelerated aging of rice. Spectral reflectance values acquired at all identified promising wavelengths were subjected to different available multivariate analyses like Partial least square regression (PLSR), principal component analysis (PCA), and multiple linear regression (MLR). Of the data used for multivariate analysis, 70% were utilized for model calibration, and the rest 30% for model validation. Maximum R^2 and minimum RMSE were considered for model selection.

3. RESULTS AND DISCUSSION

3.1. Amylose content

The initial amylose content of rice varieties varied, with Pusa Sugandh at 22.21%, and among basmati types, PB 1718 had the highest at 23.19%, followed by PB 1509 (23.12%), PB 1121 (22.79%), and PB 1637 (22.60%). During 30 days of accelerated storage at 42.6°C, all varieties showed an increase in amylose content, with PB 1509 having the least change. This increase was

most significant during the first 14 days, equivalent to 7 months of ambient storage. The elevated temperature likely caused amylopectin to convert into amylose, as observed in previous studies (Zhong et al., 2020).

Amylose content significantly influences rice cooking quality. Higher amylose levels in aged rice improve cooking quality and processing efficiency by reducing stickiness and increasing grain firmness (Thanathornvarakul et al., 2016). This is due to lower leaching of solids during cooking (Li et al., 2017) and its positive correlation with water absorption, volume expansion, fluffiness, and grain separation. In the presence of lipids, amylose acts as both a diluent and a swelling inhibitor (Zhou et al., 2002).


3.2. Volume expansion ratio (VER)

Volume expansion ratio (VER) is crucial for rice cooking quality. During accelerated storage, all varieties showed increased VER. Pusa Sugandh had the lowest initial VER at 3.15, while basmati varieties ranged from 3.51 to 3.77. After 30 days, VER increased the most for PB 1509 (4.69), followed by PB 1718 (4.65), PB 1637 (4.46), PB 1121 (4.41), and Pusa Sugandh (4.20). This trend aligns with previous findings regarding rice aging and amylopectin content reduction, affecting pasting properties and increasing VER (Thanathornvarakul et al., 2016).


Accelerated storage induces structural changes in amylose chains, resulting in harder grains and higher Volume Expansion Ratio (VER). Consequently, cooked rice becomes firmer and less sticky due to increased water absorption and VER. Study by Indiartho&Nurannisa (2021) also support the observed increase in VER during rice aging.

3.3. Water Absorption Ratio (WAR)


Water Absorption Ratio (WAR), an important cooking characteristic, mirrored the VER trend, as both are positively correlated due to increased water absorption contributing to higher volume expansion during cooking. Initial WAR values varied: PB 1121 (3.44), PB 1509 (3.33), PB 1637 (3.39), PB 1718 (3.52), and Pusa Sugandh (3.33). After 30 days of accelerated storage, WAR increased for all varieties, with PB 1121 at 3.75, PB 1509 at 3.67, PB 1637 at 3.66, PB 1718 at 3.75, and Pusa Sugandh at 3.59. Pusa Sugandh exhibited the least increase. Similar trends were reported by Zhou et al. (2015) where aging led to increased water absorption due to rice hardness and amylose-lipid complex formation.




(a)



(b)



(c)



(d)

Fig. 2 Physiochemical changes in rice during ageing a) amylose content, b) VER, c) WAR, d) KER

3.4. Kernel elongation ratio (KER)

Kernel Elongation Ratio (KER) is vital for understanding rice cooking and eating quality. PB 1121 had the highest KER at 1.62, significantly surpassing all other varieties, while Pusa Sugandh had the lowest at 1.39. Among basmati types, PB 1509, PB 1637, and PB 1718 had KER values of 1.45, 1.46, and 1.49, respectively. KER generally increased during 30 days of accelerated storage, reaching 2.32 (PB 1121), 2.25 (PB 1509), 2.17 (PB 1637), 2.31 (PB 1718), and 2.20 (Pusa Sugandh) on the 30th day. The aging effect on KER followed this order: Pusa Sugandh (58.27%), PB 1509 (55.17%), PB 1718 (55.03%), PB 1637 (48.63%), and PB 1121 (43.21%). These results align with findings from Indiartho&Nurannisa (2021).

Variability in elongation ratio is linked to amylose content, particularly in PB 1121 and PB 1509, possibly due to amylose structure and its interaction with protein and lipids during accelerated storage. Starch granule expansion without disintegration depends on amylose content (Juliano, 1985), aligning with VER and WAR trends governed by structural changes and amylose content. Aging efficiency varies with rice variety, pre-treatments, techniques, and storage conditions (Indiartho&Nurannisa, 2021).

3.5 Multivariate analysis and modeling

Spectral signatures were correlated with physicochemical parameters and aging duration, yielding correlation coefficients from -0.1 to 0.85 in the 400-1500 nm wavelength range. Correlation of raw spectral signatures with physicochemical parameters spanned wide wavelength ranges, making wavelength selection challenging. To address this, mathematical transformations, specifically 1st derivatives of spectral reflectance values, were employed. Correlation analysis revealed wavelength bands of interest: 1500-1800 nm for amylose content, 1200-1800 nm for VER, 1100-1800 nm for WAR, 500-1300 nm for KER, and 950-1400 nm for accelerated aging duration. For KER, correlation coefficients surpassed ± 0.8 at 1150-1350 nm. The derivatives demonstrated higher maximum correlation coefficients compared to raw spectral reflectance values (Table 1).

Parameter	Max. correlation	Wavelength range (nm)
Amylose content	0.7917	1500-1800
VER	0.7982	1200-1800
WAR	0.7630	1100-1800
KER	0.8463	500-1300

Table 1 Wavelengths showing maximum correlation with first derivative spectral reflectance

3.5.1 Amylose content


Amylose content changes during rice accelerated aging, affecting grain texture and cooking properties. PLSR models were developed using data in the 1500-1800 nm wavelength range, resulting in R^2 values of 0.65 (calibration) and 0.90 (validation), with an RMSE of 0.23. PCA showed that the first three principal components explained 94% of the variability. Subsequently, MLR modeling using sensitive wavelengths in the 1500-1800 nm range yielded R^2 values of 0.82 (calibration) and 0.84 (validation) with RMSE values of 0.186 and 0.210, respectively (Fig. 3a).

Matsuo et al. (2018) employed NIR spectroscopy, achieving an R^2 of 0.72 for amylose content in Japonica rice within the 850-1048 nm range. Fernández-Novales et al. (2009) determined reducing sugars in grape ripening using Shortwave-NIR spectroscopy, obtaining an R^2 of 0.92 (800-1050 nm). Bao et al. (2001) predicted rice starch quality with a spectroradiometer, yielding an R^2 of approximately 0.91 for amylose content (400-2500 nm). He et al. (2021) used NIR spectroscopy for cereals, attaining R^2 values above 0.9 in the 1923-1961 nm range for starch and amylose content.

3.5.2 Volume Expansion Ratio

Similar to the amylose content approach, VER was correlated with spectral signatures. The first three principal components explained 94.6% of VER variability, with loading values peaking between 1200-1800 nm. PLSR models had R^2 values of 0.50 (calibration) and 0.67 (validation), with RMSE of 0.21. MLR models yielded R^2 values of 0.87 (calibration) and 0.85 (validation), with RMSE values of 0.13 and 0.15, respectively (Fig. 3b).

Moghimi et al., (2010) attempted to develop calibration models for evaluating the TSS and acidity of kiwifruit. They developed a model using principal component analysis (PCA) and partial least square regression (PLS). The correlation coefficients for TSS and acidity were 0.93 and 0.943 respectively. RMSE values were 0.076% and 0.26°Brix respectively obtained between wavelength region of 400 to 1000 nm.






Fig. 3 MLR models for prediction of a) amylose content, b) VER, c) WAR, d) KER, e) ageing

3.5.3 Water expansion ratio

For water expansion ratio, the first three principal components explained 93.6% of the variability, with loading values peaking at 1100-1800 nm. PLSR models achieved R^2 values of 0.78 (calibration) and 0.70 (validation), with RMSE of 0.044. MLR models obtained R^2 values of 0.97 (calibration) and 0.90 (validation), with RMSE values of 0.21 and 0.25, respectively (Fig. 3c). In a study by Kusumiyati et al. (2021) on apple fruit quality evaluation, NIR spectroradiometer yielded an R^2 of 0.81 and RMSE of 0.009 for water content within the 702-1065 nm wavelength band.

3.5.4 Kernel elongation ratio

For kernel elongation ratio (KER), the first three principal components explained 96% of the variability, with loading values peaking at 500-1300 nm wavelengths. PLSR models achieved R^2 values of 0.64 (calibration) and 0.84 (validation), with RMSE of 0.16 and 0.15. MLR models obtained R^2 values of 0.83 (calibration) and 0.85 (validation), with RMSE values of 0.124 and 0.11, respectively (Fig. 3d). In a study by Schmilovitch et al. (2000) on mango physiological indices using NIR spectroscopy, MLR models had R^2 values of 0.92 for TSS and 0.6085 for acidity in the 1200-2400 nm wavelength region, outperforming PLSR models.

3.5.5 Ageing of rice

For accelerated aging, spectral signatures were correlated similarly to amylose content. The first three principal components explained 96% of the variability. PLSR models achieved R^2 values of 0.42 (calibration) and 0.58 (validation), with RMSE of 6.27. MLR models obtained R^2 values of 0.82 (calibration) and 0.70 (validation), with RMSE values of 4.2 and 4.5, respectively (Figure 3e). In a study by Fernández-Navales et al. (2009) on wine quality prediction during aging using NIR spectroscopy, it was found to be a promising technique for assessing grape wine quality attributes during fermentation and aging.

The chemometric analysis indicated that spectral reflectance values between 350 to 2500 nm could adequately predict quality. MLR outperformed PLSR in terms of R^2 and RMSE in both calibration and validation. MLR's superior performance is attributed to its selective wavelength choice, eliminating overfitting and collinearity issues observed in PLSR models (Table 2) (EIMasry et al., 2007; Fernández-Navales et al., 2009).

Parameters	MLR Model				PLSR Model			
	Calibration		Validation		Calibration		Validation	
	R^2	RMSE	R^2	RMSE	R^2	RMSE	R^2	RMSE
Amylose content	0.82	0.18	0.84	0.21	0.64	0.23	0.90	0.23
VER	0.87	0.13	0.85	0.15	0.50	0.21	0.67	0.21

WAR	0.97	0.21	0.90	0.25	0.77	0.04	0.70	0.04
KER	0.83	0.124	0.85	0.11	0.63	0.15	0.84	0.15
Ageing	0.82	4.2	0.70	4.5	0.42	6.27	0.57	6.27

Table 2 Summary of PLSR and MLR statistics for Amylose content, VER, WAR, KER, and Ageing

4 CONCLUSION

Physicochemical changes during accelerated rice aging were assessed both destructively and non-destructively. Spectral reflectance, including first derivatives, correlated with these changes and aging periods. PCA, PLSR, and MLR were used to develop predictive models.

- Spectral reflectance effectively captures amylose content, VER, WAR, and KER changes during rice aging, enabling age determination.
- Variabilities in physical parameters and aging were characterized within 600-1800 nm using raw spectral reflectance, and 900-1350 nm using 1st derivatives.
- MLR models outperformed PLSR models, achieving $R^2 > 0.80$ for physical parameters and age prediction (PLSR R^2 : 0.42-0.77).

References

- Juliano, B. O. A simplified assay for milled-rice amylose. *Cereal Sci. Today*. 1971; 12: 334-360.
- Sidhu JS, Gill M S, Bains GS. Milling of paddy in relation to yield and quality of rice of different Indian varieties. *J. Agric. Food Chem.* 1975; 23(6): 1183-1185. <https://doi.org/10.1021/jf60202a035>
- Azeez MA, Shafi M, Quality in Rice. Department of Agriculture, West Pakistan Technology. 1966; Bulletin. No. 13: 50.
- Bao JS, Cai, YZ, Corke H, Prediction of Rice Starch Quality Parameters by Near-Infrared Reflectance Spectroscopy. *J. Food Sci.* 2011, 66(7): 936–939. <https://doi.org/10.1111/j.1365-2621.2001.tb08215.x>
- Birth G S, Dull GG, Renfroe WT, Kays SJ, Nondestructive Spectrophotometric Determination of Dry Matter in Onions. *J Am. Soc. Hortic. Sci.* 1985, 110(2): 297–303. <https://doi.org/10.21273/JASHS.110.2.297>
- Delwiche SR, Bean MM, Miller RE, Webb BD, Williams PC, Apparent amylose content of milled rice by near-infrared reflectance spectrophotometry. 1995
- Delwiche SR, McKENZIE KS, Webb BD, Quality characteristics in rice by near-infrared reflectance analysis of whole-grain milled samples. 1996
- EIMasry G, Wang N, EISayed A, Ngadi M, Hyperspectral imaging for nondestructive determination of some quality attributes for strawberry. *J. Food Eng.* 2017, 81(1): 98–107. <https://doi.org/10.1016/j.jfoodeng.2006.10.016>
- El-Mesery H, Mao H, Abomohra A, Applications of Non-destructive Technologies for Agricultural and Food Products Quality Inspection. *Sensors* 2019, 19(4): 846. <https://doi.org/10.3390/s19040846>
- Fernández-Novales J, López MI, Sánchez MT, Morales J, González-Caballero V, Shortwave-near infrared spectroscopy for determination of reducing sugar content during grape ripening, winemaking, and aging of white and red wines. *Food Res. Int.* 2009, 2(2): 285–291. <https://doi.org/10.1016/j.foodres.2008.11.008>
- He, M, Hu J, Wu Y, Ouyang J, Determination of starch and amylose contents in various cereals using common model of near-infrared reflectance spectroscopy. *Int. Food Res. J.* 2021, 28(5): 987–995. <https://doi.org/10.47836/ifrj.28.5.12>
- Hussain, S. Z., Iftikhar, F., Naseer, B., Altaf, U., Reshi, M., & Nidoni, U. K., 2021. Effect of radiofrequency induced accelerated ageing on physico-chemical, cooking, pasting and textural properties of rice. *LWT* 139: 110595. <https://doi.org/10.1016/j.lwt.2020.110595>

- Indiarto, R., & Nurannisa, R. L., 2021. Aging Technique Of Rice And Its Characteristics. 10(01).
- Jha SN, Garg R, Non-destructive prediction of quality of intact apple using near infrared spectroscopy. *J. Food Sci. Technol.* 2010, 47(2): 207–213.
- Kaur J, Chand T, Effect of Natural Ageing of Basmati Paddy on Physico-chemical and Textural Properties of Cooked Rice. *J. Agric. Eng.* 2011, 48(2): 1–6.
- Kusumiyati K, Hadiwijaya Y, Suhandy D, Munawar AA, Prediction of water content and soluble solids content of 'manalagi' apples using near infrared spectroscopy. *IOP Conference Series: Earth and Environmental Science* 2021, 922(1): 012062. <https://doi.org/10.1088/1755-1315/922/1/012062>
- Li H, Fitzgerald MA, Prakash S, Nicholson TM, Gilbert R G, The molecular structural features controlling stickiness in cooked rice, a major palatability determinant. *Sci. Rep.* 2017, 7(1): 1–12.
- Liu F, He Y, Wang L, Pan H, Feasibility of the use of visible and near infrared spectroscopy to assess soluble solids content and pH of rice wines. *J Food Eng.* 2007, 83(3): 430–435. <https://doi.org/10.1016/j.jfoodeng.2007.03.035>
- Moghimi A, Aghkhani MH, Sazgarnia A, Sarmad M, Vis/NIR spectroscopy and chemometrics for the prediction of soluble solids content and acidity (pH) of kiwifruit. *Biosys. Eng.* 2010, 106(3): 295–302. <https://doi.org/10.1016/j.biosystemseng.2010.04.002>
- Nayak M, Mohapatra M, Effect of microwave heating on accelerated aging qualities of rice. *Oryza*, 2019, 56(2), 228–235.
- Nordey T, Joas J, Davrieux F, Chillet M, Léchaudel M, Robust NIRS models for non-destructive prediction of mango internal quality. *Sci. Hortic.* 2017, 216, 51–57. <https://doi.org/10.1016/j.scienta.2016.12.023>
- Qin J, Lu R, Measurement of the absorption and scattering properties of turbid liquid foods using hyperspectral imaging. *Applied Spectroscopy* 2007, 61(4): 388–396.
- Schmilovitch Z, Mizrach A, Hoffman A, Egozi, H, Fuchs Y, Determination of mango physiological indices by near-infrared spectrometry. *Postharvest Biol. Technol.* 2000, 19(3): 245–252. [https://doi.org/10.1016/S0925-5214\(00\)00102-2](https://doi.org/10.1016/S0925-5214(00)00102-2)
- Thanathornvarakul N, Anuntagool J, Tananuwong K, Aging of low and high amylose rice at elevated temperature: Mechanism and predictive modeling. *J. Cereal Sci.* 2016, 70: 155–163.
- Villareal CP, De La Cruz NM, Juliano BO, Rice amylose analysis by near-infrared transmittance spectroscopy. *Cereal Chem.* 1994, 71(3): 292–296.
- Yang CC, Chao K, Chen YR, Early HL, Systemically diseased chicken identification using multispectral images and region of interest analysis. *Comput. Electron. Agric.* 2005, 49(2): 255–271.
- Zhou Z, Robards K, Helliwell S, Blanchard C, Ageing of Stored Rice: Changes in Chemical and Physical Attributes. *J Cereal Sci.* 2002, 35(1): 65–78. <https://doi.org/10.1006/jcrs.2001.0418>
- Zhou Z, Wang X, Si X, Blanchard C, Strappe P, The ageing mechanism of stored rice: A concept model from the past to the present. *J. Stored Prod. Res.* 2015, 4: 80–87. <https://doi.org/10.1016/j.jspr.2015.09.004>