

Original Research Article

Influence of growth regulator and organic manure on the cultivation of baby corn

(*Zea mays L.*)

Formatted: Left

ABSTRACT

The experiment comprised of ten treatments of different combinations of organic and plant growth regulator nutrient management replicated thrice in a Randomized Block Design. The main objective of the experiment experiment's main objective was to evaluate the Influence of organic manure and plant growth regulator on the growth and yield of baby corn (*Zea mays L.*) Prayagraj condition. The three levels of of organic manure and plant growth regulator are FYM (10t/ha) and NAA (30ppm), FYM(10t/ha) and GA₃(15.47ppm), FYM (10t/ha) and Seaweed sap (Ascophyllum nodosum)5% from the present investigation it may be concluded that the profitable production of baby corn can be secured by FYM (10t/ha) and NAA (30ppm) + FYM(10t/ha) and GA₃(15.47ppm) (T₆).

Keywords: Baby corn, organic manure, growth regulator, yield and quality

Introduction

Maize(*ZeamaysL*)isthemostversatileandemergingcropwitha widerangeofadaptabilityunderdifferentagro-climaticconditions.Acrossconditions. Acrosstheglobe,itisfamousasthe queenofcerealssinceitpossesspossesseshighergeneticyield potential amongthecereal cropsas compared to likerice,wheat,oat,milletsetc. Inmostofthe developing countries,maizecontributesa majorshare foroffoodsecurity.InIndia,maizeisthethirdmostimportantcropfollowedbyriceandwheat. Itssignificanceliesinthewaythatitis not onlyutilizedforhumanconsumptionandanimal feedbutalsoitisutilizedbytheindustriesfortheproductionofcornstarch,cornoiletc. CountrieslikeThailandandTaiwanachievedsuccessfulresultsduetothe cultivationofbabycorn. Latermoreattentionisgive nonthecultivationofmaizebytheresearchersandagriculturiststotapitspotentialities forearningmoreforeignrevenueinadditiontogetmaximumreturnstothe producers. Babycornisnotageneticallydwarfmaizeasthenamesuggests,it Istheimatureearofnormalmaize. Maturedmaizeearare too hard so,can'tbeusedasvegetables. Whereasbabycorneararesoftandconsumedasvegetablebyhumanbeing (Jinjalaetal.,2016). Theyoungbabycornwasharvestedduringsilkingstage. Afterharvesting,theexternal sheathwasremovedandtheearwasusedforvegetablepurposeviz.,salad,soup,picklesetc(Muthukumar etal.,2005). Babycornisadelicious, decorative,lowcaloricnutritious vegetablewithout cholesterolandisrichinfibrecontent. ItisMaize(ZeamaysL.)isthemostversatileandemergingcropwith widerangeofadaptabilityunderdifferentagro-climaticconditions.Acrosstheglobeitisfamousasqueenofcerealsinceitpossesshighergeneticyield

Comment [DAS1]: This sentence shows no meaning here and seems useless.

Comment [DAS2]: In the Introduction part there are lots of punctuation mistakes, which are needed to be corrected. There should be a single space between full stop and starting of the next line. There should also a single space between comma and next word.

potential among other cereal crops like rice, wheat, oat, millets etc. In most of the developing countries maize contributes major share for food security. In India, maize is the third most important crop followed by rice and wheat. Its significance lies in the way that it is not only utilized for human consumption and animal feed, but also it is utilized by the industries for the production of corn starch, corn oil etc. Countries like Thailand and Taiwan achieved successful results due to the cultivation of baby corn. Later more attention is given on the cultivation of maize by the researchers and agriculturists to tap its potentialities for earning more foreign revenue in addition to get maximum return to the producers. Baby corn is not genetically dwarf maize as the name suggests, it is the immature ear of normal maize. Matured maize ears are too hard so, can't be used as vegetables. Whereas baby corn ears are soft and consumed as vegetable by human being (Jinjala *et al.*, 2016). The young baby corn was harvested during silking stage. After harvesting, the external sheath was removed and the ear was used for vegetable purpose viz., salad, soup, pickles etc (Muthukumar *et al.*, 2005).

Comment [DAS3]: These sentences are pasted twice in introduction and seems MS has been not prepared seriously.

Baby corn is a delicious, decorative, low calorie nutritious vegetable without cholesterol and is rich in fibre content. It is free from pests and diseases and it contains protein up to 15 to 18 percent, sugar 0.016 to 0.020 percent, phosphorus 0.6 to 0.9 percent, potassium 2 to 3 percent, fibre 3 to 5 percent, calcium 0.3 to 0.5 percent and ascorbic acid 75 to 80 mg/100g. As green fodder, it is the best suited for milch chain mails since it has lactogenic properties (Reenarani *et al.*, 2017). For the past few decades increased use of synthetic fertilizer reduced the usage of organic manures which affected the soil fertility and productivity. Organic way of crop production enhanced the sustainability and soil health without affecting the ecosystem.

Formatted: No Spacing, Space Before: 0 pt, Line spacing: single, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Comment [DAS4]: Clarify it.

Formatted: Highlight

Comment [DAS5]: What are the objectives for writing this MS, is missing in it. Please add it.

Materials and Methods

The current study was carried out in the Crop Research Farm, Department of Agronomy, Naini Agriculture Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj, during the Kharif season 2022, (U.P.). The experimental field is situated on the left side of the Prayagraj-Rewa Road, about four kilometers from Prayagraj city and close to the Yamuna River, at 25.57° N latitude, 87.19° E longitude and at an altitude of 98m above mean sea level. The subtropical region of Uttar Pradesh, where Prayagraj is located, has scorching summers and nice winters. The regions typical temperature ranges from 23°C to 38°C, seldom falling below 3° C or 4°C. The relative humidity levels range from 28.57% to 95%. In this location, the average annual rainfall is 1050mm. The soil chemistry analysis revealed a sandy loam texture with a p^H of 7.20, low amounts of organic carbon (0.83 percent) and potassium (208.8kg/ha) and a low quantity of accessible phosphorous (17.2kg/ha). The soil was electrically conductive and had a conductivity of 0.34 ds/m. For each of the nine treatment combinations, three replications were employed. The treatment details and treatment combinations are shown in Table 1 and 2, respectively. Organic manure, and plant growth regulator management and crop maintained according to the treatment combinations. Plant height (cm) at harvest, dry weight at harvest, number of cobs/plant, cob length, cob girth, baby corn yield (t/ha) successfully measured, and an economic analysis of each treatment was completed to determine the best treatment combination for baby corn cultivation.

Table 1. Treatments details

Organic manure	
Vermicompost	4.46t/ha
FYM	10t/ha
Goat manure	15t/ha
Plant growth regulator	
Seaweed sap (<i>Ascophyllum nodosum</i>)	5%
GA3	15.46ppm
NAA	30ppm

Comment [DAS6]: Mention few lines about each treatment.

Formatted: Font: Italic

Comment [DAS7]: What are the basis of selecting the doses of NAA and GA3? Show the references for it.

Table 2. Treatment combination

Treatment symbol	Treatment combinations
T ₁	Vermicompost(4.46t/ha) + Seaweed sap (<i>Ascophyllum nodosum</i>) 5%
T ₂	Vermicompost(4.46t/ha) + GA ₃ (15.47ppm)
T ₃	Vermicompost(4.46t/ha) + NAA (30ppm)
T ₄	FYM (10t/ha) + Seaweed sap (<i>Ascophyllum nodosum</i>) 5%
T ₅	FYM (10t/ha) + GA ₃ (15.47ppm)
T ₆	FYM (10t/ha) + NAA (30ppm)
T ₇	Goat manure (15t/ha) + Seaweed sap (<i>Ascophyllum nodosum</i>) 5%
T ₈	Goat manure (15t/ha) + GA ₃ (15.47ppm)
T ₉	Goat manure (15t/ha) + NAA (30ppm)
T ₁₀	Control (RDF) 120:60:40: NPK

Results and Discussion

Growth parameters

Plant height (cm)

Table 3 shows organic manure, plant growth regulator nutrient management and crop spacing on plant height at harvest. The data indicated that significant impact on plant height during the crop growth period. Application of T₆- FYM (10t/ha) + NAA (~~30ppm~~30 ppm) significantly influenced the plant height of baby corn at 45DAS. The maximum plant height (189.57cm) was ~~recorded~~ recorded in T₆ FYM (10t/ha) + NAA (30ppm) which was statistically at par with T₅, T₄, T₃ and minimum plant height (162.17) was recorded in the application of T₁₀ Control (RDF) 120:60:40: NPK. Application of NAA enhances photosynthesis, activates several enzymes, and ~~assimilate~~ assimilates transport to the stem. The ~~physiological~~ physiology and morphology of plants are greatly influenced by FYM. (Meena *et al.*, 2013) .A similar finding

was also made by (Iqbal *et al.*, 2016). With an increase in the organic manure application rate, they saw a considerable improvement in maize plant height and leaf area index. NAA provide the necessary ~~nutrients~~ nutrients for promoting healthy development and physiological processes in the plant system. Plant height, leaf area index, and dry matter output are all much greater when the rate of organic fertilizer application is increased. (Iguaet *et al.*, 2009) and both saw a similar outcome (Channal,2017). Higher plant height may be due to caused by enough room, nutrients, and sunshine being available, which drove the plants to grow vertically. The current findings closely resemble those of Kour *et al.* (2017); Husain *et al.* (2017); Almazet *et al.* (2017); Yahia and Hussein *et al.* (2017); Mahapatra *et al.* (2018); Qodliyatiet *al.* (2018); Ojha *et al.* (2018), and Ganvitet *al.* (2017).

Comment [DAS8]: Please improve the language

Comment [DAS9]: Please see whether this sentence is appropriate here.

Dry weight of plant (g)

Application of T6 FYM (10t/ha) + NAA (30ppm) significantly influenced the dry weight in Table 3 shows organic, plant growth regulator nutrient management and crop dry weight per plant at harvest. The data indicated that, baby corn at 45DAS. The maximum dry weight (95.47gm) was recorded in T6 : FYM (10t/ha) + NAA (30ppm) were statistically at par with T5,T4,T3 and lowest dry weight(67.87gm) was recorded in Farmers practice –RDF (120:60:40 kg/ha N, P and K). The therapies did not differ significantly from one another.

Table 3. Effect of organic manure and growth regulator on growth parameters of baby Corn.

Treatment details		Growth parameters	
Plant height (cm)	plant dry weight (g/plant)		
Vermicompost(4.46t/ha) +Seaweed sap (<i>Ascophyllum nodosum</i>) 5%	171.7188.00		
Vermicompost(4.46t/ha) + GA ₃ (15.47ppm)	176.2389.47		
Vermicompost(4.46t/ha) + NAA (30ppm)	179.6090.13		
FYM (10t/ha) + Seaweed sap (<i>Ascophyllum nodosum</i>) 5%	180.5394.47		
FYM (10t/ha) + GA ₃ (15.47ppm)	185.9795.87		
FYM (10t/ha) + NAA (30ppm)	189.5795.47		
Goat manure (15t/ha) + Seaweed sap (<i>Ascophyllum nodosum</i>) 5%	163.40 72.90		
Goat manure (15t/ha) + GA ₃ (15.47ppm)	168.6078.80		
Goat manure (15t/ha) + NAA (30ppm)	170.2382.77		
Control (RDF) 120:60:40: NPK	162.1767.87		
F-test		S	S
SEM(±)	4.36 3.74		
CD(P=0.05)		9.94	6.05

Comment [DAS10]: Italics the font for scientific name of organisms in whole MS

When organic manure were was applied, the physio-chemical characteristics of the soil may have improved, giving the soil a favorable ~~favorable~~ structure for root growth and soil enzymes (which continue to decompose organic matter in the soil to release nutrients and make them available near the rhizosphere for absorption by plant roots, thus improving quality (Chaouiet *al.* 2003). ~~Additional~~ Additionally, an increase in plant metabolism that appears to have encouraged

meristematic activities that led to apical development might be blamed for the effect of organic fertilization by vermicompost on LAI. This outcome is consistent with what atarzadeh and colleagues discovered (2013). The ultimate effect of photosynthesis activities is dry weight. The amount of sunlight that a plant gets determines how efficiently the photosynthesis process works and how many photosynthesis are produced. Larger plant organs will result from increased photosynthetic activity, which will also increase the dry weight of plants. According to Shah and Ahmad (2006), Meena *et al.* (2012), Ghimire *et al.* (2013), Kour *et al.* (2017), Kumar *et al.* (2014), Shahid *et al.* (2015), Wailare and Kesarwani, and others, proper nutrition and spacing promote higher vegetative development and more sunshine to plants (2017).

Yield parameters

Number of cobs per plant

The information on the number of cobs per plant impacted by treatments is reported in a ~~table for in general~~. The number of cobs per plant rose with crop stage progression regardless of treatment and peaked at harvest. At 60 DAS, the number of baby corn cobs per plant varied depending on the treatment combination. At 60 DAS, Numbers of cobs/plant was found significantly and highest Number cobs/plant (3.64) was recorded in T6:FYM (10t/ha) + NAA (30ppm) and lowest Number of cobs/plant (1.45) was recorded in T10 : Farmers practice –RDF (120:60:40 kg/ha N, P and K). By supplying the crop with the nutrients it needs from the beginning and increasing the supply of N, P, and K in a more synchronized way at the treatment receiving an integrated supply of nutrients from organic manure and growth regulator, which was expressed in terms of plant height, cobs per plant, cob girth, cob length, and cob weight with and without husk by virtue of organic manure, growth regulator ~~improve~~ improve the overall growth of the in accordance with Singh *et al.* (2015). Findings, increased availability of photosynthetic, metabolites, and nutrients to develop reproductive structures appears to have led to an increase in the number ~~of~~ of cobs/plant, length of cobs, weight of cobs, and yield of cobs with these nutrient management treatments similar result found Wailare and Kesarwani (2017) and Kour *et al.* (2017).

Comment [DAS11]: Which table?

Weight of cob (without husk) (g)

The data provided on length of cobs/plant (cm) without husk affected by treatments are presented in Table 4. In general the length of cobs/plant (cm) without husk differed with the advancement in crop stages, irrespective of the treatments and reached maximum at the time of harvest. The length of cobs/plant (cm) with and without husk of Baby corn was recorded in 60 DAS differed significantly with treatment combinations. At 60 DAS, length of cobs/plant was found significantly and the highest length of cobs/plant (cm) without husk (18.82) was recorded in T6 : FYM (10t/ha) + NAA (30ppm) and lowest length of cobs/plant without husk (11.2) was recorded in Farmers practice –RDF (120:60:40 kg/ha N, P and K).

Weight of cob (with husk) (g)

Table 4 shows organic manure and growth regulator on the weight of cob (with husk). Shows organic, growth regulator nutrient management and crop weight of cob (with husk). The data revealed that various treatments of highest of the highest, Cob weight (g) with husk were recorded at harvest stage. ~~The data~~ The data shown ~~showed~~ that there was a significant effect

among treatments. At 60 DAS, length of cobs/plant was found significantly and the highest length-length of cobs/plant (cm) with husk (25.13) was recorded in T6 : FYM (10t/ha) + NAA (30ppm) and lowest length of cobs/plant with husk (21.12) was-were recorded in Farmers practice -RDF (120:60:40 kg/ha N, P and K). By supplying the crop with the nutrients it needs from the beginning and increasing the supply of N, P, and K in a more synchronized way at the treatment receiving an integrated supply of nutrients from organic manure and growth regulator, which was expressed in terms of plant height, cobs per plant, cob girth, cob length, and cob weight with and without husk by virtue of organic manure growth regulator did significantly improve the overall growth of the in accordance with Singh *et al.* (2015), findings, increased increased availability of photo-synthesases, metabolites, and nutrients to develop reproductive structures appears to have led to an increase in the number of cobs/plant, length of cobs, the weight of cobs, anf-and yield of cobs with these nutrient management treatments, the similar result found Wailare Wail are and Kesarwani (2017) and Kour *et al.* (2017).

Table 4. Effect of organic manure, growth regulator on growth and yield of baby corn.

Treatment details (t/ha) (t/ha)	Yield parameters				
	No. of Cobs/ Plant	Cob length (cm)	Cob yield with husk	Cob yield without husk	Fodder yield (t/ha)
Vermicompost(4.46t/ha) +Seaweed sap (Ascophyllum nodosum) 5%	2.00	18.25	3.22	2.69	5.35
Vermicompost(4.46t/ha) + GA (15.47ppm)	2.43	18.69	3.33	2.86	5.42
Vermicompost(4.46t/ha) + NAA (30ppm)	2.50	18.95	3.35	2.85	5.55
FYM (10t/ha) + Seaweed sap (Ascophyllum nodosum) 5%	3.30	19.00	3.50	3.04	5.65
FYM (10t/ha) + GA ₃ (15.47ppm)	3.53	19.13	3.61	3.10	5.76
FYM (10t/ha) + NAA (30ppm)	3.64	19.26	3.68	3.15	5.88
Goat manure (15t/ha) + Seaweed sap (Ascophyllum nodosum) 5%	1.50	15.48	2.84	2.14	5.05
Goat manure (15t/ha) + GA ₃ (15.47ppm)	1.65	16.74	2.93	2.25	5.15
Goat manure (15t/ha) + NAA (30ppm)	1.80	17.32	3.14	2.32	5.22
Control (RDF) 120:60:40: NPK	1.45	14.72	2.72	2.10	5.01
F-test	S	NS	S	SS	
Sem(±)	0.313	1.20	0.11	0.07	0.10
CD(P=0.05)	0.93	-0.300	0.250	0.29	

Cob yield (t/ha)with and without husk

The data obtained on weight of cobs yield (kg/ha) with and without husk affected by treatments are presented in Table 4. In general the weight of cobs yield (kg/ha) with and without husk differed with the advancement in crop stage, irrespective of the treatment and reached maximum at the time of harvest. The weight of cobs yield (kg/ha) with and without husk of baby corn was recorded at 60 DAS ~~and~~ differed significantly with treatment combinations. At 60 DAS, weight of cobs yield was found significantly and highest weight of cobs yield was found significantly and highest weight of cobs yield (q/ha) with husk (3.68 q/ha) and without husk (3.15 kg/ha) was ~~reerded~~ recorded in T₆ : FYM (10t/ha) + NAA (30ppm) and lowest weight of cobs/plant (cm) with husk (2.72) and without husk (2.10) was recorded in Farmers practice –RDF (120:60:40 kg/ha N, P and K). Organic manure and growth regulator did bring about significant improvement in overall growth of the crop by providing needed ~~nutriebs~~ ~~nutrients~~ ~~fron~~ ~~from~~ ~~the~~ initial stage and increase in supply of N, P, and K in more synchronize way at the treatment receiving ~~an~~ integrated supply of ~~nutrient~~ ~~nutrients~~ from organic manure along with growth regulator and which expressed in terms of plant height, cobs per plant, cob girth, cob length, cob weight with and without husk by virtue of increased photosynthetic efficiency. Thus greater availability of photosynthetic, metabolites and nutrients to develop reproductive structures seems to have resulted in ~~an~~ increased number of cobs/plant, length of cobs, ~~the~~ weight of cobs and yield of cobs with these integrated nutrient management treatments which ~~coineides~~ ~~coincide~~ with results Shah and Ahmad (2006), Ghimire *et al.* (2013); Lone *et al.* (2013); Kumar *et al.* (2014); Ukonzeet *et al.* (2016) and Kour *et al.* (2017).

Conclusion

On the basis of results obtained in present investigation, it is concluded that the profitable production of baby corn can be secured by FYM (10t/ha) + NAA (30ppm) (T₆). These practices may be passed on to the farmers for obtaining higher returns in this agro-climatic zone. It ~~is~~ also recorded the maximum gross return, net return and benefits cost ratio.

References

- Abouziena, H.F. and Abd, El. Wahed M. S. 2013. Production capability of wheat cultivars under low lightintensity (date palm shade) conditions and some bioregulators. *Journal Applied Science. Research.*; **9**(8): 5176-5188.
- Arya, K. C. and Singh, S. N. 2000. Effect of different levels of phosphorus and Zinc on yield and nutrient uptake of maize with and without irrigation. *Indian Journal of Agronomy.* **45**(4): 717-721.

Comment [DAS12]: Carefully check all the citations, whether they are mentioned in references or not.

- Bakht, J., Ahmad, S., Tariq, M., Habib, A. and Shafi, M. 2006. Response of maize to planting methods and nitrogen fertilizer. *Journal. Agriculture. Biology. Science.*, 1: 8-14.
- Barbara M Humtsoe, Joy Dawson and Praveena Rajana 2018. Effect of nitrogen, boron and zinc as basal and foliar application on growth and yield of maize (*Zea mays* L.) *Journal of Pharmacognosy and Phytochemistry*; **7**(6): 01-04
- Bindhani. Anita., Barik, K .C. and Garnayak, L .M. 2007. Nitrogen management in baby corn (*Zea mays* L.). *Indian Journal of Agronomy*, **52**: 135-138.
- Bindhani. Anita., Barik, K. C. and Mahaptara, P. K. Productivity and nitrogen use efficiency of baby corn (*Zea mays* L.) at different levels and timing of nitrogen application under rain fed conditions. *Indian Journal of Agriculture Sciences*, **78**(7): 629-631.
- Bouyoucus, G.J. (1927). The hydrometer as the new method for the mechanical analysis of soil. *Soil Science* **23**: 343-353.
- Chauhan, S. K., Mohan, J., Dass, S. and Gadag, R. N. 2009. Evaluation and identification of suitable, **22**(3): 3-7.
- Choudhary, P.M., Patil, H.E. and Hanikare, R.H. 2006. Effect of INM in maize (*Zea mays* L.) on pattern of leaf area and dry matter production. *International. Journal. Plant Science.*, 1:17-21.
- Das, S., Dharam, P., Arora, P., Dhanju, K.S., and Mehla, J. C. 2004. Baby corn in crop diversification, National seminar on diversification of Agriculture through horticultural crops. *CCS, HAU, Uchani, Karnal*, 21-23.
- Das, S., Ghosh, G., Kaleem, M.D. and Bahadur, V. 2008. Effect of different levels of nitrogen and crop geometry on the growth, yield and quality of baby corn (*Zea mays* L.) CV. „GOLDEN BABY“. *ISHS Acta Horticulturae 809: International Symposium on the Socio-Economic Impact of Modern Vegetable Production Technology in Tropical Asia*.
- Duete, R. R. C., Muraoka, T., and Shiva, E. C. Economics viability and instalment doses of nitrogen fertilization in maize in EUTRUSTOX. *Acta Scientiarum Agronomy*, 31, 175-181,

2009.

Eajaz Ahmad Dar, Shah Nawaz Ahmad Rather, Amarjit Singh Harika Growth and Yield of Baby Corn (*Zea mays* L.) as affected by Different Crop Geometry and Level of Nitrogen Application IJSR - *International Journal Of Scientific Research* 3(8): 2277 – 8179

Eteng Ernest U. (2017). Response of Zn Uptake, Grain and Other Yield Components of Five Maize Hybrids as Influenced by Zinc Fertilization Methods in A Marginal Coastal Plain Sand Soil. *International Journal of Research Studies in Science, Engineering and Technology*, 4(12): 37-46.

Fakir, O.A., Rahman, M.A. and Jahiruddin, M. 2016. Effects of foliar application of Boron on the grain set and yield of wheat. *American Journal of Experimental Agriculture*; 12(2): 1-8.

Forlani G., Pastorelli, R., Branzoni, M. and Favilli, F., 1998. Root colonization efficiency, plant growth promoting activity and potentially related properties in plant associated bacteria. *Journal of Genetics and Breeding* 49: 343-351.

Ganesaraja, V., Rani, S., and Kavitha, M. P. 2009. Effect of drip irrigation regimes and fertilizer application methods on growth, yield and nutrient uptake of baby corn. *Journal of Maharashtra Agriculture University*, 34(1): 92-93.

George, R., and M. Schmitt. (2002). Zinc for crop production. University of Minnesota. Humtsoe B, Barbara M, Dawson J and Rajana P. (2018). Effect of nitrogen, boron and zinc as basal and foliar application on growth and yield of maize (*Zea mays* L.) *Journal of Plant Pathology*; 7(6): 01-04.

Gul, S., Khan, M.H., Khanday, B.A. and Nabi, S. 2015. Effect of sowing methods and NPK levels on growth and yield of rainfed maize (*Zea mays* L.). *Scientifica*: 1-6.

Gurminder, S., Kumar, R. Kumar, S., (2006). Effect of tillage and nitrogen levels on growth and yield of maize (*Zea mays* L.) *Annals. Agriculture. Research. New Series* 27(2): 198-199.

Ibeawuchi, I. I, Matthews –Njoku, Edna; Of or, Miriam O; Anyanwu, Chinyere P and Onyia, V. N. 2008. Plant spacing, dry matter accumulation and yield of local and improved maize

cultivar. *Journal. American. Science.* **4**(1): 11-19.

Iqbal, A., Ayoub. M., Zaman, H. And Ahmed, R., 2006. Impact of nutrient management and legumes association on agro qualitative traits of maize forage *Pakistan. Journal.Botany.* **38**, 1079-1084.

Jaliya, M. M., Ibrahim, A, Babaji, B. A. Sani, B. M. and Aminu D. (2013). Effect of nitrogen and sulphur fertilizers on maize grain protein content of QPM (Quality Protein Maize) maize varieties at Samaru Zaria. *Global Journal of Bio-Science and Biotechnology.* **2**(1): 132-134.

Jat, N. K., Kumar, A and Dhar S. 2010. Influence of sesbania green with or without wheat residues and Nitrogen fertilization on maize – wheat cropping system. *Indian Journal of Agronomy*, **55**(4): 253-258.

Jnana Bharati Palai, NC Sarkar and Jagadish Jena 2018. Effect of zinc on growth, yields, zinc use efficiency and economics in baby corn *Journal of Pharmacognosy and Phytochemistry*; **7**(2): 1641-1645

Jackson, M.L. 1973. Soil chemical analysis. Prentice Hall of India Pvt. Ltd., New Delhi. PP. 56.

Kalpana, R. and Krishnarajan, J. 2002. Effect of dose and time of potassium application on yield and quality of bay corn .*Agriculture Science Digest*, **22**(1): 59-60

Karki, T.B., Kumar, A. and Gautam, R .C. 2005. Influence of integrated nutrient management on growth, yield, content and uptake of nutrient and soil fertility status in maize. *Indian journal of Agriculture Science*, **75**(10): 682-685 -.

Kumar, A. K Sagar, G. K. Chandrika, V .and Reddy, p. M. 2009. Influence of integrated nitrogen management on yield nitrogen uptake, soil fertility status and economics of baby corn. *Indian journal of Agricultural Research*, **43**(3): 227 -229.

Kumar, A.S., Sakthivel, N., Subramanian. E., Kalpana, R., Janaki, P. and Rajesh, P. (2013). Foliar application of nutrients and plant growth regulators on growth and yield of finger millet. *Journal of Pharmacognosy and Phytochemistry*, **7**(3): 3032-3035.

- Kumar, A.S., Sakthivel, N., Subramanian. E., Kalpana, R., Janaki, P. and Rajesh, P. (2018). Foliar application of nutrients and plant growth regulators on growth and yield of finger millet. *Journal of Pharmacognosy and Phytochemistry*7(3): 3032-3035.
- Kumar, P., Desai, B. K. and Pujari, B. T. 2007. Effect of Integrated Nutrient Management on Economics of Maize Cultivation. *Karnataka Journal. Agriculture. Science.*,20(4): 831-832.
- Kunjir, S.S., Pinjari, S.S., Suryavanshi, J.S. and Bhonde, T.S. 2009. Effect of planting geometry, nitrogen levels and micronutrients on growth and yield of Sweet corn. *Bioinfolet*, 6: 22-24.
- Lamana, M. C. L. 2003. Effect of spacing between plants on growth and forage yield of two maize (*Zea mays* L.) cultivar .M. Sc. (Ag.) thesis, sudan University of science and Technology, sudan, 55.
- Liu H., Gan W, Rengel, Zhao, P. (2016). Effects of zinc fertilizer rate and application method on photosynthetic characteristics and grain yield of summer maize. *Journal of Soil Science and Plant Nutrition*, 16(2), 550-562
- Managaser,V.T. 2003 intercropped with pole sito .Population density and nitrogen fertilization of young baby cob corn in Research Journal Agriculture Department ,DMMMUS,pp.1-25.
- Margar, E., and Dawson. J., (2017) Effect of Biofertilizers level of Nitrogen and Zinc on growth and yield of hybrid maize (*Zea mays* L.) *International. current. Microbial. Applied.Science*6(9): 3614-3622.
- Mayank P., Saini, K.P., Yadav, V.and Yadav, R.K. (2018). Effect of foliar application of PGRson growth, yield and yield attributes of rice under salt stress condition. *Journal of Pharmacognosyand Phytochemistry* 2: 268-272.
- Md. Asaduzzaman ,Mrityunjoy Biswas, Md. Nazrul Islam, Mohammad Mokhlesur Rahman, Rafeza Begum, Md. Abdur Rahman Sarkar, Md. Asaduzzaman. 2014. Variety and N-Fertilizer Rate Influence the Growth, Yield and Yield Parameters of Baby Corn (*Zea mays* L.) *Journal of Agricultural Science*; 6(3): 1916-9760.
- Mona E. El-Azab. (2015). Increasing Zn ratio in a compound foliar NPK fertilizer in relation to growth, yield and quality of corn plant, *Journal. Innovations. Pharmaceutical. Biological*

science, **2**(4): 451-468.

Moreira, J. N., Silva, P .S., Silva, L. Dombroski, J. L. D. and Castro, R. S. 2019. Effect of detasseling on baby corn, green ear and grain yield of two maize hybrids. *Horticulture Brasileira*, **28**(4): 406 -411.

Motto, M and Mall, R H. 1983.Prolificacy in maize .A review .*Maydica* 23: 53-56.

Nagasubramaniam, A., Pathmanabhan, G. and Mallika, V. Studies on improving production potential of baby corn with foliar spray of plant growth regulators. *Annual. Plant Physiol.*2007; **21**(2):154-157.

Nahar. K. Ahmed, S.,Akanada, M. A. L. OndalM .A .L. Mondal, M. A. A and Islam, M .A .Genotype environment interaction for baby cob yield and maturity in baby corn .*Bangladesh journal of Agriculture Research*, **35**(3), 489 -496, 2010.

Naik, R.D., and More, S.M., (2015). Integrated nutrient management studies in finger millet. *Crop Res.*, **48**(2): 27-31.

Ochaong, P. 2005. Appropriate variety, plant density and rate and time of nitrogen fertilization for baby corn production in Amphoe Kamphaeng saen. *Thailand Research Journal of Agriculture and Biological Sciences*,**1**(4): 303-307.

Omar, H., Al-Rawi., Adil H. Abdulkafoor, Saad I. yousif. and Mustafa R. Al-Shaheen. (2018). Effect of spraying with different levels of salicylic and humic acid in some growth characteristics and yield of wheat. *ISSN: 3*:133-138.

Olsen, S.R., Cole, V.V., Watanable, F.S. and Dean L.A. 1954. Estimation of available phosphorous in soil by extraction with sodium bicarbonate. United States Department of Agriculture, Circular **939**: 1-9.

Usadadia, P. S. Mistry, N. G. Savani and K. K. Patel. 2019 Effect of Different Levels of Irrigation,Nitrogen and Foliar Application of Banana Pseudostem Sap on Drip Irrigated Sweet Corn - Green Gram Cropping Sequence, *Indian. Journal. Pure Applied. Bioscience*.**7**(5): 254-258.

- Pandey, A.K., Prakash, V., Mani, V.P. and Singh, R. D. 2000. Effect of rate of nitrogen and of application on yield and economics of baby corn (*Zea mays* L.). *Indian Journal of Agronomy* **45**(2): 338-343.
- Pandey, A. K., Mani, V. P., Prakash, V., Singh, R. D. and Gupta, H. S. 2002. Effect of varieties and plant densities on yield, yield attributes and economics of baby corn (*Zea mays* L.). *Indian Journal of Agronomy*. 47: 221-226.
- Panwar, A. K. and Munda, G. C. Response of baby corn (*Zea mays* L.) to nitrogen and land configuration in mid hills of Meghalaya .*Indian journal of Agricultural Science*, **76**(5), 293-296.
- Paramasivan, M., Kumaresan, K, R., Malarvizhi, P.,Mahimairaja, S. and Velayudham, K. 2010. Effect of different levels of NPK and zinc on yield and nutrient uptake of hybrid maize in Mayamankuruchi series of soil of Tamil nadu. *Asian journal of soil Science*, **5**(1): 157 - 161.
- Prabha, A., Parasuraman, P., Sivagamy K. and Sivakumar, B. (2016). Growth, yield and economics of irrigated finger millet as influenced by system of finger millet intensification (SFI) practices in north eastern zone of Tamil Nadu. *Journal of Pharmacognosy and Phytochemistry*, **8**(3): 600-663.,
- Bala, S. and Khoyumthem, P. 2007. Estimation of yield and economic return in baby corn. *Envirnment and Ecology*,**25**(4): 945-047,
- Rakesh Kumar, Narendra Kumawat, Sudhir Kumar, Amitesh Kumar Singh and J.S. Bohra. 2017. Effect of NPKS and Zn Fertilization on, Growth, Yield and Quality of Baby Corn-A Review. *International Journal of Current Microbiology and Applied Sciences*, ISSN: 2319-7706. **6**(3): 1392-1428.
- Ram, V., Singh, R.N. and Singh, K. 2006. Studies on integrated use of FYM, nitrogen and sulphur on growth, yield attributes and yield on winter maize (*Zea mays* L.). *Plant Archi*, **6**:749-752.

- Ramchandruppa, B. K., Nanjappa, H .V. and Soumya, T. M. 2010. Effect of stages of harvest on yield and quality of baby corn varieties. *Mysore journal of Agriculture Science*: **40**(4): 453-457.
- Ramesh S, P. Sudhakar, S. Elankavi, K. Suseendran. and Jawahar, S. (2019). Effect of Gibberellic acid (GA3) on growth and yield of rice (*Oryza sativa* L.). *Plant Archives*; **19**: 2581-6063.
- Rao, L. K., kumar, R. A. and Lal, G. M. 2009. Effect of Integrated nitrogen management on growth and yield of baby corn (*Zea mays* L.) cv. Mridula. *journal of Maharashtra Agriculture University*, **34**(3): 249-251.
- Richards, L.A. 1954. *Diagnosis and improvements of saline and alkali soils. Agriculture Handbook No. 60. USDA, Washington*
- Sahoo, S. C. and Panda, M. M. 1999. Effect of level of nitrogen and plant population on yield of baby corn (*Zea mats* L.). *Indian Journal of Agriculture Sciences***69**(2):157-158.
- Sahoo,S.C.Yield and economics of baby corn (*Zea mays* L.) as affected by varieties and levels of nitrogen .*Range Management and Agroforestry*, **32**(2): 135-137,2011.
- Sarakhsi ,H S ,Yarnia ,M and Amirniya ,R .Effect of nitrogen foliar application in different concentration and growth stage of baby corn (hybrid -704) *Advance in Environmental Biology* ,**4**(2) ,291-298 2010.
- Sathishkumar, A. N., Sakthivel, E. Subramanian, R. Kalpana, P, Janaki. and P, Rajesh. (2018). Foliar application of nutrients and plant growth regulators on growth and yield of finger millet. *Journal of pharmacognosy and phytochemistry*; **7**(3): 3032-3035.
- Scaria Dona, Rajasree G and Sudha B of baby corn (*Zea mays*). (2016). Effect of varieties and spacing on growth, yield and economics of cultivation of baby corn (*Zea mays* L.) as inter crop in coconut garden. *Research on crops*.**17**(4): 673-678.
- Shafea,L and Saffari ,M (2007). Effect of zinc (ZnSO₄) and nitrogen on chemical composition of maize grain. *International Journal Agriculture. Science*.**1**(6): 323-328.

- Shah S, Ghani G, Khan H, Arif M., Qahar, Inamullah, Ali A. and Ahmad M., (2015) response of maize cultivars to phosphorus and zinc nutrition. *journal. botany.*, 47(si): 289-292.
- Shaikh Wasim Chand, R Susheela, D Sreelatha, M Shanti and SA Hussain. 2017. Effect of zinc fertilization on yield and economics of baby corn (*ZeamaysL.*) *Journal of Pharmacognosy and Phytochemistry*; **6(5)**: 989-992.
- Shaikh Wasim Chand, R Susheela, D Sreelatha, M Shanti and SA Hussain Effect of zinc fertilization on yield and economics of baby corn (*Zea mays L.*) *Journal of Pharmacognosy and Phytochemistry* 2017; **6(5)**: 989-992
- Sharma, S. K., Swami, A A., and Singh R.K., (1992). Relative response of maize (*Zea mays L.*) varieties to zinc. *Indian Journal . Agronomy.*, **37(2)**: 361-361.
- Siam, H.S., Kader, E.M.G.A. and Alia, E.H.I. 2008. Yield and yield component of maize as affected by different sources and application rates of N fertilizers. *Research. Journal. Agriculture. Biological. Science.*, 6: 399-412.
- Singh, M. K. Singh. R. N., Singh S, P. S. P. Yadav M. K. and Singh, V. K. 2010. Integrated nutrient management and for higher yield quality and profitability of baby corn (*Zea mays L.*) *Indian journal of Agronomy*, **55(2)**, 100-104,
- Singh, D. P., Rana, N.S. and Singh R. P., (2000). Growth and yield of winter maize as influence as intercrops and nitrogen application. *Indian J. Agronomy.*, **45(3)**: 515-519.
- Singh, N.T., Viv, A. C. and Singh, R. 1985. Nitrogen response of maize under temporary flooding. *Nutrient Cycling in Agrosystem*, **6(2)**; 11-12.
- Siva, K., Shinggu C.P., Dadari, S.A., Shebayan, J.A.Y., Adekpe, D.I., Mahadi, M.A., Mukhtar, A. and SivaKumar R.G. PaThmanaban, M. K. Kalarani, Mallika Vanangamudi and P.S. Srinivasan. 2002. Effect of Foliar Application of Growth Regulators on Biochemical Attributes and Grain Yield in Pearl Millet. *Indian Journal Plant Physiol.*, **7(1)**: 79-82.
- Sobhana, V., Kumar, A., Idnani, L. K., Singh, I. and Shivadhar. 2012. Plant population and nutrient requirement for baby corn hybrids (*Zea mats L.*). *Indian journal of*

Agronomy,**57**(3): 294-296.

Srinivasan K. (1992). Effect of amendment and zinc levels on growth and yield of maize (*Zeamays L.*) *Indian Journal Agriculture*.**37** (2): 246-249.

Suresh, G., Guru, G. and Lokanadan, S. (2018). Effect of Nutrient Levels and Plant Growth Regulators on Growth Parameters of Pearl Millet. *Int. J. Pure App.Biosci.***6** (3): 271-277.

Tamrakar, S.K., Singh, P., Kumar, V. and Tirkey, T. (2018). Effect of gibberellic acid, salicylic acid, cow urine and vermiwash on corn production of *Gladiolus cv. Candyman*. *International Journal Current Microbiol Applied Science*. **6**: 677-686.

Thakur, D R., O M., and Kharawara, P.C. 1998. Effect of nitrogen and plant spacing on yield, Nitrogen uptake and economics in baby corn (*Zea mays L.*).*Indian Journal of Agronomy* **43**(4):668-71.

Thakur, D. R. and Sharma, V. 1999. Effect of varying rates of nitrogen and its schedule of split application in baby corn (*Zea mays L.*). *Indian journal of Agricultural Science*,**69**(2): 93-95.

Thakur, D. R.Prakash, O M., Kharwara P. C. and Bhalla, S. K. 1997. Effect of nitrogen and plant spacing on growth, development and yield of baby corn. *Indian journal of Agronomy*, **42**(3): 479 -483.