

Influence of growth regulator and organic fertilizers in the cultivation from Baby corn (*Zea mays* L.)

ABSTRACT

The experiment consisted of ten organic and diverse combination treatments plant growth regulator nutrient management replicated in triplicate in a randomized block design. The primary purpose of the trial was to evaluate the effect organic manure, plant growth regulator, growth and yield of baby corn (*Zea mays* L.) state of Prayagraj. Three organic levels manure and plant growth regulator are FYM (10t/ha) and NAA (30 ppm), FYM(10t/ha) and GA₃ (15.47 ppm), FYM (10t/ha) and Seaweed sap (*Ascophyllum nodosum*) 5% can be concluded from the current study that profitable production of baby corn can be ensured FYM (10t/ha) and NAA (30 ppm) + FYM(10t/ha) and GA₃ (15.47 ppm) (T₆). Baby corn is free from pesticides and its nutritional value is comparable to popular vegetables like cabbage and cucumber. Its by-products such as tassel, young husk, silk and green stalks provide good cattle food. It is high in potassium, folic acid, and is a rich source of A, B, E and many other minerals. Corn will remain one of the important, field crops in the developing countries. Considerable scopes exists from promoting baby corn technology in Asia-pacific region. The baby corn industry provides opportunities for higher income, generates employment for the rural poor potential for export. Baby corn is expected to catch the attention of more and more consumers and farmers because of its superior taste and texture.

Keywords: Baby corn, organic manure, growth regulator, yield and quality

Introduction

Maize (*Zea mays* L) is the most versatile and emergent crop with many adaptations to different agroclimatic conditions. It is famous as the queen of cereals because of its genetic yield potential among other cereal crops compared to rice, wheat, oats, sorghum and others in the Ober region. In most developing countries, maize contributes a major part of food security. Maize is the third most important crop in India after rice and wheat. Not only for human consumption and animal feed, but also for maize, corn, etc. That is why it is used in industry for production. Then more attention was paid to the cultivation of corn by researchers and agronomists. in addition to maximizing profits for producers, take advantage of the opportunity to generate more foreign exchange earnings. Baby corn, as the name suggests, is not genetic corn, but an immature ear of common corn. The ear of corn is hard and cannot be used as a vegetable. Baby corn ears are tender and eaten by humans as a vegetable (Jinjala et al., 2016). Baby corn was harvested during silking stage. After harvesting, the external sheath was removed and the ear was used for vegetable purpose viz., salad,soup, pickles etc(Muthukumar *et al.*, 2005). Baby corn is a delicious, decorative, low caloric nutritious vegetable without cholesterol and is rich in fibre content. It is free from pests and diseases and it contains protein upto 15 to 18 percent,sugar0.016to0.020 percent phosphorus 0.6to0.9 percent,potassium 2to 3percent,fibre 3to 5percent,calcium 0.3to0.5per cent and ascorbic acid 75to 80mg100/g. As green fodder, It is the

best suited for mil chainmails since it has lactogenic properties(Reenarani *et al.*, 2017).For the past few decades in creaseduse of Synthetic fertilizers have reduced the use of organic fertilizers affecting soil fertility and productivity. Organic farming methods have improved the sustainability and health of the soil without affecting the ecosystem.

Materials and methods

The current study was conducted at the crop farm, Department of Agronomy, Naini Institute of Agriculture, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj, during Kharif season 2022, (U.P.). The test site is located on the left side of the Prayagraj-Rewa road, about four kilometers from Prayagraj and near the unaamuna river, at 25.57° N latitude, 87.19° E longitude and at an altitude of 98 m above sea level. Uttar Pradesh, where Prayagraj is located, is a subtropical region with hot summers and cool winters. The average temperature in that area ranges from 23°C to 38°C, rarely dropping below 3°C or 4°C. The relative humidity ranges from 28.57% to 95%. The average annual rainfall here is 1050mm. Chemical analysis of the soil revealed a sandy loam texture with a pH of 7.20, low organic carbon (0.83 percent) and low potassium (208.8 kg/ha) and phosphorus (17.2 kg/ha). Electrically conductive soil, with a conductivity of 0.34 ds/m. Three replicates were used for each of the nine treatment combinations. Details of treatments and treatment combinations are shown in Tables 1 and 2 respectively. Organic fertilizers, plant growth regulators and plants are applied according to the combination of cultivation. Plant height at harvest (cm), dry weight at harvest, number of weeds/plants, seedling length, tillering, yield of baby corn (t/ha) were measured and economic analysis of each treatment was completed to determine the best treatment combination. . to grow baby corn.

Table 1. Treatments details

Organic manure	
Vermicompost	4.46t/ha
FYM	10t/ha
Goat manure	15t/ha
Plant growth regulator	
Seaweed-sap (<i>Ascophyllum nodosum</i>)	5%
GA3	15.46 ppm
NAA	30 ppm

Table 2. Treatment combination

Icon Treatment	Combinations Treatment
T ₁	Vermicompost (4.46t/ha) + Seaweed sap (<i>Ascophyllum nodosum</i>) 5%
T ₂	Vermicompost(4.46t/ha) + GA ₃ (15.47 ppm)
T ₃	Vermicompost(4.46t/ha) + NAA (30 ppm)
T ₄	FYM (10t/ha) + Seaweed sap (<i>Ascophyllum nodosum</i>) 5%
T ₅	FYM (10t/ha) + GA ₃ (15.47 ppm)
T ₆	FYM (10t/ha) + NAA (30 ppm)
T ₇	Goat manure (15t/ha) + Seaweed sap (<i>Ascophyllum nodosum</i>) 5%
T ₈	Goat manure (15t/ha) + GA ₃ (15.47 ppm)
T ₉	Goat manure (15t/ha) + NAA (30 ppm)
T ₁₀	Control (RDF) 120:60:40: NPK

Results and discussion

Development parameters

Plant height (cm)

Table 3 shows organic fertilizers, plant growth regulator nutrients and crop spacing in plant height at harvest. The data show a significant effect on plant height during plant growth. Application of T₆- FYM (10t/ha) + NAA (30 ppm) significantly influenced the plant height of baby corn at 45DAS. The maximum plant height (189.57cm) was recorded in T₆ FYM (10t/ha) + NAA (30 ppm) which was statistically at par with T₅, T₄, T₃ and minimum plant height (162.17) was recorded in the application of T₁₀ Control (RDF) 120:60:40: NPK. Application of NAA enhances photosynthesis, activates several enzymes, and assimilates transport to the stem. The physiology and morphology of plants are greatly influenced by FYM. (Meena *et al.*, 2013). A similar finding was also made by (Iqbal *et al.*, 2016). With an increase in the organic manure application rate, they saw a considerable improvement in maize plant height and leaf area index. NAA provide the necessary nutrients for promoting healthy development and physiological processes in the plant system. Plant height, leaf area index and dry matter output are all much greater when the rate of organic fertilizer application is increased. (Igua *et al.*, 2009) and both saw a similar outcome (Channal,2017). Higher plant height may be due to enough room, nutrients, and sunshine being available, which drove the plants to grow vertically. The current findings closely resemble those of Kour *et al.* (2017); Husain *et al.* (2017); Yahya and Husayn and others. (2017); Mahapatra *et al.* (2018); Law and others. (2018); Ojha *et al.* (2018) and Ganvit *et al.* (2017).

Dry weight of plant (g)

Application of T₆ FYM (10t/ha) + NAA (30 ppm) significantly influenced the dry weight in Table 3 shows organic, plant growth regulator nutrient management and crop dry weight per plant at harvest. Data shows that there is baby corn in 45DAS. The highest dry weight (95.47gm) was recorded in T₆: FYM (10t/ha) + NAA (30 ppm) statistically compared to T₅, T₄, T₃ and the lowest dry weight (67.87gm) in the experiment Farmer –RDF (120: 60: 40 kg/ha N, P and K) was recorded. Treatment methods were not significantly different from each other.

Table 3. Effect of organic manure and growth regulator on maize growth parameters

Details of treatment	Development parameters	
	Plant height (cm)	plant dry weight (g/plant)
Vermicompost (4.46t/ha) + Seaweed sap (<i>Ascophyllum nodosum</i>) 5%	171.71	88.00
Vermicompost(4.46t/ha) + GA ₃ (15.47 ppm)	176.23	89.47
Vermicompost(4.46t/ha) + NAA (30 ppm)	179.60	90.13
FYM (10t/ha) + Seaweed sap (<i>Ascophyllum nodosum</i>) 5%	180.53	94.47
FYM (10t/ha) + GA ₃ (15.47 ppm)	185.97	95.87
FYM (10t/ha) + NAA (30 ppm)	189.57	95.47
Goat manure (15t/ha) + Seaweed sap (<i>Ascophyllum nodosum</i>) 5%	163.40	72.90
Goat manure (15t/ha) + GA ₃ (15.47 ppm)	168.60	78.80
Goat manure (15t/ha) + NAA (30 ppm)	170.23	82.77
Control (RDF) 120:60:40: NPK	162.17	67.87
F-test	S	S
SEM(±)	4.36	3.74
CD(P=0.05)	9.94	6.05

When organic manure was applied, the physio-chemical characteristics of the soil may have improved, giving the soil a favourable root growth and soil enzyme structure (which continues to break down organic matter in the soil near the rhizosphere to remove harmful substances and be absorbed by plant roots, thereby improving quality (Chaoui et al. 2003). Additionally, an increase in plant metabolism that appears to have encouraged meristematic activities that led to apical development might be blamed for the effect of organic fertilization by vermicompost on LAI. This outcome is consistent with what Atarzadeh and colleagues discovered (2013). The ultimate effect of photosynthesis activities is dry weight. The amount of sunlight that a plant gets determines how efficiently the photosynthesis process works and how many photosynthesis are produced. Larger plant organs will result from increased photosynthetic activity, which will also increase the dry weight of plants. According to Shah and Ahmad (2006), Meena *et al.* (2012), Ghimire *et al.* (2013), Kour *et al.* (2017), Kumar *et al.* (2014), Shahid *et al.* (2015), Wailare and Kesarwani, and others, proper nutrition and spacing promote higher vegetative development and more sunshine to plants (2017).

Product parameters

Number of bulbs per plant

Data on the number of plants that grew under the influence of treatment are reported in the table. Although processed, the number of plants growing in the harvest increased and reached a maximum in the harvest. At 60 DAS, the number of baby corn bulbs per plant varied treatment combination. At 60 DAS, Numbers of cobs/plant was found significantly and highest Number cobs/plant(3.64) was recorded in T6: FYM (10t/ha) + NAA (30 ppm) and lowest Number of cobs/plant(1.45) was recorded in T10 : Farmers practice –RDF (120:60:40 kg/ha N, P and K). By

supplying the crop with the nutrients it needs from the beginning Plant height, plant head, density, length, weight with and without organic fertilizer, growth regulator Singh et al. improve overall development accordingly. (2015). The findings were similar to the increase in photosynthesis, metabolites and nutrients for the development of reproductive structures, an increase in the number of plants / plants, shoot length, shoot weight and shoot yield with this nutrient management treatment. Wailare and Kesarwani (2017) and Kour et al. (2017).

Weight of cob(without husk) (g)

The data provided on length of cobs/plant (cm) the shells affected by the treatment are shown in Table 4. In general, the plant length (cm) varied with the growth stage of the crop regardless of the treatment and reached the maximum at harvest. 60 Maize plants with and without maize (plant) length recorded at 60 DAS differed significantly with treatment combinations. At 60 DAS, bulb/plant length was found to be significant and maximum length (18.82) was recorded in T6: FYM (10t/ha) + NAA (30 ppm) and minimum length. /plants without pods (11.2) –RDF (120:60:40 kg/ha N, P and K) was recorded in farmers' experiments.

Bottle weight (with shell) (g)

Table 4 shows organic fertilizers and growth regulators by container weight. It shows organic, nutrient management that stimulates the growth and weight of the onion crop. The data revealed that different treatments were recorded with maximum cob weight (g) at harvest time. The data showed a significant interaction between treatments. At 60 DAS, plant/plant length was found to be significant and the maximum length (25.13) was recorded in T6: FYM (10t/ha.) + NAA (30 ppm) and the lowest length (21.12) was recorded in farmer practices –RDF (120: 60: 40 kg/ha N, P and K). By providing plants with the nutrients they need from the beginning and multiplying the supply of N, P and K more synchronously in the integrated nutrient treatment of organic fertilizers and growth regulators Plant height, bulb growth, density, length and density weight with. the help of organic fertilizer growth regulator, Singh et al. significantly improved overall development, according to (2015), findings show that the increase in photo-synthesis, metabolites and nutrients for the development of reproductive structures lead to an increase in the number of bulbs / plants, length of bulbs, weight of bulbs and yield. this nutritional management treatment, Vail and Kesarwani (2017) and Kour et al. (2017).

Table 4. Effect of organic manure, regulator of growth and production of baby corn.

Treatment details	Yield parameters				
	No. of cobs/plant	Cob Length (cm)	Cob Yield with husk (t/ha)	Cob yield without husk (t/ha)	Fodder Yield (t/ha)
Vermicompost(4.46t/ha) + Seaweed sap (Ascophyllum nodosum) 5%	2.00	18.25	3.22	2.69	5.35
Vermicompost(4.46t/ha) + GA (15.47ppm)	2.43	18.69	3.33	2.86	5.42
Vermicompost(4.46t/ha) + NAA (30ppm)	2.50	18.95	3.35	2.85	5.55
FYM (10t/ha) + Seaweed sap (Ascophyllum nodosum) 5%	3.30	19.00	3.50	3.04	5.65
FYM (10t/ha) + GA ₃ (15.47ppm)	3.53	19.13	3.61	3.10	5.76
FYM (10t/ha) + NAA (30ppm)	3.64	19.26	3.68	3.15	5.88
Goat manure (15t/ha) + Seaweed sap (Ascophyllum nodosum) 5%	1.50	15.48	2.84	2.14	5.05
Goat manure (15t/ha) + GA ₃ (15.47ppm)	1.65	16.74	2.93	2.25	5.15
Goat manure (15t/ha) + NAA (30ppm)	1.80	17.32	3.14	2.32	5.22
Control (RDF) 120:60:40: NPK	1.45	14.72	2.72	2.10	5.01
F-test	S	NS	S	S	S
Sem(±)	0.313	1.20	0.11	0.07	0.10
CD(P=0.05)	0.93	-	0.30	0.25	0.29

Cob yield (t/ha) and involuntarily

Bark yield weight data (kg/ha) affected by the treatment is shown in Table 4. In general, the growth of plant meat (kg/ha) is different from the growth of plant meat, which reaches the highest level in the harvest regardless of the treatment. Seed yield weight (kg/ha) was recorded at 60 DAS and differed significantly with treatment combinations. The yield weight of onion in 60 DAS was found to be significant and the maximum weight (q/ha) was recorded for both shell (3.68 q/ha) and shell (3.15 kg/ha). In T6: Farmers –RDF (120:60:40 kg/ha N, P and K). Organic fertilizers and growth regulation, early stage nutrient supply and more synchronous increased supply of N, P, and K in the treatment receiving integrated organic nutrients resulted in significant improvement in overall crop growth. Along with plant regulator and plant height, plant head, density, length, weight and container density due to increased photosynthetic efficiency. Therefore, the high availability of photosynthetic, metabolites, and nutrients for the development of reproductive structures seems to have led to increased bulb/plant, length, weight, and yield with integrated nutrient management treatment consistent with branching and yield. Ahmad (2006), Ghimire et al. (2013); Dreaming etc. (2013); Kumar et al. (2014); Ukonze et al. (2016) and Kour et al. (2017).

Conclusion

Based on the results obtained in this study, it is concluded that profitable production of baby corn can be ensured by FYM (10t/ha) + NAA (30 ppm) (T6). This practice can be transferred to farmers for higher income in these agro-climatic zones. It also recorded the highest gross profit, net profit and profit margin.

References

- Abouzienna, H.F. and Abd, El. Wahed M. S. 2013. Production capability of wheat cultivars under low light intensity (date palm shade) conditions and some bioregulators. *Journal Applied Science. Research.*; **9**(8): 5176-5188.
- Arya, K. C. and Singh, S. N. 2000. Effect of different levels of phosphorus and Zinc on yield and nutrient uptake of maize with and without irrigation. *Indian Journal of Agronomy.* **45**(4): 717-721.
- Bakht, J., Ahmad, S., Tariq, M., Habib, A. and Shafi, M. 2006. Response of maize to planting methods and nitrogen fertilizer. *Journal. Agriculture. Biology. Science.*, 1: 8-14.
- Barbara M Humtsoe, Joy Dawson and Praveena Rajana 2018. Effect of nitrogen, boron and zinc as basal and foliar application on growth and yield of maize (*Zea mays* L.) *Journal of Pharmacognosy and Phytochemistry*; **7**(6): 01-04
- Bindhani. Anita., Barik, K .C. and Garnayak, L .M. 2007. Nitrogen management in baby corn (*Zea mays* L.). *Indian Journal of Agronomy*, **52**: 135-138.
- Bindhani. Anita., Barik, K. C. and Mahaptara, P. K. Productivity and nitrogen use efficiency of baby corn (*Zea mays* L.) at different levels and timing of nitrogen application under rain fed conditions. *Indian Journal of Agriculture Sciences*, **78**(7): 629-631.
- Bouyoucus, G.J. (1927). The hydrometer as the new method for the mechanical analysis of soil. *Soil Science* **23**: 343-353.
- Chauhan, S. K., Mohan, J., Dass, S. and Gadag, R. N. 2009. Evaluation and identification of suitable, **22**(3): 3-7.

- Choudhary, P.M., Patil, H.E. and Hanikare, R.H. 2006. Effect of INM in maize (*Zea mays* L.) on pattern of leaf area and dry matter production. *International. Journal. Plant Science.*, 1:17-21.
- Das, S., Dharam, P., Arora, P., Dhanju, K.S., and Mehla, J. C. 2004. Baby corn in crop diversification, National seminar on diversification of Agriculture through horticultural crops. *CCS, HAU, Uchani, Karnal*, 21-23.
- Das, S., Ghosh, G., Kaleem, M.D. and Bahadur, V. 2008. Effect of different levels of nitrogen and crop geometry on the growth, yield and quality of baby corn (*Zea mays* L.) CV. „GOLDEN BABY“. *ISHS Acta Horticulturae 809: International Symposium on the Socio-Economic Impact of Modern Vegetable Production Technology in Tropical Asia*.
- Duete, R. R. C., Muraoka, T., and Shiva, E. C. Economics viability and instalment doses of nitrogen fertilization in maize in EUTRUSTOX. *Acta Scientiarum Agronomy*, 31, 175-181, 2009.
- Eajaz Ahmad Dar, Shahnawaz Ahmad Rather, Amarjit Singh Harika Growth and Yield of Baby Corn (*Zea mays* L.) as affected by Different Crop Geometry and Level of Nitrogen Application *IJSR - International Journal Of Scientific Research* 3(8): 2277 – 8179
- Eteng Ernest U. (2017). Response of Zn Uptake, Grain and Other Yield Components of Five Maize Hybrids as Influenced by Zinc Fertilization Methods in A Marginal Coastal Plain Sand Soil. *International Journal of Research Studies in Science, Engineering and Technology*,4(12): 37-46.
- Fakir, O.A., Rahman, M.A. and Jahiruddin, M. 2016. Effects of foliar application of Boron on the grain set and yield of wheat. *American Journal of Experimental Agriculture*; **12**(2): 1-8.
- Forlain G., Pastorelli, R., Branzoni, M. and Favilli, F., 1998. Root colonization efficiency, plant growth promoting activity and potentially related properties in plant associated bacteria. *Journal of Genetics and Breeding* 49: 343-351.
- Ganesaraja, V., Rani, S., and Kavitha, M. P. 2009. Effect of drip irrigation regimes and fertilizer application methods on growth, yield and nutrient uptake of baby corn. *Journal of*

Maharashtra Agriculture University, **34**(1): 92-93.

- George, R., and M. Schmitt. (2002). Zinc for crop production. University of Minnesota. Humtsoe B, Barbara M, Dawson J and Rajana P. (2018). Effect of nitrogen, boron and zinc as basal and foliar application on growth and yield of maize (*Zea mays* L.) *Journal of Plant Pathology*; **7**(6): 01-04.
- Gul, S., Khan, M.H., Khanday, B.A. and Nabi, S. 2015. Effect of sowing methods and NPK levels on growth and yield of rainfed maize (*Zea mays* L.). *Scientifica*: 1-6.
- Gurminder, S., Kumar, R. Kumar,S., (2006). Effect of tillage and nitrogen levels on growth and yield of maize (*Zea mays* L.) *Annals. Agriculture. Research. New Series***27**(2): 198-199.
- Ibeawuchi, I. I, Matthews –Njoku, Edna; Of or, Miriam O; Anyanwu, Chinyere P and Onyia, V. N. 2008. Plant spacing, dry matter accumulation and yield of local and improved maize cultivar. *Journal. American. Science*. **4**(1): 11-19.
- Iqbal, A., Ayoub. M., Zaman, H. And Ahmed, R., 2006. Impact of nutrient management and legumes association on agro qualitative traits of maize forage *Pakistan. Journal.Botany*. **38**, 1079-1084.
- Jaliya, M. M., Ibrahim, A, Babaji, B. A. Sani, B. M. and Aminu D. (2013). Effect of nitrogen and sulphur fertilizers on maize grain protein content of QPM (Quality Protein Maize) maize varieties at Samaru Zaria. *Global Journal of Bio-Science and Biotechnology*. **2**(1): 132-134.
- Jat, N. K., Kumar, A and Dhar S. 2010. Influence of sesbania green with or without wheat residues and Nitrogen fertilization on maize – wheat cropping system. *Indian Journal of Agronomy*, **55**(4): 253-258.
- Jnana Bharati Palai, NC Sarkar and Jagadish Jena 2018. Effect of zinc on growth, yields, zinc use efficiency and economics in baby corn *Journal of Pharmacognosy and Phytochemistry*; **7**(2): 1641-1645
- Jackson, M.L. 1973. Soil chemical analysis. Prentice Hall of India Pvt. Ltd., New Delhi. PP. 56.

- Kalpana, R. and Krishnarajan, J. 2002. Effect of dose and time of potassium application on yield and quality of bay corn .Agriculture Science Digest, **22**(1): 59-60
- Karki, T.B., Kumar, A.and Gautam, R .C. 2005. Influence of integrated nutrient management on growth, yield, content and uptake of nutrient and soil fertility status in maize. *Indian journal of Agriculture Science*, **75**(10): 682-685 -.
- Kumar, A. K Sagar, G. K. Chandrika, V .and Reddy, p. M. 2009. Influence of integrated nitrogen management on yield nitrogen uptake, soil fertility status and economics of baby corn. *Indian journal of Agricultural Research*, **43**(3): 227 -229.
- Kumar, A.S., Sakthivel, N., Subramanian. E., Kalpana, R., Janaki, P. and Rajesh, P. (2013). Foliar application of nutrients and plant growth regulators on growth and yield of finger millet. *Journal of Pharmacognosy and Phytochemistry*, **7**(3): 3032-3035.
- Kumar, A.S., Sakthivel, N., Subramanian. E., Kalpana, R., Janaki, P. and Rajesh, P. (2018). Foliar application of nutrients and plant growth regulators on growth and yield of finger millet. *Journal of Pharmacognosy and Phytochemistry***7**(3): 3032-3035.
- Kumar, P., Desai, B. K. and Pujari, B. T. 2007. Effect of Integrated Nutrient Management on Economics of Maize Cultivation. *Karnataka Journal. Agriculture. Science.*,**20**(4): 831-832.
- Kunjir, S.S., Pinjari, S.S., Suryavanshi, J.S. and Bhonde, T.S. 2009. Effect of planting geometry, nitrogen levels and micronutrients on growth and yield of Sweet corn. *Bioinfolet*, 6: 22-24.
- Lamana, M. C. L. 2003. Effect of spacing between plants on growth and forage yield of two maize (*Zea mays* L.) cultivar .M. Sc. (Ag.) thesis, sudan University of science and Technology, sudan, 55.
- Liu H., Gan W, Rengel, Zhao, P. (2016). Effects of zinc fertilizer rate and application method on photosynthetic characteristics and grain yield of summer maize. *Journal of Soil Science and Plant Nutrition*, **16**(2), 550-562
- Managaser,V.T. 2003 intercropped with pole sito .Population density and nitrogen fertilization of young baby cob corn in Research Journal Agriculture Department ,DMMMUS,pp.1-25.

- Marnagar, E., and Dawson. J., (2017) Effect of Biofertilizers level of Nitrogen and Zinc on growth and yield of hybrid maize (*Zea mays* L.) *International. current. Microbial. Applied.Science*6(9): 3614-3622.
- Mayank P., Saini, K.P., Yadav, V.and Yadav, R.K. (2018). Effect of foliar application of PGRson growth, yield and yield attributes of rice under salt stress condition. *Journal of Pharmacognosy and Phytochemistry* 2: 268-272.
- Md. Asaduzzaman ,Mrityunjoy Biswas, Md. Nazrul Islam, Mohammad Mokhlesur Rahman, Rafeza Begum, Md. Abdur Rahman Sarkar, Md. Asaduzzaman. 2014. Variety and N-Fertilizer Rate Influence the Growth, Yield and Yield Parameters of Baby Corn (*Zea mays* L.) *Journal of Agricultural Science*; 6(3): 1916-9760.
- Mona E. El-Azab. (2015). Increasing Zn ratio in a compound foliar NPK fertilizer in relation to growth, yield and quality of corn plant, *Journal. Innovations. Pharmaceutical. Biological science*, 2(4): 451-468.
- Moreira, J. N., Silva, P .S., Silva, L. Dombroski, J. L. D. and Castro, R. S. 2019. Effect of detasseling on baby corn, green ear and grain yield of two maize hybrids. *Horticulture Brasileira*, 28(4): 406 -411.
- Motto, M and Mall, R H. 1983.Prolificacy in maize .A review .*Maydica* 23: 53-56.
- Nagasubramaniam, A., Pathmanabhan, G. and Mallika, V. Studies on improving production potential of baby corn with foliar spray of plant growth regulators. *Annual. Plant Phsiol.*2007; 21(2):154-157.
- Nahar. K. Ahmed, S.,Akanada, M. A. L. Ondal M .A .L. Mondal, M. A. A and Islam, M .A .Genotype environment interaction for baby cob yield and maturity in baby corn .*Bangladesh journal of Agriculture Research*, 35(3), 489 -496, 2010.
- Naik, R.D., and More, S.M., (2015). Integrated nutrient management studies in finger millet. *Crop Res.*, 48(2): 27-31.
- Ochaong, P. 2005. Appropriate variety, plant density and rate and time of nitrogen fertilization

for baby corn production in Amphoe Kamphaeng saen. *Thailand Research Journal of Agriculture and Biological Sciences*,**1**(4): 303-307.

Omar, H., Al-Rawi., Adil H. Abdulkafoor, Saad I. yousif. and Mustafa R. Al-Shaheen. (2018). Effect of spraying with different levels of salicylic and humic acid in some growth characteristics and yield of wheat. *ISSN: 3*:133-138.

Olsen, S.R., Cole, V.V., Watanable, F.S. and Dean L.A. 1954. Estimation of available phosphorous in soil by extraction with sodium bicarbonate. United States Department of Agriculture, Circular **939**: 1-9.

Usadadia, P. S. Mistry, N. G. Savani and K. K. Patel. 2019 Effect of Different Levels of Irrigation, Nitrogen and Foliar Application of Banana Pseudostem Sap on Drip Irrigated Sweet Corn - Green Gram Cropping Sequence, *Indian Journal. Pure Applied. Bioscience*.**7**(5): 254-258.

Pandey, A.K., Prakash, V., Mani, V.P. and Singh, R. D. 2000. Effect of rate of nitrogen and of application on yield and economics of baby corn (*Zea mays* L.). *Indian Journal of Agronomy* **45**(2): 338-343.

Pandey, A. K., Mani, V. P., Prakash, V., Singh, R. D. and Gupta, H. S. 2002. Effect of varieties and plant densities on yield, yield attributes and economics of baby corn (*Zea mays* L.). *Indian Journal of Agronomy*. **47**: 221-226.

Panwar, A. K. and Munda, G. C. Response of baby corn (*Zea mays* L.) to nitrogen and land configuration in mid hills of Meghalaya .*Indian journal of Agricultural Science*, **76**(5), 293-296.

Paramasivan, M., Kumaresan, K, R., Malarvizhi, P., Mahimairaja, S. and Velayudham, K. 2010. Effect of different levels of NPK and zinc on yield and nutrient uptake of hybrid maize in Mayamankuruchi series of soil of Tamil nadu. *Asian journal of soil Science*, **5**(1): 157 - 161.

Prabha, A., Parasuraman, P., Sivagamy K. and Sivakumar, B. (2016). Growth, yield and economics of irrigated finger millet as influenced by system of finger millet

- intensification (SFI) practices in north eastern zone of Tamil Nadu. *Journal of Pharmacognosy and Phytochemistry*, **8**(3): 600-663.,
- Bala, S. and Khoyumthem, P. 2007. Estimation of yield and economic return in baby corn. *Envirnment and Ecology*,**25**(4): 945-047,
- Rakesh Kumar, Narendra Kumawat, Sudhir Kumar, Amitesh Kumar Singh and J.S. Bohra. 2017. Effect of NPKS and Zn Fertilization on, Growth, Yield and Quality of Baby Corn-A Review. *International Journal of Current Microbiology and Applied Sciences*, ISSN: 2319-7706. **6**(3): 1392-1428.
- Ram, V., Singh, R.N. and Singh, K. 2006. Studies on integrated use of FYM, nitrogen and sulphur on growth, yield attributes and yield on winter maize (*Zea mays* L.). *Plant Archi*, 6:749-752.
- Ramchandruppa, B. K., Nanjappa, H .V. and Soumya, T. M. 2010. Effect of stages of harvest on yield and quality of baby corn varieties. *Mysore journal of Agriculture Science*: **40**(4): 453-457.
- Ramesh S, P. Sudhakar, S. Elankavi, K. Suseendran. and Jawahar, S. (2019). Effect of Gibberellic acid (GA3) on growth and yield of rice (*Oryza sativa* L.). *Plant Archives*; **19**: 2581-6063.
- Rao, L. K., kumar, R. A. and Lal, G. M. 2009. Effect of Integrated nitrogen management on growth and yield of baby corn (*Zea mays* L.) cv. Mridula. *journal of Maharastra Agriculture University*, **34**(3): 249-251.
- Richards, L.A. 1954. Diagnosis and improvements of saline and alkali soils. Agriculture Handbook No. 60.USDA, Washington*
- Sahoo, S. C. and Panda, M. M. 1999. Effect of level of nitrogen and plant population on yield of baby corn (*Zea mats* L.). *Indian Journal of Agriculture Sciences* **69**(2): 157-158.
- Santosh Kumar Singh, SN Suman and Aradhna Kumari. (2018). Performance of autumn maize

crop as influenced by seaweed saps. *International journal of Chemical Studies*, **6**(2): 2341-2345, for sea weed uses.

Sahoo,S.C.Yield and economics of baby corn (*Zea mays* L.) as affected by varieties and levels of nitrogen .*Range Management and Agroforestry*, **32**(2): 135-137,2011.

Sarakhsi ,H S ,.Yarnia ,M and Amirniya ,R .Effect of nitrogen foliar application in different concentration and growth stage of baby corn (hybrid -704) *Advance in Environmental Biology* ,**4**(2) ,291-298 2010.

Sathishkumar, A. N., Sakthivel, E. Subramanian, R. Kalpana, P, Janaki. and P, Rajesh. (2018). Foliar application of nutrients and plant growth regulators on growth and yield of finger millet. *Journal of pharmacognosy and phytochemistry*; **7**(3): 3032-3035.

Scaria Dona, Rajasree G and Sudha B of baby corn (*Zea mays*). (2016). Effect of varieties and spacing on growth, yield and economics of cultivation of baby corn (*Zea mays* L.) as inter crop in coconut garden. *Research on crops*.**17**(4): 673-678.

Shafea,L and Saffari ,M (2007). Effect of zinc ($ZnSO_4$) and nitrogen on chemical composition of maize grain. *International Journal Agriculture. Science*.**1**(6): 323-328.

Shah S, Ghani G, Khan H, Arif M.,Qahar, Inamullah, Ali A. and Ahmad M.,(2015) response of maize cultivars to phosphorus and zinc nutrition pak. *journal. botany.*, 47(si): 289-292.

Shaikh Wasim Chand, R Susheela, D Sreelatha, M Shanti and SA Hussain. 2017. Effect of zinc fertilization on yield and economics of baby corn (*Zea mays* L.) *Journal of Pharmacognosy and Phytochemistry*; **6**(5): 989-992.

Shaikh Wasim Chand, R Susheela, D Sreelatha, M Shanti and SA Hussain Effect of zinc fertilization on yield and economics of baby corn (*Zea mays* L.) *Journal of Pharmacognosy and Phytochemistry* 2017; **6**(5): 989-992

Sharma, S. K., Swami, A A., and Singh R.K., (1992). Relative response of maize (*Zea mays* L.) varieties to zinc .*Indian Journal . Agronomy.*,**37**(2): 361-361.

- Siam, H.S., Kader, E.M.G.A. and Alia, E.H.I. 2008. Yield and yield component of maize as affected by different sources and application rates of N fertilizers. *Research. Journal. Agriculture. Biological. Science.*, 6: 399-412.
- Singh, M. K. Singh. R. N., Singh S, P. S. P. Yadav M. K. and Singh, V. K. 2010. Integrated nutrient management and for higher yield quality and profitability of baby corn (*Zea mays* L.) *Indian journal of Agronomy*, **55**(2), 100-104,
- Singh, D. P., Rana,N.S. and Singh R. P.,(2000). Growth and yield of winter maize as influence as intercrops and nitrogen application. *Indian J. Agronomy.*,**45**(3): 515-519.
- Singh, N.T., Viv, A. C. and Singh, R. 1985. Nitrogen response of maize under temporary flooding. *Nutrient Cycling in Agrosystem*, **6**(2); 11-12.
- Siva, K., Shinggu C.P., Dadari, S.A., Shebayan, J.A.Y., Adekpe, D.I., Mahadi, M.A., Mukhtar, A. and SivaKumarR,G.PaThmanaban, M. K. Kalarani, MallikaVanangamudi and P.S.Srinivasan. 2002. Effect of Foliar Application of Growth Regulators on Biochemical Attributes and Grain Yield in Pearl Millet. *Indian Journal Plant Physiol.*, **7**(1): 79-82.
- Sobhana, V., Kumar, A., Idnani, L. K., Singh,I. and Shivadhar. 2012. Plant population and nutrient requirement for baby corn hybrids (*Zea mats* L.). *Indian journal of Agronomy*,**57**(3): 294-296.
- Srinivasan K. (1992). Effect of amendment and zinc levels on growth and yield of maize (*Zea mays* L.) *Indian Journal Agriculture*.**37** (2): 246-249.
- Suresh, G., Guru, G. and Lokanadan, S. (2018). Effect of Nutrient Levels and Plant Growth Regulators on Growth Parameters of Pearl Millet. *Int. J. Pure App.Biosci.***6** (3): 271-277.
- Tamrakar, S.K., Singh, P., Kumar, V. and Tirkey, T. (2018). Effect of gibberellic acid, salicylic acid, cow urine and vermiwash on corn production of *Gladiolus* cv. *Candyman*. *International Journal Current Microbiol Applied Science*. **6**: 677-686.
- Thakur, D R., O M., and Kharawara, P.C. 1998. Effect of nitrogen and plant spacing on yield, Nitrogen uptake and economics in baby corn (*Zea mays* L.).*Indian Journal of Agronomy*

43(4):668-71.

Thakur, D. R. and Sharma, V. 1999. Effect of varying rates of nitrogen and its schedule of split application in baby corn (*Zea mays* L.). *Indian journal of Agricultural Science*, **69**(2): 93-95.

Thakur, D. R. Prakash, O M., Kharwara P. C. and Bhalla, S. K. 1997. Effect of nitrogen and plant spacing on growth, development and yield of baby corn. *Indian journal of Agronomy*, **42**(3): 479 -483.

UNDER PEER REVIEW