

## Original Research Article

### **“Effect of micronutrient application on growth, yield and quality of green gram growing Iron and Zinc deficient soils of Renapur Tahsil, Latur.”**

#### **Abstract**

The present investigation entitled “Effect of micronutrient application in green gram growing iron and zinc deficient soils of Renapur Tahsil, Latur” during *Kharif* season of the year, 2022-2023 at AFieldexperimentwasconductedat farmers field At. Post-Dawangaon Tq-Renapur Dist-Latur. The experiment was layout in RBD with three replications and a recommended variety of green gram BM 2003-2 as a test crop along with ten treatments. Among the various treatments the application of RDF + 25 kg ha<sup>-1</sup> FeSO<sub>4</sub> + 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub> (T<sub>7</sub>) significantly increased the growth parameters *viz.*, highest plant height (52.19 cm at 60 DAS), number of branches(10.64 at 60 DAS), leaf area(540.67 cm<sup>2</sup> plant<sup>-1</sup> at 60 DAS), dry matter content(9.54 g plant<sup>-1</sup> at 60 DAS), total chlorophyll content(9.93 mg g<sup>-1</sup> at 45 DAS) and number of nodules per plant (19.70), Yield parameters *viz.*, higher number of pods plant<sup>-1</sup>(15.43), number of grains pod<sup>-1</sup>(11.88), seed yield kg ha<sup>-1</sup>(1575.46) and straw yield kg ha<sup>-1</sup>(1864.68) at harvest and the improvement in quality parameters *i.e.* test weight (35.23 g), protein content (22.37 %) and protein yieldkg ha<sup>-1</sup> (347.27) at harvest and was found at par with treatments T<sub>4</sub> (RDF + S.A. Grade- I micro-nutrient @ 25 kg ha<sup>-1</sup> + F.A. Grade- II micro-nutrient @ 0.5 % at 25 and 40 DAS) and treatment T<sub>2</sub> (RDF + S.A. Grade- I micro-nutrient @ 25 kg ha<sup>-1</sup>).

**(Keywords: green gram, iron, zinc, growth, quality and yield)**

#### **1. Introduction**

Green gram (*Vigna radiata* L.) is a well-known domesticated legume crop grown widely all over the world providing an important economy for marginal farmers in the developing countries and a high-value commodity crop with high nutritional quality in the developed nations. Green gram comes under the family *Fabaceae*, subfamily *Papilionaceae*, genus *Vigna* and species *radiata*. This family is widespread as it occupies the third largest family of flowering plants with approximately 650 genera and nearly 20,000 species (Doyle, 1994). Green gram is alternatively known as golden gram, mungbean, moong bean, haricot mungo, mash bean, etc. It is an annual, semi-erect to erect or sometimes twining, up to 100 cm tall, deep-rooted herbaceous plant. In India, green gram is mostly grown in Andhra Pradesh, Maharashtra, Orissa, Rajasthan,

Gujarat, Punjab, Uttar Pradesh, etc. India ranks first in both area and production of all important pulses grown in the world. Pulses are grown on about 30.4 million ha. area in India with production of 14.77 million tonnes and productivity pulses is  $617 \text{ kg ha}^{-1}$ . The total area under pulses in Maharashtra is 32.69 lakh ha, with a total production of 21.44 lakh tones and productivity of  $217 \text{ kg ha}^{-1}$ . Green gram ranks third among all the pulses in India after chickpea and pigeon pea. India accounts for almost 65 percent area and 54 percent production of world mung bean. Green gram grown on about 3.57 million ha. Area in India with a total production of 17.89 metric tonnes and productivity of green gram is  $500 \text{ kg ha}^{-1}$  (Anonymous, 2021).

Micronutrient deficiencies in soil and crop plants are widespread because of increased micronutrient demand from intensive cropping practices and adaptation of high-yielding crop cultivars, enhanced crop production on marginal soils that contain low levels of essential micronutrients, increased use of high analysis fertilizers with low amounts of micronutrients, decreased use of animal manures, composts, and crop residues, use of soils low in micronutrient reserves, use of liming in acid soils, involvement of natural and anthropogenic factors that limit adequate supplies and create elemental imbalance in soil. Boradkar *et al.* (2023) reported that as much as 48, 12, 5, 4, 33, 13, and 41 percent of soils in India are affected by deficiency of Zn, Fe, Mn, Cu, B, Mo, and S, respectively. In general, farmers applied only major nutrients and not the micronutrients are lacking. Micronutrients like Iron, Zinc, Manganese, Copper, Molybdenum, and Boron play an important role in increasing legume yield through their effect on the plant itself, nitrogen-fixing symbiotic process and effective use of major and secondary nutrients. However, they are used in lower amounts as compared to macronutrients. They have a major role in cell division, development of meristematic tissues, photosynthesis, respiration, and acceleration of plant maturity. Nowadays micronutrient deficiencies are found limiting factors for crop growth and optimum yield. Hence, optimum yield potential attained through nutrient management including micronutrient iron and zinc is a basic requirement for major crops and mung bean.

## 2. Materials and Methods

A field experiment was conducted at the farmer's field of Shri. Prabhakar Ramrao Nagargoje, At/Post-Dawangaon Tq-Renapur Dist- Latur, during *Kharif*, 2022-2023, college of Agriculture, Latur, Maharashtra. It lies between  $76^{\circ}30'31''$  E Longitude and  $18^{\circ}35'26''$  N Latitude. With treatments and three replications in randomized block design. T<sub>1</sub>: RDF, T<sub>2</sub>: RDF + S.A.

Grade-I micro-nutrient @ 25 kg ha<sup>-1</sup>. T<sub>3</sub>: RDF + F.A. Grade-II micro-nutrient @ 0.5 % at 25 and 40 DAS. T<sub>4</sub>: RDF + S.A. Grade-I micro-nutrient @ 25 kg ha<sup>-1</sup> + F.A. Grade-II micro-nutrient 0.5 % at 25 and 40 DAS. T<sub>5</sub>: RDF + S.A. FeSO<sub>4</sub> @ 25 kg ha<sup>-1</sup>. T<sub>6</sub>: RDF + S.A. ZnSO<sub>4</sub> @ 25kg ha<sup>-1</sup>. T<sub>7</sub>: RDF + S.A. FeSO<sub>4</sub> + ZnSO<sub>4</sub> @ 25kg ha<sup>-1</sup>. T<sub>8</sub>: RDF + F.A. FeSO<sub>4</sub> @ 0.5 % at 25 to 40 DAS. T<sub>9</sub>: RDF + F.A. of ZnSO<sub>4</sub> @ 0.5 % at 25 to 40 DAS. T<sub>10</sub>: RDF + F.A. FeSO<sub>4</sub> + ZnSO<sub>4</sub> @ 0.5 % at 25 to 40 DAS. *i. e.* Recommended Dose of Fertilizers N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O (25:50:25 kg ha<sup>-1</sup>) + FYM were applied to green gram. The experimental soil was deep, black in colour, and good drainage, clayey in texture, highly calcareous in nature (18.45 %), moderately alkaline reaction (8.38 pH), and low in content of organic carbon (2.41 g kg<sup>-1</sup>), available nitrogen (165.55 kg ha<sup>-1</sup>), available phosphorous (17.13 kg ha<sup>-1</sup>), high in available potassium (435.45 kg ha<sup>-1</sup>), deficient in DTPA iron (2.16 mg kg<sup>-1</sup>), zinc(0.48 mg kg<sup>-1</sup>) sufficient in DTPA copper and manganese. Soil sample was collected before sowing of green gram and analysed as per standard methods. The data on growth parameters, quality parameters were recorded at 30, 45 and 60 DAS and yield parameters recorded at harvest stage. The data on various parameters recorded during the period of investigation were tabulated and statistically analysed (Panse and Sukhatme, 1996).

### 3.1 Results and discussion

#### Growth parameters

The data related to different growth parameters like plant height, number of branches, leaf area, chlorophyll content, dry matter and number of nodules as influenced by the application of micronutrient sources with certain treatments is shown below.

#### 3.1.1. Plant height

Data presented in Table 1 revealed that the plant height of green gram was significantly increased due to soil application of micronutrients in combination. The highest plant height was recorded with treatment T<sub>7</sub> (RDF + Soil application FeSO<sub>4</sub> + ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup>), increased the plant height of green gram by 39.70 cm, 46.87 cm and 52.19 cm at 30, 45 DAS and at harvest respectively, and was found at par with treatment T<sub>4</sub> and treatment T<sub>2</sub>. The treatment T<sub>1</sub> (RDF) recorded the lowest plant heights 28.60, 33.17 and 38.97 cm at 30, 45 DAS and at harvest.

Application of micronutrients as soil application along with foliar application might have enhanced the availability of nutrients, activation of nutrients increases the vegetative growth and higher chlorophyll content in the respective treatment that would help to increase the plant

height. Similar results were recorded by Gidaganti *et al.* (2019) in green gram. Ajjannavar *et al.* (2021) in chickpea. Almad *et al.* (2020) noticed that in pigeon pea.

### 3.2 Number of branches per plant

The data regarding the number of branches per plant of green gram as affected by application of micronutrient was recorded during three growth stages at 30, 45 DAS and at harvest are presented in table .1.

Branching is a crucial aspect of crop growth, bearing the plant's pods and ultimately increasing crop yield. The findings showed that the application of micronutrients might have a significant impact on the number of branches plant<sup>-1</sup> in green gram. The maximum number of branches per plant (5.93, 9.47 and 10.64 at 30, 45 DAS and at harvest) were observed in the treatment T<sub>7</sub> (RDF + 25 kg ha<sup>-1</sup> FeSO<sub>4</sub> + 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub> kg ha<sup>-1</sup>) respectively and was found at par with treatment T<sub>4</sub> and treatment T<sub>2</sub> and significantly superior over rest of the treatments. A minimum number of branches per plant (3.57, 6.27 and 7.87 at 30, 45 DAS and at harvest) were observed in treatment T<sub>1</sub> (RDF).

This might be due to the superiority of iron and zinc sulphate in maintaining higher iron and zinc concentrations in the rhizosphere. The beneficial role of synthesis of IAA, metabolism of auxin, biological activity, stimulating effect on photosynthetic pigments and enzyme activity in turn encourage vegetative growth of plants. Similar results were also reported by Almad *et al.* (2020) in pigeon pea, Ajjannavar *et al.* (2021) in chickpea. Sudhanshu *et al.* (2022) in green gram.

**Table 1: Effect of micronutrient application on plant height and Number of Branches (plant<sup>-1</sup>) of green gram.**

Treatments	Plant height (cm)			Number of Branches (plant <sup>-1</sup> )		
	30 DAS	45 DAS	At Harvest	30 DAS	45 DAS	At harvest

<b>T<sub>1</sub></b>	28.60	33.17	38.97	3.57	6.27	7.87
<b>T<sub>2</sub></b>	38.73	45.87	50.48	5.50	9.27	10.16
<b>T<sub>3</sub></b>	34.95	38.90	48.15	4.10	8.53	9.51
<b>T<sub>4</sub></b>	38.94	46.83	50.89	5.30	9.34	10.17
<b>T<sub>5</sub></b>	37.40	43.55	48.17	5.37	9.27	9.50
<b>T<sub>6</sub></b>	34.48	40.01	46.35	4.97	9.10	9.47
<b>T<sub>7</sub></b>	39.70	46.87	52.19	5.93	9.47	10.64
<b>T<sub>8</sub></b>	34.25	38.40	45.58	4.13	8.23	8.73
<b>T<sub>9</sub></b>	33.41	39.30	46.10	3.87	7.73	8.38
<b>T<sub>10</sub></b>	34.26	39.79	46.23	3.93	8.13	9.20
<b>SE(m)±</b>	<b>1.28</b>	<b>0.94</b>	<b>1.18</b>	<b>0.28</b>	<b>0.30</b>	<b>0.33</b>
<b>CD at 5%</b>	<b>3.81</b>	<b>2.79</b>	<b>3.51</b>	<b>0.85</b>	<b>0.91</b>	<b>0.98</b>

### 3.3 Leaf area

The data regarding leaf area of green gram as affected by application of micronutrient was recorded during three growth stages at 30, 45 DAS and at harvest are presented in table 2.

The highest leaf area  $\text{cm}^2 \text{ plant}^{-1}$  485.07, 532.97 and 540.67 at 30, 45 DAS and at harvest was recorded with treatment T<sub>7</sub> (RDF + 25 kg ha<sup>-1</sup> FeSO<sub>4</sub>+25 kg ha<sup>-1</sup> ZnSO<sub>4</sub> kg ha<sup>-1</sup>) respectively and was found at par with treatment T<sub>4</sub> and treatment T<sub>2</sub> and significantly superior over rest of the treatments in green gram. The lowest leaf area 315.57, 326.41 and 344.33 at 30, 45 DAS and at harvest was recorded with treatment T<sub>1</sub> (RDF).

This might be due to increased metabolic activity by the increased supply of nutrients, more dry matter accumulation in leaves helped the photosynthetic area to remain active for a longer period. Similar kinds of results were noticed by Almad *et al.* (2020) in pigeon pea. Gowda *et al.* (2014) in pigeon pea.

### 3.4 Dry matter content

The data on dry matter content  $\text{g plant}^{-1}$  of green gram as affected by application of micronutrient was recorded during three growth stages at 30, 45 DAS and at harvest are represented in table 2.

The higher dry matter was observed with the treatment  $T_7$  (RDF +  $25 \text{ kg ha}^{-1} \text{ FeSO}_4 + 25 \text{ kg ha}^{-1} \text{ ZnSO}_4 \text{ kg ha}^{-1}$ ) 4.78, 7.27 and  $9.54 \text{ g plant}^{-1}$  at 30, 45 DAS and at harvest respectively and was found at par with treatment  $T_4$  and treatment  $T_2$  and significantly superior over rest of the treatments. Further data revealed that the treatment  $T_1$  (RDF) produced lower dry matter content 2.95, 5.41, and  $7.17 \text{ g plant}^{-1}$  at 30, 45 DAS and at harvest respectively.

Higher dry matter production might be due to the increased photosynthetic rate accelerated nutrient uptake from soil, increased availability of major nutrients, and micronutrient (Zn and Fe) which increases vegetative growth and plant height. Similar findings were reported by Sudhanshu *et al.* (2022) in green gram. Ajjannavar *et al.* (2021) in chickpea and Meena *et al.* (2013) in mung bean.

**Table .2: Effect of micronutrient application on Leaf Area ( $\text{cm}^2 \text{ plant}^{-1}$ ) and Dry Matter Content ( $\text{g plant}^{-1}$ ) of green gram.**

Treatments	Leaf Area ( $\text{cm}^2 \text{ plant}^{-1}$ )			Dry Matter Content ( $\text{g plant}^{-1}$ )		
	30 DAS	30 DAS	30 DAS	30 DAS	45 DAS	At harvest
$T_1$	315.57	2.95	2.95	2.95	6.27	7.87
$T_2$	484.36	4.49	4.49	4.49	9.27	10.16
$T_3$	411.43	3.84	3.84	3.84	8.53	9.51
$T_4$	495.77	4.58	4.58	4.58	9.34	10.17
$T_5$	492.40	4.23	4.23	4.23	9.27	9.50
$T_6$	406.30	3.90	3.90	3.90	9.10	9.47
$T_7$	485.07	4.78	4.78	4.78	9.47	10.64
$T_8$	415.33	3.70	3.70	3.70	8.23	8.73
$T_9$	399.03	3.48	3.48	3.48	7.73	8.38
$T_{10}$	419.10	3.77	3.77	3.77	8.13	9.20
SE(m) $\pm$	<b>14.14</b>	<b>0.19</b>	<b>0.19</b>	<b>0.19</b>	<b>0.30</b>	<b>0.33</b>
CD at 5%	<b>42.01</b>	<b>0.57</b>	<b>0.57</b>	<b>0.57</b>	<b>0.91</b>	<b>0.98</b>

### 3.5 Chlorophyll content

The data about total chlorophyll analysed at 45 DAS are presented in Table .3.

Among all the treatments, T<sub>7</sub> (RDF + 25 kg ha<sup>-1</sup> FeSO<sub>4</sub> + 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub> kg ha<sup>-1</sup>) resulted in significantly higher chlorophyll content (9.93 mg g<sup>-1</sup>) at 45 DAS respectively and at par with treatment T<sub>4</sub> and treatment T<sub>2</sub> and significantly superior over rest of the treatments. The lowest chlorophyll content was observed with control T<sub>1</sub> RDF (6.39 mg g<sup>-1</sup>) at the flowering stage of the crop.

These results might be due to chlorophyll being a green pigment in the chloroplast of the green plants. It helps plants to create their food through photosynthesis. Iron plays a crucial role during electron transport and chlorophyll formation and zinc also plays an important role in the synthesis of chlorophyll its enzymatic role in starch formation and protein synthesis. This increase in the availability of zinc to plants might have stimulated the metabolic and enzymatic activities thereby increasing the growth of the crop. Similar findings were reported by Lokhande (2018) in green gram. Bhadru *et al.* (2019) and Meena *et al.* (2013) in green gram crop.

### 3.6 Nodulation

The data about Nodules at flowering stage (plants<sup>-1</sup>) are presented in Table .3.

The maximum number of nodules per plant was obtained under the treatment T<sub>7</sub> (RDF + 25 kg ha<sup>-1</sup> FeSO<sub>4</sub> + 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub> kg ha<sup>-1</sup>) which was (19.70) respectively and were found at par with treatment T<sub>4</sub> and treatment T<sub>2</sub> and significantly superior over rest of the treatments. The minimum number of nodules per plant was obtained under the treatment T<sub>1</sub> (RDF) which was (11.57) respectively.

The increase in several nodules might be due to increased rhizobial colonization in the rhizosphere because of the increased availability of iron, zinc plays a vital role in cellular growth, differentiation and metabolism which results in vigorous growth of plants and extensive root system leading to increased growth parameters. Similar results were also reported by Kumar *et al.* (2020) in black gram and Meena *et al.* (2013)

**Table.3: Effect of micronutrient application on Total chlorophyll Content (mg g<sup>-1</sup>) and Root nodules at flowering stage (plants<sup>-1</sup>) of green gram.**

Treatments	Total chlorophyll Content (mg g <sup>-1</sup> )	Root nodules at flowering stage (plants <sup>-1</sup> )
T <sub>1</sub>	6.39	11.57
T <sub>2</sub>	9.31	19.50
T <sub>3</sub>	7.53	16.67
T <sub>4</sub>	9.71	19.83
T <sub>5</sub>	7.46	18.38
T <sub>6</sub>	7.07	17.57
T <sub>7</sub>	9.93	19.70
T <sub>8</sub>	7.06	16.30
T <sub>9</sub>	7.08	14.93
T <sub>10</sub>	7.20	16.43
SE(m)±	<b>0.43</b>	<b>0.51</b>
CD at 5%	<b>1.28</b>	<b>1.54</b>

### Yield parameters

The data related to different yield parameters like number of pods plant<sup>-1</sup>, number of grains pod<sup>-1</sup>, seed and straw yield (kg ha<sup>-1</sup>) as influenced by the application of micronutrient sources with certain treatments has shown below.

#### 3.7 Number of pods plant<sup>-1</sup>

Data on the mean number of pods per plant of green gram as influenced by various micronutrient treatments are presented in Table 4

There was a significant increase in the number of pods plant<sup>-1</sup> with all the treatments over T<sub>1</sub> (RDF). The highest number of pods plant<sup>-1</sup> was recorded with treatment T<sub>7</sub> (RDF + 25 kg ha<sup>-1</sup> FeSO<sub>4</sub> + 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub>) at harvest (15.43) respectively which was found at par with treatment T<sub>4</sub> and T<sub>2</sub> having a number of pods per plant 15.35 and 15.33, respectively.

These results are in confirmatory with Banoth *et al.* (2022) in green gram, Gidaganti *et al.* (2019) in green gram and Meena *et al.* (2013) with green gram crop.

#### 3.8 Number of grains pod<sup>-1</sup>

Data presented in table .4 were not affected significantly with respect to the number of grains pod<sup>-1</sup> at the harvest stage.

The maximum number of grains  $\text{pod}^{-1}$  observed in treatment T<sub>7</sub> (RDF + 25 kg  $\text{ha}^{-1}$   $\text{FeSO}_4$  + 25 kg  $\text{ha}^{-1}$   $\text{ZnSO}_4$  kg  $\text{ha}^{-1}$ ) which were 11.88 grains  $\text{pod}^{-1}$  respectively and the minimum number of grains  $\text{pod}^{-1}$  (10.92) were observed in the treatment T<sub>1</sub> (RDF) in green gram.

The results obtained might be due to the application of micronutrients, iron and zinc which are playing an accumulative effect of growth characters and also yield attributes as it contains zinc which is involved in IAA synthesis and also different metabolic processes in plants. IAA promotes the prevention of pod abscission and cell elongation at suppression of pod abscission of pod was the major determining factor of the seed yield. On the other hand, auxin indirectly controlled the ethylene activity which accelerated the abscission. It also suppressed the cellulase activity, a cell-degrading enzyme that favored the abscission process. These results are in line with Gidaganti *et al.* (2019) in green gram, Banoth *et al.* (2022) in green gram, Kareti and George (2017) in chickpea and Sudhanshu *et al.* (2022) in green gram.

### 3.9 Seed yield

The data on the seed yield of green gram as influenced by the application of micronutrients are presented in table .4.

The seed yield of green gram ranged between 1020.32 kg  $\text{ha}^{-1}$  to 1575.46 kg  $\text{ha}^{-1}$ . The highest seed yield (1575.46 kg  $\text{ha}^{-1}$ ) was recorded in treatment T<sub>7</sub> (RDF + 25 kg  $\text{ha}^{-1}$   $\text{FeSO}_4$  + 25 kg  $\text{ha}^{-1}$   $\text{ZnSO}_4$  kg  $\text{ha}^{-1}$ ) respectively which was at par with treatment T<sub>4</sub> and T<sub>2</sub> and significantly superior over rest of the treatments. Whereas, the lowest seed yield (1020.32 kg  $\text{ha}^{-1}$ ) was recorded in the treatment T<sub>1</sub> (RDF) in green gram.

The increase in yield might be due to fulfillment of the demand for the crop by higher assimilation and translocation of photosynthates from source to sink and better role of iron and zinc during the reproductive phase of crop growth. However, the combined effect of iron and zinc provided sufficient nutrition to the plant and thereby more yield attributes and yield were recorded.

Similar findings were observed by Gidaganti *et al.* (2019) in green gram. Sudhanshu *et al.* (2022) in green gram. Meena *et al.* (2013) in green gram crop. Gahlot *et al.* (2020) in mungbean crop. Ajjannavar *et al.* (2021) in chickpea.

### 3.10 Straw yield

The data on the straw yield of a green gram as influenced by the application of micronutrients are presented in table .4.

Straw yield is directly related to an increase in vegetative growth of the plants. Data indicated that the application of iron and zinc markedly influenced the straw yield in the range of (1238.41 kg ha<sup>-1</sup> to 1864.68 kg ha<sup>-1</sup>). The maximum straw yield (1864.68 kg ha<sup>-1</sup>) was recorded in treatment T<sub>7</sub> (RDF + 25 kg ha<sup>-1</sup> FeSO<sub>4</sub> + 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub> kg ha<sup>-1</sup>) respectively which was found at par with treatment T<sub>4</sub> and treatment T<sub>2</sub> and significantly superior over rest of the treatments. Whereas, the lowest straw yield (1238.41 kg ha<sup>-1</sup>) was recorded in the treatment T<sub>1</sub> (RDF).

The higher straw yield might be due to improved vegetative growth and growth parameters through adequate availability of major and micronutrients in soil, which in turn, favorably influenced the physiological process and buildup of photosynthates. Similar findings were shown by Misal (2018) in green gram, Meena *et al.* (2013) in green gram crop and Gahlot *et al.* (2020) with mung bean crop.

**Table 4: Effect of micronutrient application on Yield parameter of green gram.**

Treatments	No. of pods (plant <sup>-1</sup> )	No. of grains (pod <sup>-1</sup> )	Seed yield (kg ha <sup>-1</sup> )	Straw yield (kg ha <sup>-1</sup> )
T <sub>1</sub>	11.19	10.92	1020.32	1238.41
T <sub>2</sub>	15.33	11.51	1362.58	1820.32
T <sub>3</sub>	14.05	11.21	1094.42	1401.73
T <sub>4</sub>	15.35	11.69	1414.87	1832.96
T <sub>5</sub>	14.48	11.45	1236.93	1658.49
T <sub>6</sub>	14.42	11.32	1183.40	1594.30
T <sub>7</sub>	15.43	11.88	1575.46	1864.68
T <sub>8</sub>	13.23	11.15	1038.66	1341.26
T <sub>9</sub>	13.29	11.48	1033.46	1381.41
T <sub>10</sub>	13.44	11.00	1063.20	1365.30
SE(m)±	<b>0.04</b>	<b>0.21</b>	<b>104.95</b>	<b>83.05</b>

CD at 5%	0.12	NS	311.81	246.78
----------	------	----	--------	--------

### Quality parameters

The data related to different quality parameters Test weight, Protein content and Protein yield ( $\text{kg ha}^{-1}$ ) as influenced by the application of micronutrient sources with certain treatments is shown below.

#### 3.11 Test weight

Data regarding the effect on test weight was tabulated in table .5 indicating that the application of micronutrients markedly influenced the test weight in the range of (32.01 to 35.23 g).

Test weight was not significantly influenced by the application of iron and zinc and the maximum test weight (35.23 g) was found under the treatment T<sub>7</sub> (RDF + 25  $\text{kg ha}^{-1}$  FeSO<sub>4</sub> + 25  $\text{kg ha}^{-1}$  ZnSO<sub>4</sub>  $\text{kg ha}^{-1}$ ) respectively which was highest over rest of the treatments. Whereas, the lowest test weight (32.01 g) was recorded in the treatment T<sub>1</sub> (RDF) in green gram.

An increase in test weight due to greater mobilization of photosynthesis to the developing seeds by application of micronutrients might be the reason for the increase in seed weight. An increase in this attribute by soil application might be due to the involvement of iron and zinc in enzyme activation, membrane integrity, chlorophyll formation, stomatal balance and starch utilization at early stages which enhanced accumulation of assimilate in the grains resulting in heavier grains.

Similar findings were recorded by Misal (2018) in green gram, Gidaganti *et al.* (2019) in green gram.

#### 3.12 Protein content and protein yield

The data on protein content and protein yield of green gram are presented in table .5. The mean protein content was (22.37 %). The mean protein content was not affected significantly due to different micronutrient treatments. The numerically higher protein content (22.37 %) recorded with treatment T<sub>2</sub> (RDF + S.A. Grade-I micro-nutrient @ 25  $\text{kg ha}^{-1}$ ) respectively followed by treatment T<sub>4</sub> (22.35 %), T<sub>5</sub> (22.25 %), T<sub>6</sub> (22.18 %), T<sub>7</sub> (22.04 %) respectively significantly superior over rest of the treatments. While, minimum protein content (21.37 %) was recorded at treatment T<sub>1</sub> (RDF) in green gram.

**Table No.5: Effect of micronutrient application on number of pods, number of grains, of green gram.**

Treatments	Test weight ( g )	Protein content (%)	Protein yield (kg ha <sup>-1</sup> )
T <sub>1</sub>	32.01	21.37	218.02
T <sub>2</sub>	35.04	22.37	304.59
T <sub>3</sub>	34.95	21.79	238.62
T <sub>4</sub>	35.11	22.35	315.33
T <sub>5</sub>	34.96	22.25	275.19
T <sub>6</sub>	34.30	22.18	262.82
T <sub>7</sub>	35.23	22.04	347.27
T <sub>8</sub>	33.36	21.52	223.60
T <sub>9</sub>	33.76	21.58	223.00
T <sub>10</sub>	34.47	21.68	230.55
<b>SE(m)±</b>	<b>0.73</b>	<b>0.25</b>	<b>27.77</b>
<b>CD at 5%</b>	<b>NS</b>	<b>NS</b>	<b>67.65</b>

The higher protein yield (347.27 kg ha<sup>-1</sup>) was observed with treatment T<sub>7</sub> (RDF + 25 kg ha<sup>-1</sup> FeSO<sub>4</sub> + 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub> kg ha<sup>-1</sup>) respectively which was found at par with treatment T<sub>4</sub> (315.33 kg ha<sup>-1</sup>), T<sub>2</sub> (304.59 kg ha<sup>-1</sup>), and significantly superior over rest of the treatments. Whereas, a lower protein yield (218.02 kg ha<sup>-1</sup>) was recorded in treatment T<sub>1</sub> (RDF) in green gram. The increase in protein content might be due to iron and zinc which are two important elements in enzyme structure involved in amino acids are the bases of protein synthesis, protein content increases with the application of iron and zinc. Similar findings were observed by Misal *et al.* (2018) in green gram and Lokhande *et al.*(2018) in green gram.

#### 4. Conclusion

1. Green gram crop fertilized with the application of RDF + FeSO<sub>4</sub> + ZnSO<sub>4</sub>@25 kg ha<sup>-1</sup> improved growth parameters like plant height, number of branches plant<sup>-1</sup>, leaf area, dry matter content, total chlorophyll content and number of root nodules.
2. Significantly higher, number of pods plant<sup>-1</sup>, number of grains pod<sup>-1</sup>, seed yield plot<sup>-1</sup>, seed yield ha<sup>-1</sup>, straw yield plot<sup>-1</sup>, and straw yield ha<sup>-1</sup> with the application of RDF + 25 kg ha<sup>-1</sup> FeSO<sub>4</sub> + 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub>.
3. The improvement in quality parameters was recorded with the application of RDF + 25 kg ha<sup>-1</sup> FeSO<sub>4</sub> + 25 kg ha<sup>-1</sup> ZnSO<sub>4</sub>.i.e. Test weight, protein content, protein yield.

### References

- 1 Ajjannavar, D.B., Yadahalli, V.G., Sarwad, I.M. & Patil, S.B. (2021). Growth and yield of chickpea (*Cicer arietinum* L.) and soil fertility as influenced by micronutrients and bio-fertilizers in vertisols. *Journal of Farm Science*. 34(4), 404-409.
2. Almad, S.R., Pandit S.R., Rachappa, V., Dodamani B.M. & Ananda, N. (2020). Growth, yield and economics of pigeonpea as influenced by biofortification of zinc and iron. *International Journal of Current Microbiology and Applied Sciences*. 9(2), 3088-3097.
3. Anonymous (2021). Annual report on pulses, Agric. Statistics at a glance. www.dacnet.nic.in/ean.
4. Banoth, M. K., Kumar, H. S., Guguloth, P., Naik, V. M., & Umesha, C. (2022). Influence of boron and zinc on growth and yield of green gram (*Vigna radiata* L.). *The Pharma Innovation Journal*. 11(3), 1674-1678.
5. Bhadru, P., Yadav, S.S., Bijarnia, A., Kumawat, R. & Choudhary, R. (2019). Effect of zinc and thiourea application on growth, yield and nutrient uptake of greengram [*Vigna radiata* (L.) Wilczek]. *Journal of Pharmacognosy and Phytochemistry*. 8(5), 2404-2408.
6. Boradkar, S. G., P. B. Adsul, M. S. Shelke, & Y. R. Khule. (2023). "Effect of iron and zinc application on soil properties, nutrient uptake and yield of green gram (*Vigna radiata* L.) in Inceptisol." *The Pharma Innovation Journal* 12(3): 1663-1669.
7. Doyle JJ. (1994) Phenology of the legume family: an approach to understanding the origin of nodulation. *Ann. Rev. Ecol. and Systematics*. 25: 325-349.

8. Gahlot, N., Singh, U., Moola, R., Mehriya, M. L., Borana, H. & Mandiwal, M. (2020). Biochemical assessment and yield of mungbean as influenced by zinc and iron fertilization. *Chemical Science Review and Letters*. 9(36), 949-95.
9. Gidaganti, A., Tarence, T., Smriti, R., & David, A.A. (2019). Effect of different levels of micronutrients on crop growth and yield parameters of green gram (*Vigna radiata* L.) Cv. IPM 02-03. *International Journal of Chemical Studies*. 7(3), 866-869.
10. Gowda, K.M., Halepyati, A.S., Koppalkar & Satyanarayarao (2014). Response of pigeonpea (*Cajanus cajan* L. Millsp.) to application of micronutrients through soil and foliar spray of macronutrients on yield, economics and protein content. *Karnataka Journal Agriculture Science*. 27(4), 460-463.
11. Kareti, S.S. & George, P.J. (2017). Effect of levels of phosphorus and zinc on growth and yield of kabuli chickpea (*Cicer kabulium* L.). *Journal of Pharmacognosy and Phytochemistry*. 6(4), 1013-1016.
12. Kumar, V., Yadav, R.B., Yadav, S., Verma, G. & Singh, M. (2020). Effect of micronutrients and different levels of macronutrients with rhizobium on yield and yield attributes of mungbean. *International Journal of Chemical Studies*. 8(6), 2428-2432.
13. Lokhande, P.B. (2018) *Effect of phosphorus and zinc on growth, yield, nutrient uptake and quality of green gram (Vigna radiata L.) in Inceptisol*. (Master's Thesis). Vasant Rao Naik Marathwada Krishi Vidyapeeth, Parbhani.
14. Meena, K.K., Meena, R.S., & Kumavat, S.M. (2013). Effect of sulphur and iron fertilization on yield attributes, yield and nutrient uptake of mungbean (*Vigna radiata*). *Indian Journal of Agricultural Sciences*. 83(4), 472-6.
15. Misal. B.D. (2018). *Response of macro and micro nutrient priming on growth, yield and quality of green gram (Vigna radiata L.)*. (Master's Thesis). Vasant Rao Naik Marathwada Krishi Vidyapeeth, Parbhani.
16. Panase, V. G. & Sukhatme, P. V. (1967). Statistical methods for agricultural workers. *Indian Council of Agricultural Research*, New Delhi.
17. Sudhanshu, S., Singh, R., Singh, E. & Dwivedi, K. (2022). Response of sulphur and iron fertilization on growth and yield of greengram (*Vigna radiata* L.). *The Pharma Innovation Journal*. 11(3), 259-263.