

Review Article

Transplanted v/s [no shortcut in title?](#) direct-seeded method of rice: A review

[Better to paraphrase title?](#)

Abstract

As rice traditionally grown as transplanted crop in puddled land leading to various problems because of labour shortage *i.e.*, unavailability of labours during the peak period of work [what is the relation between transplanted plant and labour shortage?? Clarify once](#). Other than that rising of nursery needs extra care and management. So there is a need for other methods of establishment of rice which produce higher or similar yields compare to transplanting. Direct seeding is one of the method which includes wet and dry seeding. Long term research has to be done in order to know the sustainability of direct seeded rice over transplanting.

Key words: Rice, semidry, drum seeded, transplanting, [one more keyword?](#)

Introduction

Rice (*Oryza sativa* L.) is one of the world's predominant staple food crops in about 40 countries and for more than 65 per cent of the population in India. Hence, rice can be called as 'The stuff of life'. World rice production demand was projected to increase by 25% from 2001 to 2025 to keep pace with population growth.

Traditional method of transplanting of rice is becoming difficult due to acute shortage of labour, especially during the peak periods of operation apart from higher labour wages. But rice growers across the country seek out elevated levels of productivity to counterbalance increasing costs of production. Also in transplanting method of establishment, the major operations such as nursery preparation, seedling rising, pulling from nursery, transporting and distribution of seedlings to main field and transplanting operation consumes 30-40% of total cost of cultivation (Rani and Jayakiran, 2003 [what is recent trends??](#)). Besides being laborious, this method of establishment also causes drudgery to womenfolk (Budhar and Tamilselvan, 2002). To reduce the labour requirement and cost on production, a need has been felt to replace the manual transplanting with scientific, economically feasible and environmentally safe establishment

method (Sanjay *et al.*, 2006). Hence an alternative to transplanting, direct seeding of rice can be practiced to overcome the above problems with an additional advantage of reduced crop duration and comparable grain yield (Sharma *et al.*, 2005) [already studies late before why not adopted till date discuss something on this also??](#).

1.1 Effect of crop establishment methods on growth of rice

[Make relation of one sentence with other?? In whole chapter??](#) Akkas Ali *et al.* (2012) reported that significantly taller plant height was recorded under transplanted rice (106.1 cm) compared to direct-seeded rice (98.5 cm).

Thirumurugan and Malarvizhi (2015) reported that puddled manual transplanting recorded significantly higher plant height (96 cm) over unpuddled machine planting (82.6 cm).

Bheruet *et al.* (2016) reported that, the higher plant height (71.09 cm) and dry matter (1371.92 g m^{-2}) was recorded in transplanted rice compared to direct seeded rice. The higher plant height (72.54 cm) and drymatter (945.90 g m^{-2}) at harvest was with drum-seeded rice over dry seeding (Sanjay *et al.*, 2018).

Prakhar *et al.* (2018) revealed that growth parameters *viz.*, plant height, number of tillers m^{-2} and dry matter production at harvest were on par between transplanting, direct seeded wet and dry rice. Among rice establishment methods, normal transplanting [what is normal and other transplanting?? Write just transplanting??](#) recorded significantly higher leaf area index at 90 DAS (4.44), plant height (100.64 cm), number of tiller (589.04 m^{-2}) and dry matter accumulation (999 g m^{-2}) at harvest than drum seeded and direct seeded rice (Rajnesh *et al.*, 2018). [Add](#)

[However or else??](#) Ravindra Nath and Jai Dev (2018) observed significantly higher plant height (82.42 cm) and leaf area index (3.05) in transplanting over drum seeded and direct seeded method of establishment.

1.2 Effect of crop establishment methods on yield and yield parameters of rice

Jamil and Hussain (2000) observed that transplanting of rice produced significantly more number of productive tillers (14.51 hill^{-1}), more number of spikelets ($124.53 \text{ panicle}^{-1}$) at harvest and paddy yield (2.77 q ha^{-1}) than direct sowing. The wet seeding by drum seeder was recorded

higher grain yield (56.57 q ha⁻¹) and it was on a par with the transplanting method of establishment (Budhar and Tamilselvan, 2002).

Ali *et al.* (2006) [recent references??](#) observed that among establishment methods, transplanting recorded higher number of filled spikelets (68 panicle⁻¹) and panicle length (25.8 cm) at harvest as compared to drum seeding and dry seeding, however, the grain yield was on par. The higher grain yield (5329 kg ha⁻¹) and straw (8247 kg ha⁻¹) yield recorded in transplanting method of rice establishment over drum seeded rice (Hugar *et al.*, 2009).

Jagtap *et al.* (2012) reported that among crop establishment methods, transplanting recorded significantly higher growth attributes, grain (52.58 q ha⁻¹) and straw yield (61.98 q ha⁻¹) over dibbling of seeds.

Sandhya *et al.* (2014) reported that among establishment methods, transplanting recorded significantly higher panicles (430 m⁻²), total grains panicle⁻¹ (135) and filled grains panicle⁻¹ (125) over drum seeded rice, but was on par with semi dry rice. However, grain and straw yields were on par among establishment methods.

Bheru *et al.* (2016) study revealed that significantly higher effective tillers (360.58 m⁻²), length of panicle (21.07 cm), weight of panicle (2.15 g), filled grains per panicle (97.08), test weight (22.24 g), grain (41.97 q ha⁻¹) and straw (47.42 q ha⁻¹) yield were in transplanting method of rice establishment as compared to drilling. [Relate with other sentences??](#)

Sanjay *et al.* (2018) recorded significantly higher effective tillers (274.89 m⁻²), grain (40.62 qha⁻¹) and straw (53.31 qha⁻¹) yield in drum seeded rice than dry seeded rice. The higher grain yield of 5.63 t ha⁻¹ was in transplanting over direct seeded rice (Ravi and Ramakichenin, 2018).

Prakhar *et al.* (2018) observed that panicles m⁻², number of grains panicle⁻¹, panicle length, panicle weight, grain weight panicle⁻¹, harvest index and 1000-grain weight were on par between transplanting, direct seeded wet and dry rice establishment methods. However the straw yield (52.48 q ha⁻¹) and grain yield (39.11 q ha⁻¹) were significantly higher in transplanting over direct dry seeding, but was on par with direct wet seeding. The higher grain yield of 3833 kg ha⁻¹ obtained in transplanting and lower with wet drum seeding after dry tillage (Vinay *et al.*, 2018).

Ravindra Nath and Jai Dev (2018) observed significantly higher grain panicle⁻¹ (112.81), length of panicle (23.23 cm), grain (51.45 qha⁻¹) and straw (64.24 qha⁻¹) yield in transplanting over drum seeded and direct seeded method of rice establishment. However, test weight was found non-significant.

Bhardwaj *et al.* (2018) observed significantly higher panicles (282.00 m⁻²), panicle length (23.46 cm), grains panicle⁻¹ (124.10), filled grain panicle⁻¹ (106.27), grain (44.18 qha⁻¹) and straw yield (68.43 qha⁻¹) in conventional transplanting followed by drum seeding of sprouted seeds. Among rice establishment methods, normal transplanting performed good by registering higher yield attributes and grain yield of 42.36 q ha⁻¹ (Rajnish *et al.*, 2018).

1.3 Effect of crop establishment methods on nutrient uptake and nutrient use efficiency of rice

Chander and Pandey (1997) noticed significantly higher N (112.8 kg ha⁻¹), P (17.0 kg ha⁻¹) and K (172.3 kg ha⁻¹) uptake under transplanting method of establishment than direct seeded rice.

Sandhya *et al.* (2014) reported that significantly higher N (87.8 kg ha⁻¹), P₂O₅ (35.1 kg ha⁻¹) and K₂O (143.7 kg ha⁻¹) uptake at harvest by rice were in transplanting than drum seeded rice, but was on par with semidry rice. The higher total N (82.80 kg ha⁻¹), P (16.05 kg ha⁻¹), K (102.21 kg ha⁻¹) uptake were observed in transplanted method of rice establishment than direct seeded rice (Rajnish *et al.*, 2018).

1.4 Economic feasibility of crop establishment methods in rice

Budhar and Tamilselvan (2002) revealed that rice registered the higher net income of Rs.19,039 and Rs 18,587 ha⁻¹ with a B:C ratio of 2.33 and 2.29 in manual broadcasting and drum seeded rice, respectively.

Hugar *et al.* (2009) recorded net returns of Rs. 35,903 and 30,387 ha⁻¹ in transplanting and drum seeded methods of establishments, respectively. The higher B:C ratio of 1.87 recorded in drum seeded method of establishment.

Sandhya *et al.* (2014) reported that, gross return of Rs.57,200,56,389 and 53,882 ha⁻¹ with a B:C ratio of 1.25, 1.65 and 1.45 in transplanting, semidry rice and drum seeder methods of establishment, respectively.

Bhardwaj *et al.* (2018) observed that net returns of Rs 45,704 and Rs 49,294 with the B:C ratio of 2.34 and 1.80 in drum seeded and transplanting method of rice establishment, respectively. The higher net return and B:C ratio (Rs 20,574 and 0.58, respectively) in transplanting method and B:C ratio (0.77) in direct seeded wet and dry rice method of establishment (Prakhar *et al.*, 2018).

Vinay *et al.* (2018) observed that higher B:C ratio of 1.70 in dry seeded rice with drum seeder and lower with wet drum seeding after dry tillage.

Conclusion

Though transplanting is the popular method of establishment it is cumbersome. nursery raising, transporting of seedlings to the main field and transplanting. Direct wet seeding and Direct dry seeding better options of crop raising as it saves considerable amount of time, labour and water requirement and there is a possibility of increasing the cropping intensity in case of direct wet sowing. Transplanting can be replaced by direct seeding under puddle condition.

References

- [Ali, A. M., make middle letter small?? In all names ??](#) ALI, A. M., LADHA, J. K., RICKMAN, J. AND LALES, J. S., 2006, Comparison of different methods of rice establishment and nitrogen management strategies for lowland rice. *J. Crop Improv.*, **16**(1): 173-189.
- ANBUMANI, S., CHANDRASEKHARAN, B. AND KUPPUSWAMY, G., 2004, Evaluation of establishment methods and NPK levels in rice and their impact on succeeding crops. *Agric. Sci. Digest*, **24**(3): 190-193.
- ANONYMOUS, 2008, Status paper on rice in Karnataka.
- BANERJEE, H., PAL, S. AND MAITI, S., 2006, Integrated effect of organic and inorganic fertilizer on the productivity and profitability of rice grown under rice-rice crop sequences in the Gangetic West Bengal. *J. Crop Weed*, **2**(1): 40-44.

- BHARDWAJ, R., SINGH, M. K., SINGH, C. S. AND SINGH, A. K., 2018, Efficacy of crop establishment methods on yield and economics of rice (*Oryza sativa* L.) under puddled condition. *Int. J. Curr. Microbiol. App. Sci.*, Special Issue-7: 3625-3631.
- BHERU, L. K., VIRESH GOVIND CHAVAN, V. A., RAJEMAHADIK, V. M., KANADE, R.V., DHOPAVKAR, H. K., AMETA AND TILEKAR, R. N., 2016, Effect of different rice establishment methods on growth, yield and different varieties during *kharif* season. *IJPJX-CAS-USA*, **6**(2): 127-131.
- BUDHAR, M. N. AND TAMILSELVAN, N., 2002, Effect of stand-establishment techniques on yield and economics of lowland irrigated rice (*Oryza sativa*). *Indian J. Agron.*, **47**(1): 57-60.
- CHANDER, S. AND PANDEY, J., 1997, Nutrient removal by scented basmati rice (*Oryza sativa*) and associated weeds as affected by nitrogen and herbicides under different rice cultures. *Indian J. Agron.*, **42**(2): 256-260.
- HUGAR, A. Y., CHANDRAPPA, H., JAYADEVA, H. M., SATHISH, A. AND MALLIKARJUN, G. B., 2009, Influence of different establishment methods on yield and economics of rice. *Agri. Sci. Digest.*, **29**(3): 202-205.
- JAGTAP, D. N., MAHADKAR, U. V., CHAVAN, L. S. AND SRINIVASARAO, C., 2012, Performance of rice (*Oryza sativa* L.) influenced by different crop establishment methods and fertilizer sources. *Indian J. Dryland. Agric. Res. Dev.*, **27**(2): 55-58.
- JAMIL, M. AND HUSSAIN, A., 2000, Effect of different planting methods and nitrogen levels on growth and yield of rice (BASMATI 385). *Pak. J. Agri. Sci.*, **37**(1-2): 83-87.
- PRAKHAR, K. S. K., CHOUDHARY AND ABHINANDAN, S., 2018, Effect of nitrogen scheduling on yield and yield attributing characters in rice under different establishment methods. *Int. J. Curr. Microbiol. App. Sci.*, Special Issue-7: 5153-5160.

- RAJNESH, T., SINGH, N. B., VIPUL, S. AND DEVESH, K., 2018, Effect of planting methods and integrated nutrient management on growth parameters, yield and economics of rice. *J. P. P.*, **7** (2): 520-527.
- RANI, T. S. AND JAYAKIRAN, K., 2003, Evaluation of different planting techniques for economic feasibility in rice. *Electronic J. Environ. Agric. Food Chem.*, **9**(1): 150-153.
- RAVI, S. AND RAMAKICHENIN, B., 2018, Evaluation of different nutrient management practices for enhancement of the productivity in different rice establishments methods. *Int. J. Curr. Microbiol. App. Sci.*, **7**(3): 1726-1731.
- RAVINDRA NATH AND JAI DEV., 2018, Effect of establishment methods under different weed management technique on rice and associated weeds. *J. Pharmacognosy Phytochemistry*, **7**(5): 807-810.
- SANDHYA, K. M., RAMANA, A. V. AND RAMANA MURTHY, K. V., 2014, Effect of different crop establishment techniques and nutrient doses on nutrient uptake and yield of rice (*Oryza sativa* L.). *Karnataka J. Agric. Sci.*, **27**(3): 293-295.
- SANDHYA, K. M., 2014, Appraisal of nutrient requirement of rice (*Oryza sativa* L.) under different crop establishment techniques. *M.Sc. (Agri.) Thesis*, Acharya N. G. Ranga Agricultural University.
- SANGEETHA, C. AND BASKAR, P., 2015, Influence of different crop establishment methods on productivity of rice—a review. *Agri. Review*, **36**(2): 113-124.
- SANJAY, K., ASHOK, K. S., SINGH, G., NARESHMANI, P., AJIT, K. AND DINESH K., 2018, Effect of establishment methods and weed management practices on the growth and yield of direct seeded rice (*Oryza sativa* L.). *J. Pharmacognosy Phytochemistry*, **7**(1): 164-168.

SANJAY, M. T., SETTY, T.K.P. AND NANJAPPA, H. V., 2006, Productivity, energetics and economics of different systems of crop establishment in rice. *Crop Res.*,**31**(3): 350-353.

SHARMA, S. K., PANDEY, D. K., GANGWAR, K.S. AND TOMAR, O.K., 2005, Effect of crop establishment methods on performance of rice (*Oryza sativa*) cultivars and their effect of succeeding wheat (*Triticum aestivum*). *Indian J. Agron.*,**50**(4): 253-255.

THIRUMURUGAN. S., AND MALARVIZHI. P., 2015, Establishment methods, crop residues and nitrogen effect on yield and economics of rice (*Oryza sativa* L) in Cauvery old delta. *Trends in Biosciences.*, **8**(10): 2537-2542.

VINAY, K., SHAILENDRA, S., VIDYASAGAR AND MAURYA, M. L., 2018, Evaluation of different crop establishment method of rice on growth, yield and economics of rice cultivation in agro-climatic condition of eastern Uttar Pradesh. *J. Pharmacognosy Phytochemistry*,**7**(3): 2295-2298.