

Mycological assessment of selected swimming pools of recreational centers around the University of Port Harcourt, Nigeria.

ABSTRACT

Aims: This study was carried out to determine the physicochemical properties and mycological safety of four swimming pools within and around the University of Port Harcourt, Rivers State, Nigeria.

Study design: Random sampling design was employed.

Place and Duration of Study: Department of Microbiology Laboratory, University of Port Harcourt, between August and October, 2023.

Methodology: A total of 8 samples were collected before and after patronage by patrons from 4 swimming pool. Standard methods were employed for the physicochemical parameters such as pH, turbidity and residual chlorine as well as the enumeration for fungi using potato dextrose agar.

Results: The pH values ranged from 4.50 to 10.00. The Nephelometric Turbidity Unit ranged from 0.45 to 0.90 NTU while the residual chlorine ranged from 0.36 to 355.00 mg/l. The fungal counts ranged from 1 to 9 cfu/ml, while the isolated species on the basis of cultural morphology and microscopic characteristics were identified as species of *Fusarium*, *Penicillium*, *Aspergillus*, *Histoplasma*, *Rhodotorula*, *Microsporium* and *Trichophyton*.

Conclusion: This study has revealed a potential health risk associated with the use of water of the examined recreational facilities due to none compliance with the WHO standard of < 1 cfu/100ml for fungal and the other parameters, suggesting a public health concern and need for effective interventions.

Keywords: Chlorine, molds, physicochemical properties, swimming pool, yeast

1. INTRODUCTION

Water is among the most important natural resources crucial for life processes in both plants and animals playing prominent roles in food production and processing, transportation, power generation, cooling industrial machines, irrigation, recreation amongst others [1, 2]. However, contaminated water, irrespective of the intended use poses serious threats to humans by serving as a medium for disease transmission [3].

A swimming pool is an enclosed body of water of limited size contained in a holding structure, patronized by different classes of people for leisure activities such as sports or rehabilitative treatment [4]. Pools can be built into the ground (in-ground pools) or built above ground; as a freestanding construction or as part of a building or other larger structures, aboard ocean-liners and cruise ships. Pools in most hotels or other guest houses are exposed to contamination from body fat and human waste materials such as nasal secretions, saliva, sweat, faecal, urine and body lotions and creams [5].

Public pools can infect individuals based on the fact that public pools do not use environmentally sound disinfectant agents in the water, rather brominating agents are used, when these brominating agent gets mixed with carbon- friendly substances such as urine, hair, skin, cosmetics and sunscreen it becomes toxic [6]. Reported injuries from swimming pools include asthma [7, 8], skin and eye irritation, neurological disease conditions [9, 10, 11] and other skin infections, especially fungal foot disease or athlete's foot and genital infections resulting from bacterial, fungal, protozoan and viral agents [4, 11, 12, 13, 14, 15, 16, 17].

The fungal species previously isolated from swimming pool included *Cephalosporium* spp., *Fusarium* spp., *Penicillium* spp., *Rhizopus* spp., *Aspergillus* spp., *Trichophyton mentagrophytes*, *Mucor* spp., *Candida albicans*, *Aspergillus niger*, *Alternaria* spp., *Absidia* spp. and *Trichophyton* spp.[18, 19, 20, 21].

The assessment of a variety of physical-chemical parameters at the time of collection, such as pH, turbidity, and free accessible chlorine, as well as the presence of fungi could also predict the quality of pool water which is the focus of this present study.

2. MATERIAL AND METHODS

2.1 Study area

The four swimming pools used for this study, were hotel swimming pools in communities around the University of Port Harcourt, Port Harcourt. The test samples were obtained directly from the swimming pools **before and after patronage by patrons.**

2.2 Determination of Physicochemical parameters

The physicochemical parameters analyzed were pH, turbidity and residual chlorine. These were all carried out according to the analytical methods described in Standard Methods for Water and Wastewater Analysis [22].

2.3 Enumeration of Total Fungi

Samples of the swimming pool water samples were serially diluted in ten folds. Then the molten potato dextrose agar at 45°C was supplemented with 1% lactic acid and poured into the Petri dishes containing 0.1 ml of the appropriate dilution for the isolation of the total fungi. They were swirled to mix and colony counts were taken after incubating the plates at room temperature (29±2 °C) for 48 to 72 h. Colonies from primary plates were purified on freshly prepared **potato dextrose** agar supplemented with 1% lactic acid for identification

2.4 Characterization and Identification of Fungal Isolates

Fungal isolates were examined macroscopically and microscopically using the needle mounts technique. Their identification was performed according to the scheme of Barnett and Hunter[23] and Larone [24].

3. RESULTS AND DISCUSSION

The presence or absence of pathogenic molds and yeast in recreational water has a crucial role in determining the health risk connected with swimming, as it can spread infectious diseases such as skin, eye, and ear infections [25].

3.1 Physicochemical parameters

Recreational water does not provide drinkable water to humans anywhere in the world. Its quality; however, needs to be on par with drinking water due to the high danger of microbial contamination from the environment and the fact that many swimmers inadvertently consume it while swimming. The results of the examined parameters are presented in Table 1. The turbidity, with the exception of a single pool water collected before patronage were within the WHO acceptable limit (0.5). The values obtained in this present study were within the range (0.22-12.61 NTU) reported by Ajadi et al. [21] in examined swimming pools in Osogbo metropolis, western Nigeria but are below the 38.00 to 90.00 NTU reported by Agomuo and Amadi [26] from swimming pools in Owerri, the Imo State capital, eastern Nigeria.

The pH of some pools was either below or above the WHO acceptable limit (7.2 -7.8). Some of the values obtained in this study were comparable to the 5.6 to 6.7, 6.8 to 7.1 and 4.48 to 7.70 reported by Agomuo and Amadi [26], Eze et al. [20] and Ajadi et al. [21], respectively. The low pH of 4.50 to 5.00 in two pools, will make the water more acidic; hence, patrons may experience burning eyes, itchy skins and easily ripped swimming clothes.

The residual chlorine in this present study (0.36-355.00) with the exception of a sample were higher than the WHO limit <3.0. Similar higher values of 221±6.0 to 294±10.0 have been reported by Eze et al. [20]. Elevated chlorine levels could lead to higher operational expenses and an increased risk of corrosion and scaling.

Table 1. Physicochemical parameters of swimming pool water

Parameters	USC Pool		MBL Pool		MH Pool		HH Pool		WHO Limit
	BS	AS	BS	AS	BS	AS	BS	AS	
Residual Chlorine (mg/l)	355.00	0.36	21.30	7.10	177.50	35.5	14.20	7.10	<3.0
Turbidity (NTU)	0.45	0.45	0.45	0.45	0.90	0.45	0.45	0.45	0.5
pH	10.00	6.68	5.00	5.00	7.10	7.10	4.50	4.50	7.2-7.8

BS=Before swimming; AS=After swimming

Osei-Adjei et al. [27] have posited that chlorine's ability to react with foreign particles diminishes as pH rises and that only about 20% of the chlorine applied to the pool can be utilized at pH 8.0 because the chlorine reacts with carbonates to generate scale. The water in the pool turns murky and aggravates the skin. Free (residual) chlorine levels should be maintained within limits set by national, local regulations or WHO recommendations as low free chlorine encourages the growth of microorganisms, including fungi in pool water, while high amounts can cause irritation to the skin, eyes, and upper respiratory tracts as well as the production of disinfection by-products such trihalomethanes [28].

Table 2. Identification of the fungal isolates

Codes	Cultural characteristics/appearance on PDA	Microscopic features using low power magnification and lactophenol cotton blue	Tentative organism
MH (BT) a	Brown sporing/granular surface and light cracked reverse	Vesicles are hemispherical and phialides are produced from a primary row of metulae. Phialides produce globose to elliptical conidia arranged in chains.	<i>Aspergillus</i> spp.
MH (BT) b	White woolly surface and light reverse	Hyphae are small and septate and give rise to phialides that produce single-celled microconidia.	<i>Fusarium</i> spp.
MH (BT) c	Black sporing/granular surface and light cracked reverse	Septate hyphae with long conidiophores that support spherical vesicles that give rise to large metulae from which long chains of conidia are produced.	<i>Aspergillus</i> spp.
MH (BT) d	Dull white, coarse woolly surface and light reverse	Smooth club-shaped, thin-walled macroconidia with 3-8 septa.	<i>Trichophyton</i> spp.
MH (BT) e	Dull white woolly surface and light reverse	Smooth club-shaped, thin-walled macroconidia with 3-8 septa.	<i>Trichophyton</i> spp.
MH (AT) a	Flat, white with a dense suede-like to downy surface	Large spindle shape-shaped multisegmented macroconidia with curved ends	<i>Microsporium</i> spp.
UPSC (AT)a	Black sporing/granular surface and light cracked reverse	Septate hyphae with long conidiophores that support spherical vesicles that give rise to large metulae from which long chains of conidia are produced.	<i>Aspergillus</i> spp.
UPSC (BT) a	Dense woolly white surface and light reverse	Large, rounded, single-celled, 8-14 μ m in diameter, tuberculate macroconidia formed on short, hyaline, undifferentiated conidiophores.	<i>Histoplasma</i> spp.
UPSC (BT) b	White woolly surface and light reverse	Hyphae are small and septate and give rise to phialides that produce single-celled microconidia.	<i>Fusarium</i> spp.
UPSC (BT) d	Dull white, coarse woolly surface and light reverse	Smooth club-shaped, thin-walled macroconidia with 3-8 septa.	<i>Trichophyton</i> spp.

MBLH (BT) a	Orange, 2mm, round, raised, smooth, shiny, white surface and light reverse	Oval shaped cells appearance single, budded, clustered.	<i>Rhodotorulaspp.</i>
MBLH (BT) b	Dull white, coarse woolly surface and light reverse	Smooth club-shaped, thin-walled macroconidia with 3-8 septa.	<i>Trichophyton spp.</i>
MBLH (AT) a	Blue velvet surface and light reverse	Hyphae are hyaline and septate and produce brush-like conidiophores.	<i>Penicillium spp.</i>
MBLH (AT) b	Dull white, cottony surface and light reverse	Smooth club-shaped, thin-walled macroconidia with 3-8 septa.	<i>Trichophyton spp.</i>
MBLH (AT) c	White fluffy surface and light reverse	Hyphae are small and septate and give rise to phialides that produce single-celled microconidia.	<i>Fusarium spp.</i>

UNDER PEER REVIEW

3.2 Mycological count and identity

The study revealed that the mycological quality of the examined swimming pool water (1 to 9 cfu/ml) generally exceeded WHO's limit (<1 cfu/100ml). The presumptive identities of the isolated molds and a yeast are presented in Table 2. The predominant fungal was *Trichophyton* spp. (33.33%), followed by *Aspergillus* spp. and *Fusarium* spp. (20.00% and the least occurring were jointly *Penicillium*, *Histoplasma*, *Microsporium* and *Rhodotorula* (6.67%). These molds and some yeasts have been reported in swimming pools by several researchers from different locations and countries [18, 19, 20, 21, 29]. Among these are *Aspergillus*, *Penicillium* and *Trichophyton* which have been implicated in human illness. According to Alice [30], *Aspergillus niger* is the causative agent of aspergillosis, which is typically an external ear infection (otomycosis) that can cause tympanic membrane perforation and ear canal ulceration. Additionally, *Aspergillus* has been documented to serve as a gateway for the spread of diseases in people with weakened immune systems [31]. *Trichophyton mentagrophyte* is the etiologic agent of human nails and feet infections. It is responsible for ringworm in the buttocks, groin, beard hair, and scalp. Prescott et al. [32] reported that *Fusarium* is known to cause eye infections in both humans and animals.

4. CONCLUSION

The results of this study revealed that the physicochemical parameters examined and fungi counts from the pools exceeded the WHO's acceptable limits thereby, underscoring the necessity of heightened monitoring of swimming pool water quality. Maintaining ideal pH, and chlorine levels, as well as using flocculants to keep water clear, are all important aspects of proper water management that are essential to stopping the spread of fungal illnesses. Regular inspection and testing of the swimming pools will guarantee all year-round, low-maintenance and safe swimming pool.

REFERENCES

1. Mackereth FH, Heron J, Talling JF. Water analysis. Ambleside: Fresh Water Biological Association Scientific Publication; 2003.
2. Vanloon GW, Duffy SJ. The hydrosphere. In: Environmental Chemistry: A Global Perspective 2nd Ed. New York: Oxford University Press; 2005.
3. Okoro HK, Adeyinka A, Jondiko OE, Ximba BJ, Kakalanga SJ. Assessment of heavy metals contamination in groundwater: A case study of central industrial district in Ilorin Kwara State, Nigeria. Int J Phys Sci 2012;7(28):5078-5088.
4. Amala SE, Aleru CP. (2016). Bacteriological quality of swimming pools water in Port Harcourt Metropolis. Nat Sci. 2016; 8: 79-84.
5. Saberianpour S, Momtaz H, Ghanbari F, Mahmodi F. Assessment of bacterial and fungal contamination in public swimming pools in Shahrekord, Iran. J. Trop. Dis. 2015; 4(2), 1-4.
6. Agbagwa OE, Young-Harry WM. Health implications of some public swimming pools located in Port Harcourt, Nigeria. Pub Health Res. 2012; 2 (6): 190-196.

7. Thickett KM, McCoach JS, Gerber JM, Sadhra S, Burge PS. Occupational asthma caused by chloramines in indoor swimming-pool air. *European Resp J*. 2002;19(5):827-832
8. Bernard A, Carbonnelle S, Michel O, Higuët S, De Burbure C, Buchet J, et al. Lung hyperpermeability and asthma prevalence in school children: Unexpected associations with the attendance at indoor chlorinated swimming pools. *Occup Environ Med*. 2003;60(6):385-394
9. Chiswell B, Wildsoet CF. The causes of eye irritation in swimming pools. *Water Sci Technol*. 1989;21(2):241-244
10. Erdinger L, Kirsch F, Sonntag HG. Irritierende Wirkung von Nebenprodukten der Schwimmbadwasserdesinfektion. *Zentralbl. Hyg. Umweltmed*. 1998; 200:491-503.
11. Schets FM, Engels GB, Evers EG. *Cryptosporidium* and *Giardia* in swimming pools in the Netherlands. *J. Water Health*. 2004; 2(3): 191-200.
12. Hilderbrand JM, Maguire HC, Holliman RE, Kangosu E. An outbreak of *Escherichia coli* O557: H7 infection linked to paddling pools. *Commun Dis, Rep. CDR Rev*. 1996; 6 (2): R33-R36.
13. Papapetropoulou M, Vantarakis AC. Detection of adenovirus outbreak at a municipal swimming pool by nested PCR amplification. *J. Infect*. 1998; 36: 101-103.
14. Centre for Disease Control (CDC). An outbreak of norovirus gastroenteritis at a swimming club-Vermont. *Morbidity and Mortality Weekly Report*. 2004.
14. Tate D, Mawer S, Newton A. Outbreak of *Pseudomonas aeruginosa* folliculitis associated with a swimming pool inflatable. *Epidemiol. Infect*. 2003; 130: 187-192.
15. Glauner T, Kunz F, Zwiener C, Frimmel FH. Elimination of swimming pool water disinfection by-products with advanced oxidation processes (AOPs). *Acta HydrochimHydrobiol*. 2005;33(6):585-594.
16. World Health Organization (WHO). Water, Sanitation, and Health Team. Guidelines for safe recreational water environments – Vol. 2 swimming pools and similar environments. Geneva: World Health Organization. 2006.
17. Barna Z, Kádár M. The risk of contracting infectious diseases in public swimming pools. A review, *Ann Ist Super Sanità*. 2012; 48(4): 374-386.
18. Papadopoulou C, Economou V, Sakkas H, Gousia P, Giannakopoulos X, Dontorou C, et al. Microbiological quality of indoor and outdoor swimming pools in Greece: Investigation of the antibiotic resistance of the bacterial isolates. *Int J Hyg Environ Health*. 2008; 211(3-4): 385-397.
19. Bello OO, Mabekoje OO, Egberongbe HO, Bello TK. Microbial qualities of swimming pools in Lagos, Nigeria. *Int J Appl Sci Technol*. 2012; 2(8): 89-96.
20. Eze VC, Onwuakor CE, Ikwuegbu AL. Microbiological and physicochemical characteristics of swimming pool water in Owerri, Imo State, Nigeria. *J Appl Environ Microbiol*. 2015; 3(1): 6-10.

21. Ajadi FA, Bakare MK, Oyedeji O. Assessment of the physicochemical and microbiological qualities of swimming pools in selected hotels in Osogbo metropolis, southwestern Nigeria. *Ife J Sci.*2016; 18 (4):831-843.
22. American Public Health Association (APHA). *Standard Methods for the Examination of Water and Wastewater* (23rd ed.). Washington DC: American Public Health Association. 2017.
23. Barnett HL, Hunter BB. *Illustrated genera of fungi, imperfecti*(3rd ed.) Burgess Publishing Company, Minnesota, U.S.A. 1972.
24. Larone BH. *Important Fungi: A Guide to identification*, Harper and Row Publishers, Hagerstown, Maryland.1986.
25. Lagerkvist BJ, Bernard A, Blomberg A, Bergstrom E, Forsberg B, Holmstrom K, et al. Pulmonary epithelial integrity in children: relationship to ambient ozone exposure and swimming pool attendance. *Environ Health Perspect.* 2004; 112(17): 1768-1771.
26. Agomuo EN, Amadi PU. Swimming pools quality risk assessment for heavy metal deposition and intake via oral and dermal exposure. *Environ Anal Health Toxicol.*2019; 34(3):1-9.
27. Osei-Adjei G, Sarpong SK, Laryea E, Tagoe E. Bacteriological quality assessment of swimming pools in the Osu-Labadi Area, Accra. *J Nat Sci Res.*2014; 4(19): 126-129.
28. Bilajac L, Lušić DV, Jelinic JD, Rukavina T. Microbiological and chemical indicators of water quality in indoor hotel swimming pools before and after training of swimming pool operators. *JWater Health.* 2012; 10(1): 108-115.
29. Okafor UC, Nworji KE, Obubu M. Antimicrobial susceptibility pattern of microorganisms from some selected swimming pools in Awka, Anambra State, Nigeria. *Int J Biomed Clin Sci.*2020; 5(3): 145-152.
30. Alice LS. *Principles of microbiology.* (5th ed.) CV Mosby, London. 1977.
31. Buot G, Toutous-Trellu L, Hennequin C. Swimming pool decks as environmental reservoir of *Fusarium*. *Med Mycol.* 2010; 48 (5): 780-784.
32. Prescott LM, Harley JP, Klein DA. *Microbiology.* (5th ed.) Report 71. The Bacteriological examination of water supplies. (4th ed.) London: HMSO, 1969.: McGraw-Hill Book, New York. 2002.