

EVALUATION AND COMPARISON OF THERAPEUTIC EFFICACY OF VARIOUS HORMONAL PROTOCOLS IN ANOESTRUS BUFFALO HEIFERS OF SEMI-ARID REGION

ABSTRACT

To assess and compare different hormonal protocols' treatment efficacy in anoestrus buffalo heifers in a semi-arid region. Forty buffalo heifers with a history of anoestrus were chosen randomly from the semi-arid region of Banaskanth and divided into four equal groups. Group-I: Heatsynch protocol (n = 10), Group-II Ovsynch protocol (n = 10), Group-III Doublesynch protocol (n = 10) and Group-IV Control (n = 10) are followed by artificial insemination at a predetermined period in treated buffalo heifers and insemination during spontaneous estrus in group-IV buffalo heifers. In Groups I, II, and III, the corresponding percentages of estrus induction were 90, 80, and 80 percent. Major symptoms such as edoema, erythema, dampness of vulva and estrus mucus secretion were present during the expression of estrus. Fifty percent of the buffalo heifers in Group IV displayed estrus at varying intervals. After three consecution services, Group II had the highest overall fertility rate (50%) followed by Group III (40%), Group I (30%), and Group IV (30%). In true anoestrus buffalo heifers, the Heatsynch, Ovsynch, and Doublesynch procedure can be applied with success to induce estrus. Buffalo heifers that follow the Ovsynch protocol may have a higher conception rate.

KEYWORDS: Heatsynch, Ovsynch, Doublesynch, buffalo heifers, anoestrus.

1 INTRODUCTION

Buffalo is crucial to the upkeep of a sustainable food production system in emerging nations and India is not an exception. 53% of the world's milk produced by buffaloes comes from India (Sethi *et al.*, 2014). Since buffaloes often exhibit both silent and anestrus estrus, buffalo farmers typically have difficulty identifying when their animals are in estrus. Moreover, their incapacity to exhibit clear symptoms of estrus, seasonality, delayed puberty onset and variable duration of estrus impede their improved reproductive control and genetic advancement. Prolonged anestrus and postpartum acyclicity also cause buffalo breeders to suffer significant financial losses.

Anestrus is one of the most commonly occurring reproductive problems in cattle and buffalo of India, affecting livestock productivity and economics to a great extent. It is a functional disorder of the reproductive cycle that is characterized by the absence of overt signs of estrus manifested either due to lack of expression of estrus or failure of its detection. In heifers, it poses a herd problem possibly due to low

plane of nutrition, the stress of seasonal transition or extremes of climatic conditions. The clinical survey revealed higher incidences of anoestrus and inactive ovaries in buffaloes (55.5 and 19.4%, respectively) than in cows (43 and 17.2%, respectively) (Tanwar *et al.*, 2003). Incidence in India has been reported between 2.13–67.11 and 9.09–82.50 percent in indigenous cattle and buffaloes (Thakor and Patel, 2013). Incidence in heifers has been reported between 12.37 to 64.66 percent (Sinha *et al.*, 1987). Variable duration of estrus (4-16 hours) and difficulty in prediction time of ovulation make AI application in buffaloes difficult (Baruselli, 2001). This consideration indicates the need for estrus synchronization using fixed-time insemination for implementation of breeding programs in buffaloes (Presicce *et al.*, 2004).

To improve reproductive efficiency, several protocols of estrus induction and ovulation synchronization have been developed. These procedures are based on manipulating the Corpus luteum, either to induce premature luteolysis using prostaglandins or to prolong the luteal phase using progestogens. Estrus synchronization has many advantages and is becoming mandatory in modern animal husbandry practices in indigenous cattle and buffaloes, which are known for anoestrus and silent estrus. Looking at the above fact the comparative study of various synchronization protocols in ovulation was decided. Hence, this study was planned to evaluate the comparative efficacy of heatsynch, ovsynch and doublesynch protocols for fertility enhancement in anestrus buffalo heifers.

2 MATERIAL AND METHODS

2.1 LOCATION OF THE EXPERIMENTAL AREA

The present study was carried out at Banaskantha district at Department of Veterinary Gynecology and Obstetrics, College of Veterinary Science and Animal Husbandry, Sardarkrushinagar, Dantiwada, SDAU with Collaboration of All India Co-ordinated Research Project (AICRP) on Nutritional and Physiological Intervention for enhancing reproduction performance in animal, Dept. of Animal Reproduction, Gynecology & Obstetrics, College of Veterinary Science and Animal Husbandry, AAU, Anand.

2.2 EXPERIMENTAL PROTOCOL

2.2.1 Selection of the Buffalo Heifer

A total 40 Anestrus buffalo heifers with an average body condition score between 3 to 5 with a history of anestrus were selected from villages of Banaskantha milk-shed areas of Gujarat. The buffalo heifers were screened gynaeco-clinically for their reproductive status. Detailed history and rectal palpation findings were recorded. The anestrus buffalo heifers having small, smooth, inactive ovaries with normal genitalia and no palpable CL on either ovary, relaxed uterine horn was selected for study. The reproduction status of their buffalo heifers was re-examined after 10 days to conformation to the condition.

2.2.2 Grouping of Buffalo Heifers

All the 40 buffalo heifers were randomly divided into four groups as follows.

2.2.2.1 Group-I (HeatSynch Protocol, n=10).

The Selected buffalo heifers of this group were administered with Intramuscular Injection (Inj.)- of 10 µg of GnRH analog i.e. Buserelin Acetate 10 µg (Receptal , @ 2.5 ml, Intervet India Pvt Ltd, New delhi) on day 0, followed by Inj. of 500 µg PGF_{2α} analogue i.e. Cloprostenol sodium (Estrumate, @ 2 ml, Intervet India Pvt Ltd) and estradiol benzoate 1 mg I/M (Sigma, USA) on days 7 and 8, respectively, and fixed time artificial inseminated(FTAI) twice at 48 and 60 hrs post-Estradiol injection.

2.2.2.2 Group-II (Ovsynch Protocol, n=10):

The selected buffalo heifers of this group were administered with Intramuscular Inj. of 10 µg of Buserelin acetate on day 0, Inj. of 500 µg Cloprostenol sodium, on day 7 and second Inj. of 10 µg of Buserelin acetate on day 9 followed by fix time artificial insemination twice at 12 and 24 hrs later.

2.2.2.3 Group-III (Double Synch Protocol, n=10):

The buffalo heifers of this group were administered with Inj. of 500 µg Cloprostenol sodium on -3 day (3 days before the first injection of GnRH), I/M followed Inj. of 10 µg Buserelin acetate on day 0, Inj. of 500 µg Cloprostenol sodium on 7th day and second Inj. of 10 µg Buserelin acetate on 9th day, the buffalo heifers were fix time artificial inseminate twice at 12 and 24 hrs later following the second injection of buserelin acetate.

2.2.2.4 Group-IV (Anoestrus control, n=10):

The selected buffalo heifers of this group were injected with normal saline 5 ml, i.m on days 0, 7 and 9 and were observed for onset of natural estrus and artificial insemination was performed on detection of estrus. The group served as an untreated control.

2.3 ESTRUS INDUCTION AND ESTRUS INTENSITY

The buffalo heifers were closely observed for the exhibition of estrus symptoms at the time of fixed time insemination. The intensity of induced estrus was recorded based on the expressions of estrus symptoms. Estrus symptoms of group IV were recorded at the spontaneous estrus. The estrus intensity of selected buffalo heifers were monitored based on their behavior changes i.e. bellowing, excitement, off-feed, estrus mucus discharge as well as edema, erythema and wetness of the vulva.

2.4 FERTILITY RESPONSE

The buffalo heifers which did not show signs of estrus following the treatment were examined rectally for pregnancy diagnosis on 60th-day post-insemination to confirm the pregnancy. The buffalo heifers that did not conceive at the fixed time insemination were re-inseminated up to two subsequent estrus cycles. The conception rate up to 3rd insemination was recorded.

2.5 STATISTICAL ANALYSIS OF DATA

The data collected were suitably tabulated and analyzed following standard statistical method shown by Steel and Torrie (1981).

3 RESULTS AND DISCUSSION

3.1 INDUCTION AND INTENSITY OF ESTRUS

When the records of all 40 buffalo heifers were pooled it was observed that 75 percent (30/40 buffalo heifer) heifers showed induction of estrus. The induction of estrus along with the intensity in respect to the expression of estrus signs like bellowing, excitement, off feed, estrus mucus discharge and edema, erythema and wetness of vulva were recorded. The details of observation (Induction of estrus (%)) are depicted in figure 1.




Fig. 1: Induction of estrus (%) in the buffalo heifers treated with different treatment and control group.

The recorded estrus intensity at the time of artificial Insemination in Group-I (HeatSynch Protocol) was 90 percent (9/10) induction of estrus with excitement (55.55%), estrus mucus discharge (EMD) (100%), edema, erythema and wetness of vulva (88.88%). The findings (bellowing, excitement, mucus discharge and swollen) of the present study are similar of Mohan *et al.* (2009) in buffaloes following Heatsynch protocol.

Group-II (Ovsynch Protocol) expresses 80percent (8/10) induction of estrus with excitement (87.50 %), bellowing (11.1 %), estrus mucus discharge (100%) and edema, erythema and wetness of vulva (100%). Atanasov *et al.* (2011) observed mucus discharge in buffaloes at the time of AI following Ovsynch protocol. Similarly Malik *et al.* (2011) and Yotov *et al.* (2012) also observed 40 and 66.7 percent anestrus buffaloes showing estrus mucus discharge following Ovsynch protocol.

In Group-III (Double Synch Protocol) estrus induction was 80percent (8/10) with excitement (75%), estrus mucus discharge (100%) and edema, erythema and wetness of the vulva (100%). Hoque *et al.* (2014) observed diverse intensities of estrus signs in Ovsynch-treated water buffaloes. They observed Bellowing, Frequent urination, swelling of the vulva and pink-colored vestibule along with uterine tone at time of insemination in all the buffaloes at induce estrus with different protocols like modified Ovsynch and Double-ovsynch.

Whereas in group IV (control anestrous buffalo heifer) 50 percent (5/10) heifer showed estrus at different time intervals. i.e 20 days (2 buffalo heifer), 35 days (1 buffalo heifer) and 55 days (2 buffalo heifer) post last injection of NS(Normal saline) with expressions of excitement (100%), bellowing (60%), estrus mucus discharge (100%) and edema, erythema and wetness of vulva (100%).

In Heatsynch protocol total of 90 percent (9/10) of buffalo heifers showed estrus symptoms at the time of insemination. A similar result was recorded by Buhecha *et al.* (2016) in anestrued buffaloes. However, 100 percent estrus was recorded following heatsynch protocol in buffalo by Ali *et al.* (2012) and Mohan *et al.* (2014). Kandiel *et al.* (2012) also reported 100 percent estrus induction by Heatsynch protocol in buffalo heifers.

In Ovsynch protocol total of 80 percent (8/10) buffalo heifers showed estrus induction at the time of insemination. A similar result was recorded by Jabeen *et al.* (2012), Savalia *et al.* (2014) and Buhecha *et al.* (2016) in anoestrus buffaloes. Lower estrus response was observed by Kandiel *et al.* (2012), Kalwar *et al.* (2015) and Waqas *et al.* (2016) in buffaloes. However, 89 percent estrus was recorded following Ovsynch protocol in buffalo by Neglia *et al.* (2003) and Paters *et al.* (2003). The 100 percent estrus induction was recorded following Ovsynch protocol in buffalo by Atanasov *et al.* (2011) and Nakrani *et al.* (2014). Paul and Prakash, (2005) reported 90 percent estrus synchronization in buffalo following Ovsynch protocol.

In Doublesynch protocol total of 80 percent (8/10) of buffalo heifer showed estrus symptoms at the time of insemination. Similar results were recorded by Ozturk *et al.* (2010) in dairy cows. However, 100 percent estrus was recorded following doublesynch protocol in buffalo by Mirmahmoudi and Prakash, (2012).

In the control group total of 50 percent (5/10) buffalo heifer showed estrus symptoms at different time intervals within 90 days. Lower estrus response was observed by Parmar *et al.* (2012), Nakrani *et al.* (2014) and Buhecha *et al.* (2016) in anestrous buffaloes.

In this study maximum estrus induction was obtained in the Heatsynch group followed by Ovsynch and Doublesynch group. The higher success rate in Buffalo heifer observed in Heatsynch group using estradiol benzoate could be due to the higher sensitivity of buffalo heifer to this treatment.

3.2 FERTILITY RESPONSE

The buffalo heifers of the treatment group were inseminated at fixed time and the control group was inseminated at spontaneous estrus by the semen of a buffalo bull. The buffalo heifers that failed to conceive at 1st insemination were re-insemination for two subsequent estrus. The fertility response of buffalo heifer is shown in table 1 and figure 2.

Table 1: Fertility response of buffalo heifers following different treatments.

Group	Conception/service			Overall l	Percent t (%)
	1 st	2 nd	3 rd		
Group I (n=10)	1 (10.00%)	2 (22.22%)	-	3/10	30.00
Group II (n=10)	2 (20.00%)	2 (25.00%)	1 (16.66%)	5/10	50.00
Group III (n=10)	2 (20.00%)	1 (12.50%)	1 (14.28%)	4/10	40.00
Group IV (n=10)	1 (10.00%)	2 (22.22%)	-	3/10	30.00

The recorded overall Fertility response of group-I, II, III and IV was 30.00, 50.00, 40.00 and 30.00 percent, respectively. The 1st insemination conception rate of group I, II, III&IV was 10.00, 20.00, 20.00 and 10.00 percent, respectively. Whereas the similar fingers for 2nd insemination conception rate were 22.22, 25.00, 12.50 and 22.22 percent, respectively. Following the third insemination recorded conception rate was 0, 16.66, 14.28 and 0 percent, respectively in group I, II, III and IV.




Fig. 2: Fertility rate (%) in buffalo heifers treated with hormonal protocol and control group.

In this study 30 percent Fertility response was achieved by using Heatsynch protocol in anestrous buffalo heifers. A similar conception rate was obtained by Mohan *et al.* (2009) using Heatsynch protocol during the summer and winter seasons in anestrous buffaloes. An almost similar conception rate was obtained by Buhecha *et al.* (2016) in buffaloes using Heatsynch protocol.

Fertility response obtained was 50 percent following Ovsynch protocol in anestrous buffalo heifers. Similar results were obtained by

Berber *et al.* (2002) and Naikoo *et al.* (2010) in buffalo. However, contrary to the findings lower conception rate was recorded in post-partum anestrus buffaloes (Atanasov *et al.*, 2011 and Buhecha *et al.*, 2016). The higher conception rate following Ovsynch protocol was recorded by Kandiel *et al.* (2012), Nakrani *et al.* (2014) and Savalia *et al.* (2014) in anestrus buffaloes.

In Doublesynch protocol obtained fertility response was 40 percent in anestrus buffalo heifers. A higher conception rate was obtained by Mirmahmoudi and Prakash, (2012) in anestrus buffaloes. Ozturk *et al.* (2010) reported 72 percent conception rate in anestrus dairy cow following Doublesynch protocol.

In control group overall fertility response was 30 percent in anestrus buffalo heifers. Contrary to the present study higher conception rate (Nakrani *et al.*, 2014 and Buhecha *et al.*, 2016) in anestrus buffaloes and a lower conception rate (10.00%) in buffaloes (Yotov *et al.*, 2012) was observed.

CONCLUSIONS

According to the study's findings, true anoestrus buffalo heifers can be successfully induced to undergo estrus using the Heatsynch, Ovsynch, and Doublesynch methodology. The most noticeable symptoms of induced estrus in buffalo heifers were erythema, edema, the wetness of vulva, and estrus mucus secretion. In buffalo heifers, comparatively improved fertility response can be obtained by using the Ovsynch protocol followed by Doublesynch, and Heatsynch protocols.

REFERENCES

- Ali, R.; Shukla, S.P. and Nema, S.P. (2012). Hormonal induction of ovarian cyclicity and conception rate in postpartum anoestrus buffaloes. *Indian Journal of Field veterinarian*. **7**(4): 44-46.
- Atanasov, A.; Yotov, S.; Antonov, A. and Kolev, P. (2011). Induction of oestrus and conception rates in Bulgarian Murrah buffaloes after fixed-time artificial insemination (A Preliminary Study). *Bulgarian Journal of Veterinary Medicine*. **14**(3): 165–170.
- Baruselli, P. S. (2001). Control of follicular development applied to reproduction biotechnologies in buffalo, p. 128-146. In Proceedings of the “I: Congresso Nazionale Sull’Allevamento Del Bufalo”, Eboli, Italy.
- Berber, R. C. A.; E. H. Madureira, and P. S. Baruselli. (2002). Comparison of two ovsynch protocols (GnRH vs. LH) for fixed timed insemination in buffalo (*Bubalus bubalis*). *Theriogenology*. **57**:1421-1430.
- Buhecha, K. V.; Dhama, A. J.; Theodore, V. K.; Thakor, R. and Parmar, S. C. (2016). Effect of Various Ovulation Synchronization Protocols on Estrus Response,

Conception Rate and Blood Biochemical Profile in Anoestrus Buffaloes. *International Journal of Advanced Veterinary Science and Technology*. **5** (1): 232-238.

- Hoque, M. N.; Talukder, A. K.; Akter, M. and Shamsuddin, M. (2014). Evaluation of ovsynch protocols for time artificial insemination in water buffaloes in Bangladesh. *Turkish Journal of Veterinary and Animal Science*. **38**:418-424.
- Jabeen, S.; Anwar, M.; Andrabi, S. M. H.; Mehmood, A.; Murtaza, S. and Shahab, M. (2012). Determination of Ovsynch Efficiency for Oestrus Synchronization by Plasma LH and P4 Levels in Nili Ravi Buffalo during Peak and Low Breeding Seasons. *Pakistan Veterinary Journal*. **33** (2): 221-224.
- Kalwar, Q.; Memon¹, A. A.; Bhutto, M. B.; Kunbhar H. K.; Mirani, A. H.; Anwar, M. and Wagan, S. A. (2015). Estrus response and fertility rate in Kundhi buffaloes following estrus synchronization in breeding season. *Journal of advanced Veterinary & animal research*. **2**(3): 362-365.
- Kandiel, M. M. M.; Basssuoni, L. A.; Sosa, G. A.; Abou Ahmed, M. M. and Ghallab, A. M. (2012). Comparative Efficacy of Ovsynch and Heatsynch protocols Assessed by transrectal Ultrasonography and serum progesterone in Egyptian buffalo heifers. *Theriogenology*. **2**(3):178-183.
- Malik, R. K.; Singh, P.; Singh, L. J.; Sharma, R. K.; Phulia, S. K.; Tuli and Chandolia, R. K. (2011). Ovarian response and fertility of ovsynch-treated post-partum anestrous murrah buffaloes. *Buffalo Bulletin*. **30**(4): 272-276.
- Mirmahmoudi, R. and Prakash, B.S. (2012). The endocrine changes, the timing of ovulation and the efficacy of the Doublesynch protocol in the Murrah buffalo (*Bubalus bubalis*). *General and Comparative Endocrinology*. **177**: 153–159
- Mohan, K.; Nitu and Prakash, B. S. (2014). Evaluation of the Heatsynch protocol in murrah buffaloes (*Bubalus bubalis*) during hot summer season. *Journal of animal research*. **4**(2):241-246.
- Mohan, K.; Sarkar, M. and Prakash, B. S. (2009). Efficiency of Heatsynch Protocol in Estrous Synchronization, Ovulation and Conception of Dairy Buffaloes (*Bubalus bubalis*). *Asian-Australian Journal of Animal Science*. **22**(6): 774-780.
- Naikoo, M.; Patel, D.M.; Sarvaiya, N.P. and Killader, A. (2010). Estrous synchronization in postpartum anestrous mehsana buffaloes using different hormone protocols. *Indian Journal of Field Veterinarian*. **6**(2): 1-4.
- Nakarani, B. B.; Panchal, M. T.; Dharni, A. J.; Hadiya, K. K.; Patel, J. A. and Gosai, R. K. (2014). Effect of control breeding techniques on fertility and plasma profile of biochemical and mineral constituent in anestrous buffaloes. **1**(4B); 299-304.

- Neglia, G.; Gasparini, B.; Di Palo, R.; De Rosa, C.; Zicarelli, L. and Campanile, G. (2003). Comparison of pregnancy rates with two estrus synchronization protocols in Italian Mediterranean Buffalo cows. *Theriogenology*. **60**:125-133.
- Ozturk, O. A.; Cirit, U.; Baran, A. and Ak, K. (2010). Is Doublesynch protocol a new alternative for timed artificial insemination in anestrus dairy cows. *Theriogenology*. **73**: 568-576.
- Parmar, K. H.; Shah, R. G.; Tank, P. H. and Dhama, A. J. (2012). Effect of hormonal and non-hormonal treatment on reproductive efficiency and plasma progesterone, bio-chemical and macro-minerals profile in postpartum anoestrus buffaloes. *Indian Journal of Field Veterinarian*. **8**(2): 48-54.
- Paul, V and Prakash, B. S. (2005). Efficacy of the Ovsynch protocol for synchronization of ovulation and fixed-time artificial insemination in Murrah buffaloes (*Bubalus bubalis*). *Theriogenology*. **64**: 1049-1060.
- Peters, M. W. and Pursley, J. R. (2003). Timing of final GnRH of Ovsynch protocol affects ovulatory follicle size, subsequent luteal function and fertility in dairy cows. *Theriogenology*. **60**:1197-1204.
- Presicce, G. A.; Senatore, E. M.; Bella, A.; De Santis, G.; Barile, V. L.; De Mauro, G. J.; Terzano, G. M.; Stecco, R. and Parmeggiani, A.(2004). Ovarian follicular dynamics and hormonal profiles in heifers and mixed-parity Mediterranean Italian buffaloes (*Bubalus bubalis*) following an estrus synchronization protocol. *Theriogenology*. **61**: 1343–1355.
- Savalia, K. K.; Dhama, A. J.; Hadiya, K. K.; Patel, K. R. and Sarvaiya, N.P. (2014). Influence of controlled breeding techniques on fertility and plasma progesterone, protein and cholesterol profile in true anoestrus and repeat breeding buffaloes. *Veterinary World*. **7** (9): 727-732
- Sethi, R. K.; Singh, K. P.; Balhara, A. K. and Sunesh. (2014). Home. Available at <http://www.buffalopedia.cirb.res.in/>
- Sinha, B. P.; Sinha, S. N. and Singh, B. (1987). Incidence of anestrus in crossbred cattle in field and farm conditions. *Livestock Advisor*. **12**(32):43-48.
- Steel, R. G. D. and Torrie, J. H. (1981). Principles and Procedures of Statistics, A Biometric Approach. 2nd Edn. Mc Graw Hill, Int. Book Agency, Singapore.
- Tanwar, P. S.; Rakha, N. K. and Phogat, J. B. (2003). Challenges in buffalo infertility. *Intas polivet*. **4** (2): 121–27.
- Thakor, D. and Patel, D. (2013). Incidence of infertility problems in cattle and buffaloes . dairy cattle. <http://en.engormix.com>.
- Waqas, M.; Mehmood, M. U.; Shahzad, Q.; Kausar, R.; Sattar, A. and Naseer, Z. (2016). Comparative efficacy of G6G and Ovsynch protocols on synchronization and pregnancy rate in Nili–Ravi buffalo. *Animal reproduction science*. **166**: 9-14.

Yotov, S.; Atanasov, A. and Ilieva, Y. (2012). Therapy of ovarian inactivity in postpartum Bulgarian Murrah buffaloes by PRID and Ovsynch estrus synchronization protocols. Asian Pacific Journal of Reproduction.1(4): 293-299.

UNDER PEER REVIEW