

## ***Trichoderma* spp. isolates from agriculture field identified internal transcribed spacers (ITS), with chitinase activity and biocontrol activity against *Fusarium* spp.**

### **Abstract**

Twelve *Trichoderma* isolates were obtained from various Uttar Pradesh sites. Internal transcribed spacer (ITS) amplification and sequencing were used to identify isolates of *Trichoderma* that showed 99–100% identification with *Trichoderma harzianum*, *Trichoderma asperellum*, and *Trichoderma longibrachiatum*, the three species of *Trichoderma*. Tests were conducted in vitro to evaluate the biocontrol potential of *Trichoderma* isolates against *Fusarium* species. After seven days of incubation, the isolate TBT6 (*T. harzianum*) exhibited the greatest antagonism against *Fusarium oxysporum*, with inhibition rates of 87.1% by TBT7 (82.2%), and the least amount of inhibition (59.7%) by *T. longibrachiatum* isolate TBT10. Under field conditions, the antagonistic *T. harzianum* isolate TBT6 can be employed as a bio-control agent against *Fusarium* spp. and for the development of formulations based on *Trichoderma*. Using the dual plate approach, twelve highly antagonistic *Trichoderma* isolates were selected for chitinolytic activity. Observing the breakdown of chitin substrates, *Trichoderma* isolates TBT6 was shown to have strong chitinolytic enzyme activity, making it a good candidate for endochitinase gene isolation. Using two particular primers, the genomic DNA of the *Trichoderma* isolate TBT6 was isolated and amplified.

**Key words:** *Trichoderma*, internal transcribed spacers (ITS), antagonism, *Fusarium oxysporum*, Chitinase activity, Biocontrol.

### **INTRODUCTION**

Farmers use chemical pesticides as their first option to manage plant diseases and maximise crop yields. (Junaid *et al.*, 2013). It was estimated that 12% of crop loss is due to plant pathogens (Sharma *et al.*, 2012). Long-term pesticide use can lead to serious health and environmental issues in addition to being very expensive for developing nations. The famous saprophytic fungus *Trichoderma* may be isolated from any soil and, because of its high rate of colonisation and reproduction, is an effective biocontrol agent. (Pandya *et al.*, 2011; Pal and Gardener, 2006; El-Hassan *et al.*, 2013; Rao *et al.*, 2015). Numerous fungal phytopathogens, including *Phythium*, *Phytophthora*, *Macrophomina*, *Aspergillus*, *Rhizoctonia*, and *Fusarium*, are combated by *Trichoderma*. Employing a variety of biocontrol techniques, such as mycoparasitism, antibiotic synthesis, and pathogen competition for resources (food and space). (Benítez *et al.*, 2004; Rayatpanah *et al.*, 2011; Krishnamurthy and Shashikala, 2006). Recently, the global attention has been paid to *Trichoderma* as a safe alternative of pesticides and this led to an increase in the number of *Trichoderma* biocontrol products (Woo *et al.*, 2014). In agriculture, 37% of crop loss is due to pests, out of which 12% is due to pathogens. Fungi are responsible for more than 70% of all major crop diseases. Significant loss has been observed in several crop species like rice, wheat, barley, cotton, and groundnut due to fungi. In agriculture, annual crop losses due to pre and post-harvest fungal diseases exceeds 200 billion euros, in the United States alone, over \$600 million are annually spent on fungicides. (Sharma *et al.*, 2012).

*Fusarium oxysporum* species are ubiquitous soil-borne pathogens of a wide range of horticultural and food crops which cause destructive vascular wilts, rots, and damping-off diseases and several strains of pathogen to humans or animals (Gordon, 2017).

Chemical pesticides are used by the farmers for the management of fungal and pest control. The chemical pesticides which used have been affecting both humans and the environment. Chitinase is one of the enzymes that has been effectively used for the management of fungal diseases. It targets chitin of the cell wall for the management of plant pathogens. Plants have been known to implicate chitinase in defense against plant pathogens. Chitinase provides an alternate solution over harmful chemicals to plant defense against fungal pathogens. *Fusarium oxysporum* and *Fusarium graminearum* are the representative species known as plant-pathogenic *Fusarium* (Arie, 2019). The majority of *Fusarium* species are soil-inhabiting fungi. *Fusarium* conidia can be dispersed by water in rain splash and via irrigation systems but become airborne when dried, which makes them well-suited for atmospheric dispersal over long distances and which contributes to their worldwide distribution (Lin *et al.*, 2013). Although *Fusarium* utilizes multiple infection strategies, these fungi are considered to be hemibiotrophs capable of transitioning to necrotrophs depending on specific environmental and metabolic (Perfect and Green, 2001). As plant pathogens, they

cause root and stem rot, vascular wilt, or fruit rot in ~~a number of several~~ economically crop species resulting in major yield losses (MT ha<sup>-1</sup>) and in economic losses that value over \$1 Billion (Xia *et al.*, 2020). *Fusarium* toxins are the most abundant natural contaminants of diets containing cereals and other grains (Kumar *et al.*, 2013). Additionally, in clinical, several species are considered to be opportunistic pathogens in immune-compromised humans (Guarro, 2013).

Plant diseases management is a significant cost component in crop production. Traditionally, the approaches to dealing with disease in agricultural ecosystems include breeding resistant varieties of the crops species, hygiene to prevent the spread of contaminated soil or seed, and fungicides to kill potentially infecting fungi. However, increasing concerns about the effects of fungicides in the environment and residues in food have resulted in ~~the~~ deregistration of ~~a number of several~~ fungicides. Moreover, resistance of pathogens to fungicides has rendered certain fungicides ineffective. There is a need to ~~streng~~~~strengthen~~ ~~the~~ practices and components of IPM ~~in order~~ to reduce the dependence on synthetic agrochemicals. Biological control is an essential part of these strategies as a substitute ~~of~~ ~~for~~ agrochemicals. Biological control is the suppression of disease by the application of a Biocontrol Agent (BCA) usually a fungus, bacterium, or virus, or a mixture of these to the plant or the soil. The biological control agent acts to prevent infection of the pathogen by the plant. The main advantage of using a biological control agent they are highly specific for a pathogen and hence are considered harmless to non-target species.

The Biocontrol agents exercise several antagonistic mechanisms such as nutrient competition, antibiotic production, mycoparasitism and induction of systemic resistance has been proposed as the major antagonistic mechanism by *Trichoderma* produces enzymes like chitinase and  $\beta$ -1, 3-glucanase which degrades the cell wall materials and also releases ~~a number of several~~ toxic substances that can inhibit the growth of the pathogens fungal. (Puyam, 2016). These chitinase genes have functions in the biocontrol mechanism such as cell wall degradation, hyphal growth, and parasitic activity. (Lidia *et al.*, 2014). Chitinase helps in ~~the~~ breakdown of the glycoside bonds. Glucose oxidase ~~eatalyse~~~~catalyzes~~ D-glucose to D-glucono-1, 5-lactone, and hydrogen peroxide are known to have antifungal effects. Xylanase helps in breaking hemicelluloses a major component of plant cell walls. ~~The Chitinase-chitinase~~ gene (*ech42*) ~~was~~ produced by the biocontrol agent *Trichoderma harzianum* which was responsible for mycoparasitism. These genes have their unique functions in the biocontrol mechanism such as cell wall degradation, hyphal growth, stress tolerance, and parasitic activity. ~~Role-The role~~ of genes play a major the biocontrol process by regulating some signals and ~~lead~~~~ing~~ to the secretion of some enzymes that help in the degradation of the pathogens ~~and~~ hence they are known as biocontrol genes. Increased expression of the genes helps in enhanced biocontrol activity which helps in promoting ~~the~~ plant growth and prevents the plant from pathogen attack. The gene *ech42* from *T. harzianum* codes for endochitinase with significantly higher inhibitory activity against a broad range of phytopathogenic fungi than other chitinolytic enzymes. The *ech42* gene involved in the biocontrol activity of *Fusarium* disease incidence has not been reported. Thus the genetic characterization of *ech42* ~~is still~~ remains an open field for ~~the~~ researchers.

It is also known to have the ability to interact with plants, inducing resistance to biotic and abiotic stresses and promoting plant growth. These characteristics of fungi are useful in ~~the~~ research of *Trichoderma* strains as biocontrol agents. *Trichoderma harzianum* ~~being is~~ the most cited species as an active agents of commercial biopesticides and biofertilizers. Chitinases are found in a wide range of organisms including bacteria, fungi, higher plants, insects, and some vertebrates. Chitinases have been isolated from the *Trichoderma* species (Ganiger *et al.*, 2009). So the *Trichoderma* chitinase encoding gene/s will help in crop improvement and disease management. They will also minimize the use of chemical pesticides.

It was reported that ~~the~~ identification of *Trichoderma* based on morphological characters can give misleading results (Fahmi *et al.*, 2016). Recently, molecular identification based on internal transcribed spacers (ITS) amplification and sequencing ~~is has been~~ common and highly trusted (Savitha and Sriam, 2015; Fahmi *et al.*, 2016; Oskiera *et al.*, 2015; Jiang *et al.*, 2016). In this study, 10 isolates of *Trichoderma* spp. were isolated from soil rhizosphere of different locations in Egypt, characterized on ~~a~~ molecular level and screened for their antagonistic *Fusarium* oxysporum.

**Table 1. Isolation and identification of *Trichoderma* isolates based on ITS data**

---

Isolate code	State	GenBank Acc. No. (ITS sequence)	BLAST results identity	results identity (%)
<b>Molecular</b>				

## identification

---

TBT1	U.PMW776752	99 <i>T. asperellum</i>
TBT2	U.PMW776753	100 <i>T. asperellum</i>
TBT3	U.P.MW776754	99 <i>T. asperellum</i>
TBT4	U.PMW776755	99 <i>T. asperellum</i>
TBT5	U.PMW776756	100 <i>T. longibrachiatum</i>
TBT6	U.PMW776756	99 <i>T. harzianum</i>
TBT7	U.PMW776758	99 <i>T. harzianum</i>
TBT8	U.PMW776759	99 <i>T. harzianum</i>
TBT9	U.PMW776760	99 <i>T. longibrachiatum</i>
TBT10	U.PMW776761	99 <i>T. longibrachiatum</i>
TBT11	U.P-	- <i>Trichoderma spp.</i>
TBT12	U.P-	- <i>Trichoderma spp.</i>

## MATERIALS AND METHODS

### Isolation of *Trichoderma* isolates

twelve 12 isolates of *Trichoderma* spp. were recovered from the soil rhizosphere of various locations in Uttar Pradesh. ~~that~~ That were grown with various crops. ~~A number of~~ Several soil samples were taken at a depth of 15 cm, sealed in sterile bags, and brought into the lab for the isolation procedure. Fahmi et al. (2016) 1g air-dried soil sample ~~was~~ were added into 9 ml sterile water in a test tube to make a 1:10 dilution ( $10^{-1}$ ). The mixture was vigorously shaken on a vortex mixture for 5-10 minutes to obtain a uniform suspension. One ml of soil suspension was transferred into a fresh sterile test tube containing 9 ml sterile water under aseptic conditions to make  $10^{-2}$  dilution. Further,  $10^{-3}$  dilution was made by pipetting 1 ml suspension into another fresh test tube containing 9 ml sterile water. One ml of soil suspension was taken and uniformly spread on PDA and TSM-containing Petri plates and incubated at  $25 \pm 2^\circ\text{C}$  for 5-7 days in the dark. After incubation as the mycelial growth ~~were~~ appeared, the hyphal tips from the advancing mycelium were cut and transferred into the fresh PDA medium for further purification. The purified isolates were cultured on PDA slants and maintained at  $4^\circ\text{C}$  for further use.

### Mycelial growth kinetics of different *Trichoderma* isolates

Growth kinetics of different *Trichoderma* isolates performed on PDA medium. Recorded data of growth kinetics observed by ~~studied~~ studying their regular time ~~interval~~ interval for 12 h. By studying growth *Trichoderma* isolate viz., TBT1, TBT2, TBT3, TBT4, (*Trichoderma asperellum*), TBT5, TBT9, TBT10 (*T. longibrachiatum*), TBT6, TBT7, TBT8, (*T. harzianum*) TBT11 (*T. koningii*) and TBT12, (*T.*

*koningiopsis*). Among different *Trichoderma* TBT6, TBT7 and TBT9 groups were fast-growing were shown in (Table. 2)

**Table 2- Growth kinetics data mean value of different *Trichoderma* isolates (cm)**

Isolates	12hr	24hr	36hr	48hr	60hr	72hr	84hr
TBT1	1.63	2.43	4.46	5.40	7.20	7.43	8.63
TBT2	1.26	2.30	4.30	5.20	7.15	7.27	7.83
TBT3	1.36	2.36	4.43	5.37	7.50	7.47	8.60
TBT4	1.46	2.40	4.40	5.27	7.30	7.50	8.50
TBT5	1.30	2.63	4.30	5.30	7.30	7.60	8.80
TBT6	1.90	2.80	4.80	5.90	7.80	8.20	9.00
TBT7	1.86	2.73	4.43	5.60	7.60	7.63	8.93
TBT8	1.36	2.60	4.80	5.20	7.50	7.50	8.87
TBT9	1.50	2.50	4.26	5.70	7.70	7.80	8.90
TBT10	1.10	2.10	4.20	5.10	7.10	7.30	8.30
TBT11	1.43	2.50	4.50	5.20	7.20	7.60	8.73
TBT12	1.33	2.40	4.30	5.23	7.13	7.50	8.87
CD@5%	0.138	0.150	0.238	0.324	0.210	0.205	0.242
C V	5.536	3.539	3.165	3.564	1.684	1.602	1.641

Values are the mean of three replications

### Soil borne pathogens

The University of Svpuat Meerut's Faculty of Agri-Biotech kindly provided by isolated of *Fusarium* spp.

### DNA extraction from *Trichoderma* isolates

*Trichoderma* DNA was isolated according to Al-Samarrai and Schmid's (2000) instructions. The sharp, separate bands served as a quality indicator for the DNA.

### Isolation of Genomic DNA

The genomic DNA from twelve *Trichoderma* isolates was extracted by CTAB methods with some modification. The Genomic DNA of *Trichoderma* was isolated from cell which were in vegetative phase.

The cell suspension culture obtained by incubation of Potato Dextrose Broth (PDB)

*Trichoderma* mycelium and incubated for 36 hrs genomic DNA successfully isolated.

### Purification of crude genomic DNA

Isolated DNA samples were treated with 1µl of RNase (10mg/ml) mixed gently and incubated in boiling water bath at 37°C for 1 h. This was done to single stranded RNA. The purified DNA was precipitated and dissolve in 30µl TE buffer (pH 8.0). The quality of purified DNA of *Trichoderma* was checked on 0.8% agarose gel electrophoresis. The purified DNA did not show shearing or presence of RNA contamination. This DNA was purified.

### Molecular identification of *Trichoderma* isolates

Polymerase chain reaction (PCR) was utilized to amplify the internal transcribed spacer regions of *Trichoderma* using ITS1 (5' – TCC GTA GGT GAA CCT GCG G - 3') and ITS4 (5' - TCC TCC GCT TAT

TGA TAT GC - 3') primers. PCR conditions were performed as described by Loc et al. (2011). PCR products were first purified using QIAquick PCR Purification Kit (QIAGEN Cat. No. 28104). Sequencing was performed using Big Dye Terminator v3.1 Cycle Sequencing Kit in a total volume of 20  $\mu$ L using 3500 GeneticAnalyzer, Applied Biosystems (Daejeon, Korea).

#### PCR amplification of chitinase gene

According to Loc et al. (2011), the chitinase gene was amplified using two particular primers: CHI-F (5-ATG TTG GGC TTC CTC GGA-3) and CHI-R (5-TTC GGG ATG GTT GTC ATA CTG-3).

#### Antagonism of *Trichoderma* against *Fusariumoxysporum*

The pathogen and antagonist were inoculated in the same Petri plate opposite with each other and observation was recorded at every 24 hours after inoculation of both organisms up to 96 hours. The interaction was observed between *Trichoderma* isolates and *Fusariumoxysporum* at 4<sup>th</sup> day. The growth of *Fusariumoxysporum* was increases at and 5<sup>th</sup> day thereafter this was checked by the antagonist at 6<sup>th</sup> and 7<sup>th</sup> day.



Figure 1. Antagonism of *T. harzianum* against *Fusarium* spp.

All *T. harzianum* isolates exhibited higher growth inhibition activity of *Fusariumoxysporum* compared to the other species isolates. Among the *T. harzianum* isolates highest growth inhibition percentage was recorded in TBT6 (87.1%) followed by TBT7 (82.2%), whereas the minimum growth inhibition TBT10 (59.7%) was recorded in *T. longbrachiatum* isolates TBT10 after 7 DAI (Table 3.). *T. harzianum* isolate TBT6 showed initial faster growth, overgrew on *Fusariumoxysporum* after 5 days of inoculation.

Table 3- Growth inhibition of *Fusariumoxysporum* in dual culture by different *Trichoderma* isolates at  $28 \pm 2^{\circ}$  C

<i>Trichoderma</i> isolates	Growth inhibition efficiency of <i>Trichoderma</i> spp. in dual culture					
	5 <sup>th</sup> days after inoculation		6 <sup>th</sup> days after Inoculation		7 <sup>th</sup> days after inoculation	
	Radial Growth (cm)*	Growth Inhibition (%) <sup>#</sup>	Radial Growth (cm)*	Growth Inhibition (%) <sup>#</sup>	Radial Growth (cm)*	Growth Inhibition (%) <sup>#</sup>
TBT1	2.33	43.6	2.00	68.8	1.76	80.4
TBT2	3.66	11.4	3.40	48.2	2.83	68.6
TBT3	2.93	29.1	3.03	53.6	3.03	66.3
TBT4	3.03	26.6	2.86	56.1	2.63	70.8
TBT5	3.00	27.4	2.46	62.0	2.10	76.7
<b>TBT6</b>	<b>2.63</b>	<b>36.3</b>	<b>2.33</b>	<b>64.0</b>	<b>1.16</b>	<b>87.1</b>

TBT7	2.66	35.6	2.43	62.5	1.60	82.2
TBT8	2.63	36.3	2.53	61.0	2.16	76.0
TBT9	3.83	36.3	3.66	44.3	3.36	62.7
<b>TBT10</b>	<b>4.03</b>	<b>2.4</b>	<b>3.86</b>	<b>41.4</b>	<b>3.63</b>	<b>59.7</b>
TBT11	3.26	21.1	2.63	59.5	2.36	73.8
TBT12	3.66	11.4	3.5	46.7	3.13	65.2
Control	4.13	-	6.66	-	9.00	-
C.D.	0.20	-	0.19	-	0.176	-
C.V.	3.13	-	3.41	-	3.55	-

\*Values are the mean of three replications, #Values are the mean percentage inhibition

### Data and cluster analysis

The NCBI website (<http://www.ncbi.nlm.nih.gov/>) was used to compare the findings of *Trichoderma* sequencing with known sequences using BLASTn. MEGA version 6 was used for the alignment and phylogenetic analysis (Tamura et al., 2013). The [eostat-Costas6.3](#) version programme was used to [analyse](#) the antagonistic data that was obtained. Use of the Duncan's multiple range test (1955) guided the analysis of variance and means comparison at the 5% level of significance.

### Quantitative estimation of chitinase activity of *Trichoderma* isolates

All *Trichoderma* isolates were used for biochemical assay for the production of cell wall degrading enzymes. Chitinase activity was assayed by measuring the release of reducing saccharides from colloidal chitin. A [standard](#) curve generated with N-acetyl- $\beta$ -D-glucosamine (NAGA) was used to determine reducing saccharides concentration. The activity ranged from 5.77 unit/ml to 21.15 unit/ml commercial chitin-derived colloidal chitins of the TBT5 and TBT6 isolates respectively; while TBT7 isolate presented 20.12 unit/ml chitinase enzyme activities

**Table 4. Quantitative estimation of chitinase activity (U/ml) from different species of *Trichoderma***

S. N.	Isolates	Chitinase activity (Unit/ml)
TBT1	<i>Trichodermaasperellum</i>	12.30
TBT2	<i>Trichodermaasperellum</i>	18.23
TBT3	<i>Trichodermaasperellum</i>	14.26
TBT4	<i>Trichodermaasperellum</i>	9.17
TBT5	<i>Trichodermalongibrachiatum</i>	6.18
TBT6	<i>Trichodermaharzianum</i>	21.15
TBT7	<i>Trichodermaharzianum</i>	20.12

TBT8	<i>Trichodermaharzianum</i>	18.34
TBT9	<i>Trichodermalongibrachiatum</i>	18.63
TBT10	<i>Trichodermalongibrachiatum</i>	5.77
TBT11	<i>Trichodermakoningii</i>	17.28
TBT12	<i>Trichodermakoningiopsis</i>	7.29


## RESULTS AND DISCUSSION

### Isolation and molecular identification of *Trichoderma* isolates

From rhizosphere soil samples taken from several locations in Uttar Pradesh, 12 strains of *Trichoderma* were identified (Table 1). The ITS region was amplified by PCR using ITS primers, yielding a single band of roughly 600 bp. After the PCR products were sequenced, the sequencing data was uploaded to the NCBI website for BLAST searches and comparison with previously released ITS data. Four isolates were found to belong to *Trichodermaasperellum* (TBT1, TBT2, TBT3, TBT4), three isolates were classified as *Trichodermaharzianum* (TBT6, TBT7, TBT8), and two isolates were identified as *Trichodermalongibrachiatum*. The amplified ITS regions of the *Trichoderma* isolates showed 99 to 100% identity with three species of *Trichoderma*. (Table 1). The ITS identity was validated by the phylogenetic analysis.


### Phylogenetic analysis of *Trichoderma*

Phylogenetic relationship among the *Trichoderma* species was determined using the ITS sequence information of 12 *Trichoderma* spp. and 23 other *Trichoderma* sequences, including *T. asperellum*, *T. harzianum*, *T. longibrachiatum*, *T. koningii* and *T. koningiopsis* available in the NCBI database. A dendrogram was constructed by using the ClustalW and Mega7.0 software with the Neighbor-Joining (NJ) method.



**Figure 2.** Phylogeny analysis of *Trichoderma* isolates based on ITS data using MEGA.

UNDER PEER REVIEW




**Fig.3** Amplified PCR products of ITS region of 12 native *Trichoderma* spp. isolates from soil using ITS universal primers. Lane M: 100 bp ladder (Promega, USA), Lane 1-12: TBT1-TBT12

#### **Amplification of chitinase gene from *Trichoderma harzianum***

The genomic DNA isolated from *T. harzianum* was PCR amplified with specific primers for chitinase gene. The concentration (10 pM/μl) of respective primers was used in the PCR amplification. Through amplification, a good unique band was observed without any primer dimers. The large-scale amplification of putative full-length of chitinase gene approx (1.3 kb).

#### **4.9. Gel elution of the PCR amplicon**

The PCR products were separated in a low melting agarose gel of 1.2% along with 1kb DNA molecular marker. Very sharp bands of expected sizes were obtained. The 1.3 kb amplicon corresponding to chitinase gene were eluted out using QIAEX II Gel Extraction Kit.



**Figure 4.** Amplification of chitinase gene in *Trichoderma* isolates based CHI-F and CHI-R primers, M; 1 kb DNA ladder.

### **Chitinase gene of *Trichoderma***

**Chitin** is an essential part of these pathogen cell walls, *Trichoderma* secretes extremely potent chitinases that may degrade and feed on them. (Seidl-Seiboth et al., 2014; Hassan et al., 2015; Prasetyawan et al., 2018). Chitinase 42 is one *Trichoderma* endochitinase that can hydrolyze the  $\beta$ -1, 4-glycosidic bonds that hold the N-acetyl glucosamine residues in chitin together. (Hassan et al., 2015). Using PCR-based specialist primers (CHI-F and CHI-R), the chitinase 42 gene in *Trichoderma* isolates was amplified and discovered in this experiment, as shown in Figure 4. The presence of this gene was confirmed in all investigated isolates by the PCR, which a single band around 1500 bp similar by Loc et al. (2011). Since the primers are unique to *Trichoderma* chitinases, the amplification of the chitinase gene verifies that these isolates are *Trichoderma*.

### **Conclusion**

Identification of *Trichoderma* based on ITS markers exhibited high efficiency in discrimination among different *Trichoderma* spp. isolates. *Trichoderma* isolates showed variability in their aggressiveness against *Fusarium oxysporum*. In general, the isolate TBT6 of *T. harzianum* showed the best antagonism against *Fusarium oxysporum* and while the isolate. *Trichoderma* isolates improved. These results confirm the efficacy of *Trichoderma* as excellent biocontrol agent and also as plant growth promoting. *Trichoderma* under *in-vitro* conditions, against *Fusarium oxysporum* dual culture bioassay. Among the *T. harzianum* isolates highest growth inhibition percentage was recorded in TBT6 (87.1%) followed by TBT7 (82.2%), whereas the minimum growth inhibition (59.7%) was recorded in *T. longibrachiatum* isolates TBT10 after 7 Days. Chitinase activity was assayed by measuring the release of reducing saccharides from colloidal chitin. Standard curve generated with N-acetyl- $\beta$ -D-glucosamine (NAGA) was used to determine reducing saccharides concentration. The activity ranged from 5.17 unit/ml to 21.15 unit/ml commercial chitin derived colloidal chitins of the TBT10 and TBT6 isolates respectively; while TBT7 isolate presented 20.12 unit/ml chitinase enzyme activities.

### **REFERENCES**

Abdel-lateif K, El-

Zanaty AF, Helwa M (2017). Genetic diversity of *Trichoderma* isolates and their antagonism against *Rhizoctonia Solanica* and *Phytophthora infestans*. *Menoufia Journal of Agriculture Biotechnology* 2:63-73.

Agrawal T, Kotasthane AS (2012). Chitinolytic assay of indigenous *Trichoderma* isolates collected from different geographical locations of Chhattisgarh in Central India. *SpringerPlus* 1:73-82

Al-Samarrai TH, Schmid J (2000). A simple method for extraction of fungal genomic DNA. *Letters in Applied Microbiology* 30:53-56.

Arie T. (2019). *Fusarium* diseases of cultivated plants, control, diagnosis, and molecular and genetic studies. *Journal of Pesticide Science*, 44: 275-281.

Athira K (2017). Efficacy of fungicide and bio-control agents against root rot of black Gram (*Vigna mungo* L.) caused by *Macrophomina phaseolina* (Tassi) Goid. *International Journal of Current Microbiology and Applied Sciences* 6(10):2601-2607.

Benítez T, Rincón AM, Limón MC, Codón AC (2004). Biocontrol mechanisms of *Trichoderma* strains. *International Microbiology* 7:249-260

- Chen S, Dickson DW (2004). Biological control of nematodes by fungal antagonists. In: Nematology Advances and Perspectives. Nematode Management and Utilization, vol. II. (Chen ZX, Chen SY, Dickson DW, eds). Wallingford: CAB International, UK pp. 977-1039.
- Daykin ME, Hussey RS (1985). Staining and histopathological techniques in nematology. pp. 39-48. In: K.R. Barker, C.C. Carter, J. N. Sasser, eds. An Advanced Treatise on *Meloidogyne*: Volume II. Raleigh, NC: North Carolina State University Graphics.
- Duncan B (1955). Multiple range and Multiple F test. *Biometrika* 11:1-42.
- Elad Y, Hadar Y, Chet I, Henis Y (1981). Biological control of *Rhizoctonia solani* by *Trichoderma harzianum* in carnation. *Plant Disease* 65:675-677.
- El-Hassan SA, Gowen SR, Pembroke B (2013). Use of *Trichoderma hamatum* for biocontrol of lentil vascular wilt disease: efficacy, mechanisms of interaction and future prospects. *Journal of Plant Protection Research* 53(1).
- Epstein E (1972). *Mineral Nutrition of Plants: Principles and Perspectives*. John Wiley, New York.
- Ganiger M. C., Bhat S., Chetri P. and Kuruvinashetti M. S. (2009). Production of endoglucanase by *Trichoderma* for control of phytopathogenic fungus *Sclerotium rolfsii*. *Journal of Applied Sciences Research*, 5(7): 870-875
- Goody JB (1957). Laboratory method for work with plant and soil nematodes. Ministry of Agric., Fisheries and Food: Tech. Bull. 2 London, England 44 p.
- Gordon T. R. (2017). *Fusarium oxysporum* and the *Fusarium* wilt syndrome. *Ann. Rev. Phytopathol.* 55 : 23-39.
- Guarro J. (2013). Fusariosis, a complex infection caused by a high diversity of fungal species refractory to treatment. *European Journal of Clinical Microbiology & Infectious Diseases*. 32, 1491–1500.
- Hartman K, Sasser JN (1985). Identification of *Meloidogyne* species on the basis of differential hosts test and perineal pattern morphology. In: Barker KR, Carter CC, Sasser JN, editors. An advanced treatise on *Meloidogyne*. Volume II. North Carolina State University Graphics, Raleigh, North Carolina: Methodology. A cooperative publication of the Department of Plant Pathology and the United States Agency for International Development pp. 69-77.
- Hassan M, EL-fiky ZA, Said MA, Ahmed NF (2015). Chitinase and cellulase genes sequencing for some Egyptian *Trichoderma* species isolated from rhizosphere and assay of their activity. *Egyptian Journal of Genetics and Cytology* 44:15-30.
- Hussey RS, Barker KR (1973). Comparison of methods for collecting inocula of *Meloidogyne* spp., including a new technique. *Plant Disease Report* 57:1025-1028.
- Izuogu NB, Abiri TO (2015). Efficacy of *Trichoderma harzianum* T22 as a biocontrol agent against root-knot nematode (*Meloidogyne incognita*) on some soybean varieties. *Croatian Journal of Food Science and Technology* 7(2):47-51.
- Jansson HB, Jeyaprakash A, Zuckerman BM (1985). Control of root-knot nematodes on tomato by the endoparasitic fungus *Mericoniospora*. *Journal of Nematology* 17:327-329.

- Junaid JM, Dar NA, Bhat TA, Bhat AH, Bhat MA (2013). Commercial biocontrol agents and their mechanism of action in the management of plant pathogens. *International Journal of Modern Plant and Animal Sciences* 1(2):39-57.
- Karsen G, Moens M (2006). Root-knot nematodes. In: Perry, R.N. and Moens, M. (Eds). *Plant Nematology*. CAB Publishing pp. 59-90.
- Krishnamurthy YL, Shashikala J (2006) Inhibition of aflatoxin B1 production of *Aspergillus flavus*, isolated from soybean seeds by certain natural plant products. *Letters in Applied Microbiology* 43:469-474.
- Kumar A., Sahu T. K., Bhalla A. and Jain, A. K. (2013) Morphological characterization of *Trichoderma harzianum* from Madhya Pradesh. *Annals of Plant Protection Sciences* 22(1): 190-239.
- Lidia B., Siwulski M., Sobieralski K., Lisiecka J. and Jędryczka M. (2014). *Trichoderma* spp. application and prospects for use in organic farming and industry. *Journal of plant protection research* 54(4) 2014-0047.
- Lin B., Bozorgmagham A., Ross S. D., Schmale and D.G. (2013). Small fluctuations in the recovery of fusaria across consecutive sampling intervals with unmanned Aircraft 100 m above Ground Level. *Aerobiologia*, 29: 45-54
- Mascarin GM, Junior MFB, Filho JV de A (2012). *Trichoderma harzianum* reduces population of *Meloidogyne incognita* in cucumber plants under greenhouse conditions. *Journal of Entomology and Nematology* 4:54-57.
- Pal KK, Gardener BM (2006). *Biological Control of Plant Pathogens*. The Plant Health Instructor. <https://www.apsnet.org/edcenter/advanced/topics/Pages/BiologicalControl.aspx>.
- Pandya JR, Sabalpara AN, Chawda SK (2011). *Trichoderma*: a particular weapon for biological control of Phytopathogens. *Journal of Agricultural Technology* 7(5):1187-1191.
- Perfect S. E. and Green, J.R. (2001). Infection structures of biotrophic and hemibiotrophic fungal plant pathogens. *Molecular Plant Pathology*, 2: 101-108
- Prasetyawan S, Sulistyowati L, Aulanni'am (2018). Glucanase and chitinase from some isolates of endophytic fungus *Trichoderma* spp. *IOP Conference Series: Materials Science and Engineering* 299:012026.
- Puyam A. (2016). Advent of *Trichoderma* bio-control agent. *Journal of Applied and Natural Science* 8 (2) 1100 - 1109
- Rao KLN M, Raju KS, Ravisankar H (2015). Cultural conditions on the production of extracellular enzymes by *Trichoderma* isolates from tobacco rhizosphere. *Brazilian Journal of Microbiology* 47(1):25-32
- Rayatpanah S, Nanagulyan SG, Alav SV, Razavi M, Ghanbari-Malidarreh A (2011). Pathogenic and genetic diversity among Iranian isolates of *Macrophomina phaseolina*. *Chilean Journal of Agricultural Research* 72:40-44.
- Seidl-Seiboth V, Ihrmark K, Druzhinina I, Karlsson M (2014). Molecular evolution of *Trichoderma* chitinases. *Biotechnology and Biology of Trichoderma* 2014:67-78.
- Sharma R. A. (2012). Brief review on mechanism of *Trichoderma* fungus use as biological control agents. *International Journal of Innovations in Biological and Chemical Sciences* 2: 200-210.
- Sharma RA (2012). Brief review on mechanism of *Trichoderma* fungus use as biological control agents. *International Journal of Innovations in Biological Sciences* 2:200-210..

- Spiegel Y, Sharon E, Bar-Eyal M (2007). Evaluation and mode of action of Trichoderma isolates as biocontrol agents against plant-parasitic nematodes. *IOBC.WPRS.Bulletin*30:129-133.
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013). MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution*30:2725-2729.
- Woo SL, Ruocco M, Vinale F, Nigro M, Marra R, Lombardi N, et al (2014). Trichoderma-based Products and their Widespread Use in Agriculture. *The Open Mycology Journal*8(1).
- Wu Q, Sun R, Ni M, Yu J, Li Y, Yu C, Dou K, Ren J, Chen J (2017). Identification of a novel fungus, *Trichoderma asperellum* GDFS1009, and comprehensive evaluation of its biocontrol efficacy. *PLOS ONE*12(6):e0179957.
- Xia R., Schaafsma A.W. F., and Hooker D. C. (2020). Impact of the improvements in Fusarium head blight and agronomic management on economics of winter wheat. *World Mycotoxin Journal*, 1–18.