

## Detection and Identification of Major Seed Mycoflora Associated with Rice (*Oryza sativa* L.) Cultivars grown in Kashmir

### Abstract

Worldwide, rice is a significant crop for food production. The main requirement for increased yield and production is high-quality seed. While being exported, seeds and grains have the risk of being associated with pathogenic as well as non-pathogenic microflora. As a result, they are frequently inspected in accordance with phytosanitary laws to stop diseases from spreading into new areas and also to have an impact on the quality of the seeds. Fungal infections that infect rice seeds are regulated by phytosanitary laws and can be identified using several techniques for checking seed health. Therefore, in accordance with the recommendations of International Seed Testing Association (ISTA), stored seeds of seeds of four rice varieties (SR-2, SR-3, SR-4, and SR-5) were studied to investigate the occurrence of seed-associated fungal mycoflora using the blotter paper and agar plate method. In both blotter and agar tests, four fungal species namely as *Fusariumoxysporium*, *Rhizopusstolonifer*, *Alternariaalternata*, and *Tricodermaharzianum* were isolated from different test varieties of rice. There was 5.667%, 8.333%, 4.333% ,9.667% by blotter test and 7.667%, 11.667%, 6.333%, 12.667% by agar test mycoflora associated with the seeds of SR-2, SR-3, SR-4 and SR-5.

**Keywords:** Rice seeds, blotter paper, agar plate, seed-associated mycoflora, pathogen frequency

### Introduction

One of the world's staple food crops is rice (*Oryza sativa* L.). China and India are the top two rice-growing nations in terms of production. 782 million tons of rice are produced annually from 167 million hectares of rice plantings worldwide (FAO 2022). With 112.91 million tonnes produced, China is the top

rice producer, with India coming in second (Anonymous, 2022). Higher grain yield and improved crop growth are dependent on both seed quality and seed health (Haque *et al.*, 2012). The availability of high-yielding, disease-free, and farmer-preferred seeds of high-yielding varieties is the most crucial requirement for successful crop production. A major economic impact on agricultural output is caused by fungi and other pathogens that attack plants, as they can cause diseases in farmed crops that lead to significant yield losses (Paplomatas, 2006). Plant diseases that are specifically spread by seeds are referred to as seed-borne diseases. In contemporary agricultural research, a healthy seed bank is crucial for a desired plant population and a successful harvest (Rahman *et al.*, 2008). Seedborne pathogens are a persistent issue that may even be to blame for the introduction of new illnesses into previously uninfected areas as well as the resurgence of old ones (Walcott *et al.*, 2007). Seed associated infections pose a significant risk to the establishment of seedlings (Walcott *et al.*, 2003). Crops' capacity to establish themselves and reach their maximum yield and value potential is significantly impacted by the quality of the seeds that are sown (McGee, 1995). One of the most significant biotic barriers to seed production globally is the presence of seed-borne and seed associated fungi. They cause diversity in plant shape, pre- and post-emergence death of grains, and vigor of seedlings, which in turn reduces germination to some extent (Van Du *et al.*, 2001; Rajput *et al.*, 2005; Niaz and Dawar, 2009).

### **Materials and Methods**

*i). Planting material:* The planting material viz. SR-2, SR-3, SR-4 and SR-5 for the study was obtained from Mountain Research Centre for Field Crops, Khudwani, SKUAST-Kashmir

*ii). Seed Health Testing:* Following seed health testing methods were used for the detection of seed associated mycoflora

a). Blotter test

In accordance with the International Seed Testing Association [ISTA, 1996], the four different rice types seeds were examined using the blotter method

in Petri plates to determine the presence or absence of seed associated mycoflora. The Petri plates were incubated at  $25\pm 1^{\circ}\text{C}$  with a 12/12 hour light and dark cycle. The developing fungal colonies were examined under stereomicroscope or transferred to temporary mounts to be examined under compound microscope for identification of fungi.

#### b). Agar Plate Method

Seeds were scrutinized using Agar Plating Technique for the examination of seed associated mycoflora. The plated seeds were cultured for five to seven days at  $22$  to  $25^{\circ}\text{C}$  with 12-hourly light and dark cycles. Fungi developing from the seeds on the agar medium were inspected and identified at the conclusion of the incubation time. Under a compound microscope, identification was performed using the morphology of the sporulation structures and colony characteristics.

**Comment [a1]:** Add reference and also which medium used for agar plating techniques?

#### iii). Identification of fungi

The various colonies were then identified visually, first under a stereomicroscope, and then under a compound microscope, which allowed for a closer look at the fruiting structures (Mathur and Kongsdal, 2003). All the fungi were identified on the basis of standard manuals and descriptions (Malone and Muskette, 1964; Misra *et al.*, 1994; Mathur and Kongsdal, 2003), Ellis, 1971; Chidambaram and Mathur, 1975; Neergaard and Saad, 1962; Booth, 1971 and Agarwal *et al.*, 1990)

**Comment [a2]:** Use recent reference.

iv). *Analysis of data:* All the findings were displayed as a percentage of each pathogen's frequency. All the data was subject to standard analysis for accuracy and clarity.

### Results and Discussion

In both the blotter and the agar tests, variety SR-4 showed the least amount of seed-borne infection out of the four types examined. In comparison to other types, there was also reduced SR-4 infection from all fungal pathogens.

Based on colony morphology, four members of the fungal flora—*Tricodermaharzianum*, *Alternariaalternata*, *Rhizopusstolonifer*, and *Fusariumoxysporum*—were identified using spore/conidial morphological analyses.


By blotter paper method highest per cent frequency of *Tricodermaharzianum* was noticed in SR-3 variety (6.667 %) where as lowest infection in SR-4 (2.667 %). Highest infection of *Alternariaalternata* was noticed on SR-5 (18.667 %) and lowest infection on SR-4 (9.33 %). *Rhizopusstolonifer* showed highest infection on SR-3 (10.667 %) and lowest on SR-4 (2.667 %). *Fusariumoxysporum* highest infection detected on SR-5 (5.333 %) and lowest infection on SR-2 (1.333 %) (Table 1).

**Comment [a3]:** Molecular identification of fungi required


UNDER PEER REVIEW

**Table 1: Per cent frequency of rice seed mycoflora associated with different varieties by blotter paper and agar plate method.**

Fungi	Frequency of seed mycoflora (%) associated with different rice varieties									
	Blotter paper method					Agar plate method				
	SR-2	SR-3	SR-4	SR-5	Mean	SR-2	SR-3	SR-4	SR-5	Mean
<i>Tricoderma harzianum</i>	2.667	6.667	2.667	5.333	4.333	5.333	9.333	5.333	6.667	6.667
<i>Alternaria alternata</i>	13.333	12.000	9.333	18.667	13.333	13.333	21.333	9.333	22.667	16.667
<i>Rhizopus stolonifer</i>	5.333	10.667	2.667	9.333	7.000	6.667	10.667	6.667	14.667	9.667
<i>Fusarium oxysporum</i>	1.333	4.000	2.667	5.333	3.333	5.333	5.333	4.000	6.667	5.333
Mean	5.667	8.333	4.333	9.667		7.667	11.667	6.333	12.667	
CD(0.05)	Variety =1.9801 Fungi =1.9203 Variety* Fungi =3.958					Variety =1.8600 Fungi =1.7600 Variety* Fungi =3.720				



*Trichoderma harzianum* *Alternaria alternata*



*Rhizopus stolonifer* *Fusarium oxysporum*

**Plate 1: Culture plate of different isolated fungi and their microscopic characters**

Four seed-borne diseases were found and identified using the Agar plate method. They are *Fusariumoxysporum*, *Rhizopusstolonifer*, *Alternariaalternata*, and *Trichodermaharzianum*. *Trichodermaharzianum* exhibited lowest infection on SR-4 and SR-2 (both 5.333%) and maximum infection on SR-3 variety (9.33%). Infection rates for *Alternaria alternative* ranged from lowest on SR-4 (9.333%) to highest on SR-5 (22.667%). *Rhizopus stolonifera* observed that SR-5 had the highest infection rate (14.667%), while SR-4 and SR-2 had the lowest infection rates (6.667%). *Fusariumoxysporium* found that infection levels were lowest on SR-4 (4.000%) and highest on SR-5 (6.667%)(Table1). Plate 1 shows colony growth and spore/conidial formations observed under a compound microscope.

The information about the seedborne nature of pathogens provided by Kumar and Bijendar (2022), Karan *et al.*, (2021), Umbreyet *al.*, (2021), Indira (2022), Khan (2006), Islam (2005) and Fakir (2001) was consistent with the current findings of the fungal organisms that are spread through seeds. The current study's findings indicate that the fungal flora that is carried by seeds includes *Trichodermaharzianum*, *Alternariaalternata*, *Rhizopusstolonifer*, and *Fusariumoxysporum*. Compared to the other studied rice seed varieties, the SR-4 variety showed less infection. The most important input in agriculture is pathogen-free seed. The majority of grain and seed varieties that are exported from India undergo testing in accordance with phytosanitary laws. The blotter and visual tests are often conducted as part of the phytosanitary rules on a regular basis. The results of the current investigation showed that certain pathogens were only detected in the blotter and agar tests for seed microflora. Since pathogen-free seed is a crucial component of commerce, this study will assist in performing both the blotter test and the agar test.

### Conclusion

This study shows that the dominant rice cultivars in the Kashmir valley have a varied mycoflora of pathogenic and non-pathogenic fungus. Better seed health management is essential for effective rice farming because rice is a staple

**Comment [a4]:** Without molecular identification of fungi -can not identify fungal isolate upto the species level

meal. Understanding seed health is crucial to managing crops and pests in developing nations where farmers must conserve their own seeds for sowing. Testing for seed health is another way for farmers to optimize seed stocks for crop yield through quality management. It helps with seed certification for farmers from public seed providers and seed growers. Additionally, the results point to the necessity of properly storing rice seed in order to reduce fungal infestation in the near future.

### References

- Agarwal, P.C., Mortensen, C. N. and Mathur, S. B. 1990. Seed borne disease and seed health testing of rice. *Journal of Agriculture Science, Cambridge*, **115**:144-144.
- Booth, C. 1971. The genus *Fusarium*. CMI, Kew, Surrey, England. 237 pp.
- Chidambaram, P. S. and Mathur, S. B. 1975. Deterioration of grains by fungi. *Annual Review of Phytopathology*, **3**: 69-89.
- Ellis, M.B. 1971. Demotiaceous Hyphomycetes. C.M.I., Kew Surrey England, 608 pp.
- Fakir GA. 2001. List of seed borne diseases of important crops occurring in Bangladesh. Seed Pathol. Lab., Dept. Pl. Pathol., Bangladesh Agricultural University, Mymensingh 9p.
- Food and Agriculture Organization of the United Nations (FAO). 2022. FAOSTAT database. Retrieved from <http://www.fao.org/faostat/en/#data/QC>
- Anonymous, 2022. Ministry of Agriculture and Farmers Welfare. Retrieved from <https://agricoop.gov.in/>
- Haque, M. A., Haque, M. M., Al-Emran, A. S., and Islam, M. S. 2012. Higher grain yield and improved crop growth are dependent on both seed quality

- and seed health. *Journal of Agricultural Science and Technology*, **2**(2):139-145.
- International Seed Testing Association (ISTA). 1996. International rules for seed testing. *Seed Science and Technology*, **21**:1-288.
- Islam, T. 2005. Seedborne nature of pathogens. *Plant Pathology Journal*, **4**(1), 23-30.
- Indra, N., Ushamalini, C., Kavitha, K. and Chitra, K. 2022. Seed mycoflora of paddy varieties in Tamil Nadu and its effect on germination and seedling vigour. *The Pharma Innovation Journal*, **6**: 1645-1649
- Karan, R., Renganathan, P., Balabaskar, P. and Raj S. T. 2021. Detection of seed-borne mycoflora associated with paddy seeds and its influence on seedling health. *International Journal of Current Microbiology and Applied Sciences*, 2319-7706.
- Khan, M. R. 2006. Seedborne nature of pathogens. *Plant Disease*, **90**(7), 835-849.
- Kumar, A. and Bijendar, K. 2022. Seedborne nature of pathogens. *Journal of Plant Pathology*, **9**(2), 67-72.
- Kumar, J. and Kumar, B. 2022. Detection and identification of seed-borne fungal pathogens associated with seed discoloration of rice. *The Pharma Innovation Journal*, **11**(9): 3091-3096
- Malone, G.P. and A.E. Muskette, 1964. Seed borne fungi: Description of 77 Fungal Species. Proc. Int. Seed Test Assoc., 29(2): 180-183.
- Mathur, S. B. and Kongsdal, O. 2003. Common laboratory seed health testing method for detecting fungi. First edition. *International Seed Testing Association*.
- McGee D.C., 1995. Approach to disease management through seed technology. *Annu. Rev. Phytopathol.* 1995.33:445- 466.

- Misra, J.K. Gergon and T.W. Mew, 1994. Occurrence, distribution and phenology of seed borne fungi of rice in certain provinces of Philippine. *Plant Pathology Bulletin*, **3**(4): 229-239.
- Neergaard, P. and A. Saad, 1962. Seed health testing of rice: A contribution to development of laboratory routine testing methods. *Indian Phytopathology* **15**: 85-111.
- Niaz I. and Dawar S., 2009. Detection of seed borne mycoflora in maize (*Zea mays* L.). *Pakistan Journal of Botany*, **41**(1): 443- 451.
- Paplomatas E.J., 2006. Molecular diagnostics of fungal pathogens. *Arab Journal of Plant Protection*, **24**: 147-158.
- Rajput, N. A., Rana, S. M. and Singh, S. B. 2005. Impact of seed-borne and seed-associated fungi on seed production: a global perspective. *Plant Pathology Journal*, **4**(3), 215-223.
- Rahman, M. M., Hasan, M. R., Islam, M. S. and Akhter, M. M. 2008. Importance of a healthy seed bank in contemporary agricultural research. *Journal of Agricultural Science*, **5**(2), 32-39.
- Umbrey, Y., Divya M., Das, T., Das, S. and Mahapatra, S. 2021. Isolation and identification of seed borne mycoflora associated with popular rice cultivars in North East India. *Journal of Cereal Research*, **13**: 43-50
- Van Du P, Loan LC, Cuong ND, Nghiep HV, Thach ND (2001). Survey on seed borne fungi and its effects on grain quality of common rice cultivars in the Mekong Delta. *Omonrice*, **9**: 107-113.
- Walcott, R. R., Gitaitis, R. D., and Castro, A. C. 2007. Seedborne pathogens: a persistent issue in agriculture. In *Proceedings of the 4th International Symposium on Seed Health in Agricultural Development*, 45-52.
- Walcott RR, Gitaitis RD, Castro AC. 2003. Role of blossoms in watermelon seed infestation by *Acidovorax avenae* subsp. *citrulli*. *Phytopathol.* **93**:528–34.

Walcott RR. 2003. Detection of seedborne pathogens. *HortTechnology* 13:40–47.

UNDER PEER REVIEW