

## Effect of Varieties, Solar Drying and Zeer Pot Refrigeration Lining Media on **Phytochemical** Properties of Chili Pepper Fruits

### ABSTRACT

Chili peppers are rich in **phytochemical** compounds such as carotenoids, beta carotene, phenols, flavonoids, capsaicinoids which protect human body against diseases such as cancer, heart diseases, diabetes but deteriorate more rapidly. Therefore, the main objective of this study **was** to investigate the effect of varieties, solar drying and zeer pot lining media on phytochemical characteristics of chili peppers. A 2 x 3 x 3 factorial experimental design in Randomized Complete Block Design (RCBD) was used **to collect data from the experimental field for laboratory analyses of phytochemicals**. Results of effect of varieties on phytochemical characteristics of chili pepper fruits revealed that, red cayenne was significantly greater in total carotenoids (9222ug/100g), beta carotene (934.23ug/100g), total flavonoids (50.04mg/100g), total capsaicinoids (247.30mg/100g) but scotch bonnet was significantly higher in total phenols (301.74mg/100g). Again, on results of effect of solar drying on **phytochemical** characteristics of chili pepper fruits showed that, unblanched chili pepper fruits had significantly ( $p < 0.05$ ) higher total phenols (386.64mg/100g), total flavonoids (51.97mg/100g) and total capsaicinoids (260.72mg/100g) while controlled chili pepper fruits had significantly higher total carotenoids (8035.4ug/100g) and beta carotene (812.34ug/100g). Also, on results of effect of lining media of zeer pot refrigeration on **phytochemical** characteristics of chili pepper fruits; sand lining media of zeer pot refrigeration had significantly ( $p < 0.05$ ) higher total phenols (304.30mg/100g), total flavonoids (49.99mg/100g) and total capsaicinoids (242.17mg/100g) while styrofoam lining media had significantly higher total carotenoids (8506.1ug/100g) and beta carotene (861.64ug/100g).

**Keywords:** Varieties, solar drying, zeer pot, **phytochemicals**, chili peppers

### INTRODUCTION

Chili peppers are cultivated all over the world with leading countries been China, Mexico and Turkey which they contribute about 70% of the world's chili pepper cultivation[1]. Ghana is ranked eleventh (11<sup>th</sup>) producer of chili pepper in the world and second (2<sup>nd</sup>) producer in Africa with estimated total production in terms of metric tons been 88, 000 metric tons [2].In

Ghana, it is normally cultivated by small scale or commercial farmers for export or for domestic consumption [3].

They contain various food nutrients such as protein, carbohydrate, vitamin A and C, folic acid, fibre and low in sodium and caloric content [4] and phytochemical substances such as carotenoids, beta carotene, flavonoids, phenols, capsaicinoid etc which are antioxidants which defend human body against disease infections such as cancer, diabetes and heart diseases, diarrhoea and arthritis [5], [6]. They are added to food such as meat, fish, vegetable, salad and soup to make it spicy and add flavour to food [7]. Heat is extracted from chili peppers to prepare various alcoholic beverages and culinary as well as pharmaceutical products [8]. Capsaicin is also extracted from chili peppers to prepare tear gas or pepperspray which is used to control crowd or riot by law enforcement officers. They are also used to prepare organic insecticides or pesticides which is environmental friendly and do not pose health threats to humans after consumption of the product [9].

Solar dryer is a drying device which harvest the sun radiation and use for drying agricultural products hygienically [10] by reducing contamination by dust, rain, insect or pests there by improving the quality of agricultural products [11].


Zeer pot also referred to as pot – in – pot or evaporative cooler is a storage device use for storing agricultural products which is capable of reducing temperature of the agricultural products by 15 - 40<sup>0</sup>F resulting in the extension of shelf – life of the agricultural products without using electricity or fossil fuel. It is low cost as compared to electrical refrigerator and environmental friendly [12], [13].

Chili peppers deteriorate quickly after harvest, during processing and storage which results in scarcity of chili peppers in the dry season there by depriving humans of the use of chili peppers which cause many ailments associated with inadequate consumption of chili peppers and increase the price of chili peppers in the lean seasons. Global annual postharvest loss of vegetables is estimated to be 30 – 70% [14]. In Ghana postharvest loss of fruits and vegetables is also estimated to be 20 – 50% [15]. Therefore, it is important to investigate the effect of varieties, solar drying and zeer pot lining media on phytochemical characteristics of chili peppers, to make chili peppers available all year round for human use.

## **MATERIALS AND METHODS**

### **Study Area**

The study was conducted in the Dormaa – East District with total land area of 456 Square Kilometres. It lies within the middle belt of Ghana, with latitude between 7<sup>0</sup> 08<sup>1</sup> North and 7<sup>0</sup> 25<sup>1</sup> degrees and longitude 2<sup>0</sup> 35<sup>1</sup> West and 2<sup>0</sup> 48<sup>1</sup> West [16].



**Figure 1. Map of the study area**

## **Source of Chili Pepper Fruits and the Various Lining Media for Solar Drying and Zeer**

### **Pot Refrigeration**

One bag of fresh scotch bonnet and one bag of red cayenne were bought from chili pepper traders in the various major market centres in the respective communities, Asuotiano, Wamfie and Wamanafa. The various lining materials (one (1) bag each) of sand (0.02mm), styrofoam (0.05mm) and wood shavings (thickness = 0.02mm and length = 4.5 cm) were collected from each respective communities.

### **Preparation of Chili Pepper Fruits**

Chili pepper fruits were sorted to remove diseased fruits, washed and separated into two (2). One (1) group was blanched for three (3) minutes at 100<sup>0</sup>C using coalpot and thermometer. The other group was unblanched.

### **Procedures for Solar Drying Chili Pepper Fruits**

Blanched chili peppers were dried from 8:00 am to 6:00 pm for two (2) weeks and unblanched chili pepper fruits were dried for three (3) weeks. After drying of chili pepper fruits, the fruit stalks were removed or destalked.



**Figure 2. Solar dryer**

### **Procedures for Zeer Pot Refrigeration of Chili Peppers**

The zeer pots were labelled with the use of tape and marker. The bottom of the outer clay pots were filled to a height of 6 cm with the various lining materials of zeer pot. The inner clay pots were placed in outer clay pots and the space between inner and outer clay pots were filled with various lining media (sand (sieved), styrofoam (crushed) and wood shavings) to the top level. The weight of the zeer pots were measured using weighing scale with zeer pots filled with styrofoam, wood shavings and sand weighing 6.96kg, 7.18kg and 15.52kg respectively. Solar dried chili pepper varieties were placed in the various labelled zeer pot refrigerators. Each zeer pot refrigerator was kept 14cm apart (14 cm between rows and 14 cm within rows and each experimental set up was kept in each three (3) communities for three (3) month under room temperature (26 - 34<sup>0</sup>C ). Water (600ml) was used to replenish zeer pots every day **in** each morning.



**Figure 3. Zeer pot refrigerators with different lining media**

### **Phytochemical Properties Analyses**

Phyto – chemical characteristics analyses were carried - out to determine the effect of varieties, solar drying and zeer pot refrigeration lining media on phytochemical characteristics of chili pepper fruits.

### **Determination of total carotenoids**

The total carotenoid was calculated using [17]method. Glass wool was used to filter the decanted liquid from a 1 g sample that had been pulverised using 50 ml of acetone in a 50 ml volumetric flask. The sample was ground until no more colour could be extracted from it and the extract was colourless. After the filtrate was moved, 30 millilitres of petroleum ether were previously put to a 250 millilitre separating funnel.

A little over 250 millilitres of distilled water were added to the mixture gradually while allowing it to run over the funnel's walls. The aqueous (lower) phase was eliminated once the two phases separated. To get rid of any leftover acetone, the top phase was cleaned four times with 250 ml of distilled water each time. During the final washing, all of the lower phase was thrown away, leaving none of the upper stages behind. After that, the petroleum ether phase was collected in a volumetric flask (50 ml) and any remaining water was removed by passing it through a tiny funnel filled with 10g of anhydrous sodium sulphate.

Petroleum ether was used to wash the separator funnel. The washings were collected in a volumetric flask and passed through the funnel with sodium sulphate. Petroleum ether was used to make the solution up to the 50 ml level. Utilising a spectrophotometer (UVMS Excellence UV5, Switzerland) to measure the sample's absorbance at 450 nm, the following formula was used to determine the total carotenoid concentration.

$$\text{Total carotenoids (ug/g)} = \frac{\text{Absorbance} \times \text{Total volume} \times 10^{-4}}{\text{Sample weight} \times 2592}$$

### **Determination of beta carotene**

Using acetone, 2g of each sample was extracted to assess the amount of beta-carotene. Using a mortar and pestle, the samples were crushed until the residue lost all colour. After the extract was run through a funnel that was packed with glass wool, 25 millilitres of the extract were placed in a flask with a circular bottom and dried at a temperature of roughly 600 degrees Celsius. To dissolve the beta-carotene, 1 millilitre of petroleum ether was added to the evaporated sample. Column chromatography was then used to elute the solution. A slurry composed of silica gel with a mesh size ranging from 60 to 120 and petroleum ether was used to prepare a 15-cm glass column that was equipped with glass wool at the elution point. Following the settling of the slurry, anhydrous  $\text{Na}_2\text{SO}_4$  was packed into the column, and 1 millilitre of 100% ethanol was added to activate the silica gel and anhydrous  $\text{Na}_2\text{SO}_4$ . Petroleum ether was then used to evaporate the mixture till a volume of 25 ml was obtained. In a UV-VIS spectrophotometer (UVMS Excellence UV5, Switzerland), the eluted absorbance was measured at 450 nm. A standard curve was created by plotting the absorbance of five standard solutions of beta-carotene, ranging in concentration from 0.4 to 2.4 ug/g, against their corresponding concentrations. The absorbance was measured at the same wavelength [18].

### **Determination of total phenol**

Calculation of total phenols was done using method described by [19]. After homogenising the fruits of the chilli peppers in 0.3N HCl in methanol, the mixture was centrifuged at 10,000g for ten minutes. After collecting the supernatant, the pellet was once more extracted using 0.3N HCl in methanol and centrifuged. On a water bath, the supernatants were collected and allowed to evaporate. The resultant residue was dissolved in 5 millilitres of distilled water. 0.5 ml of the Folin-phenol reagent was added to this volume, and it was violently shaken. One millilitre of a 35% sodium carbonate solution was added after three minutes, shaken, and let to stand for one hour. Using a UV-Vis Spectrophotometer (UVMS Excellence UV5, Switzerland), absorbance was measured at 650 nm. Tannic acid concentrations were varied to create a calibration curve that was used to calculate the total quantities of phenolics.

### **Determination of total flavonoids**

The measurement of total flavonoid content was conducted using the [20] method. 80% ethanol-extracted chilli peppers that were stored for a night. Using a UV-Vis Spectrophotometer (UVMS Excellence UV5, Switzerland), 0.5 ml of fruit extract and 0.5 ml of 2%  $\text{AlCl}_3$  in ethanol were added, and the mixture was left to stand for an hour. The number of total flavonoids was calculated using a calibration curve that was made with varying concentrations of quercetin.

### **Determination of total capsaicinoids**

The method by which capsaicinoids were isolated from the chilli pepper samples is detailed by [21]. After heating 0.5 g of chilli peppers for 4 hours at 80°C in 8 ml of acetonitrile, the capsaicinoids were extracted. Throughout the extraction procedure, the suspensions were frequently shaken every 30 minutes. The suspended material settle and cool. The supernatant was utilised for HPLC injections after being filtered through a 0.45 µm membrane filter (Millipore) into a 2 ml glass vial.

Using a pump unit (Hitachi L 6200 A) solvent equipped with an autosampler (Hitachi AS 2000), a fluorescence detector (Hitachi FL), and an integrator (HP 3393 A), liquid chromatography was carried out. At room temperature, the separation was carried out on a column (Phenomenex ODS, 5 µm, 250 x 4.6 mm i.d.).

The HPLC operating conditions that were utilised by [22] were as follows. At a flow rate of 1.2 ml/min, the eluent was a combination of a solution of water, and acetic acid (100: 100: 1). The run lasted 35 minutes in total. Ten µL was the injection volume. Every capsaicinoid was measured at 280 nm with a 0.4 mL/min flow rate. Moreover, U/Vis spectra were captured at a spectrum collection rate of 1.25 scans/s (peak width 0.2min) in the 200–600 nm range.

### **Experimental design and treatments**

A 2 x 3 x 3 factorial experimental design in Randomized Complete Block Design (RCBD) with eighteen (18) treatments (Two (2) levels of chili pepper varieties, three (3) levels of solar drying processes and three (3) levels of types of lining media used in zeer pot storage). The two (2) levels of chili pepper varieties were scotch bonnet and red cayenne, the three (3) levels of solar drying processes were control, blanched and unblanched chili peppers and the three (3) levels of zeer pot lining media were sand, styrofoam and wood shavings. The

experiment was replicated three (3) times with total sample population of fifty – four (54). After three (3) months of storage, samples of stored chili peppers were taken into labelled zip bags and placed in ice chest cooler, transported within four (4) hours to Food Science Laboratory, Kwame Nkrumah University of Science and Technology (KNUST) for Laboratory analyses.

## **RESULTS AND DISCUSSION**

### **Effect of Varieties on **Phytochemical** Characteristics of Chili Pepper Fruits**

**Phytochemical** analyses were carried - out to determine effect of varieties on **phytochemical** characteristics of chili pepper fruits. Table 1 showed the results of effect of varieties on **phytochemical** characteristics of chili pepper fruits. The results indicated that, there were significant difference ( $p < 0.05$ ) in **phytochemical** characteristics of chili pepper fruits.

#### **Total carotenoids of chili pepper fruits**

There was significant effect ( $p < 0.05$ ) on the total carotenoids of chili pepper fruits with red cayenne been the highest (9222.4ug/100g) and the least was recorded in scotch bonnet (4296.1ug/100g)[23].

The increased in total carotenoids in red cayenne could be due to environmental conditions such as light intensity and temperature in which red cayenne was grown [24] and good agronomical practices such as application of fertilizers, irrigation which increased the size of the fruits to cause increased in total carotenoid content in red cayenne [25]. Carotenoids act as antioxidants which protect the human body against diseases by improving the human immune system [26].

#### **Beta carotene of chili pepper fruits**

There was significant difference ( $p < 0.05$ ) in beta carotene which followed **the** same trend as total carotenoids (9222.4ug/100g) with red cayenne recorded significantly ( $p < 0.05$ ) the best beta carotene (934.23ug/100g) but the least beta carotene (429.30ug/100g) been recorded in scotch bonnet[27].

The increased in beta carotene in red cayenne could be attributed to genetic characteristics of red cayenne and the environmental conditions such as light intensity and temperature were suitable for cultivating red cayenne [24]. Beta carotene is a precursor for vitamin A which is good for vision or eye health, healthy skin, mucous membrane, lower the risk of heart diseases, cancer and stronger immune system [6].

#### **Total phenols of chili pepper fruits**

Significantly, there was difference ( $p < 0.05$ ) in total phenols of chili pepper fruits with scotch bonnet been the best in total phenols (301.74mg/100g) whereas red cayenne recorded the least total phenols (229.93mg/100g) of chili peppers[28].

The increased in total phenols in scotch bonnet could be as a result of the cultivar, growing conditions, fruit stage of maturity and postharvest handling of the scotch bonnet fruits [29]. Total phenols protects the human body against diseases such as diabetes, cancer and cardiovascular diseases. It has anti – bacterial and anti – fungal properties which reduce the activities of bacteria and fungi [5].

### Total flavonoids of chili pepper fruits

Total flavonoids had similar trend as total carotenoids (9222.4ug/100g), beta carotene (934.3ug/100g) where red cayenne had significantly ( $p < 0.05$ ) higher total flavonoids (50.04mg/100g) and the least total flavonoids (26.28mg/100g) been recorded in scotch bonnet[30].

High total flavonoids in red cayenne may be due to the genotype of red cayenne and the ripening stage of the red cayenne at which it was harvested [31]. Flavonoids function as detoxifying agent, antimicrobial defensive compound, capable of neutralizing free radicals, protect the human body against cancer, alzheimer, atherosclerosis, cardiovascular diseases such as heart attacks, ulcer, inflammation, osteoporosis, diarrhoea and arthritis [32].

### Total capsaicinoids of chili pepper fruits

Total capsaicinoids were significantly ( $p < 0.05$ ) higher in red cayenne (247.30mg/100g) which also continued similar trend as total carotenoids (9222.4ug/100g), beta carotene (934.23ug/100g) and total flavonoids (50.04mg/100g) whereas scotch bonnet recorded the least total capsaicinoids (174.56mg/100g)[33].

The increased in total capsaicinoids in red cayenne could be attributed to intrinsic genetic characteristics of red cayenne and the environmental conditions where the red cayenne fruits were cultivated [34]. Capsaicinoids are used for flavouring and preserving food due to high pungency as well as used as medicinal purposes. It is used as food additive and also used for treating various pathogenic infections [35]. Again, heat in pepper fruits is added to a wide range of alcoholic beverages to give better taste and flavour [8], [36]. Capsaicin is extracted from chili peppers to use as active ingredient in preparation of tear gas or pepper spray and organic insecticides or pesticides [9].

**Table 1: Effect of varieties on phytochemical characteristics of chili pepper fruits**

Varieties	Carotenoids (ug/100g)	Beta Carotene	Phenol	Flavonoids (mg/100g)	Capsaicinoids
Scotch bonnet	4296.1b	429.30b	301.74a	26.28b	174.56b
Red cayenne	9222.4a	934.23a	229.93b	50.04a	247.30a
HSD (0.05)	75.47	7.76	1.62	0.41	0.76
CV	2.02	2.06	1.10	1.95	0.65

Means with the same letters (s) in the column are not significantly different from each other ( $p > 0.05$ , according to Tukey's HSD)

### **Effect of Solar Drying on Phytochemical Characteristics of Chili Pepper Fruits**

Phytochemical analyses were done to determine the effect of solar drying on phytochemical characteristics of chili pepper fruits. The results were demonstrated in table 2 which suggested that, there were significant effect ( $p < 0.05$ ) on phytochemical characteristics of chili pepper fruits.

#### **Total carotenoids of chili pepper fruits**

There were significant difference ( $p < 0.05$ ) in total carotenoids of chili peppers with controlled chili peppers significantly ( $p < 0.05$ ) had higher total carotenoids (8035.4ug/100g) followed by blanched – solar dried (6647.0ug/100g) chili pepper fruits and the least total carotenoids been unblanched (5595.5ug/100g) chili pepper fruits[37].

The high carotenoids in controlled chili peppers could be due to genotype which determines the specific carotenoids biosynthetic enzymes [38]. Carotenoids act as antioxidants which protect the human body against diseases by improving the human immune system [26].

#### **Beta carotene of chili pepper fruits**

Significantly, the higher beta carotene was noted in controlled (812.34ug/100g) followed by blanched – solar dried (670.78ug/100g) and the least beta carotene was recorded in unblanched – solar dried (562.18ug/100g) of chili pepper fruits[39].

Increased in beta carotene in controlled chili peppers could be due to difference in cultivars, maturity, growing practices, climates and postharvest handling which increased beta carotene [40]. Beta carotene is a precursor of vitamin A which is good for vision or eye health, healthy skin, mucous membrane, lower the risk of heart diseases, cancer prevention and stronger immune system [6].

#### **Total phenols of chili pepper fruits**

The best total phenols was significantly ( $p < 0.05$ ) recorded in unblanched – solar dried (386.64mg/100g) chili pepper samples which was followed by blanched – solar dried (327.11mg/100g) chili pepper samples and the lowest total phenols was determined in controlled chili pepper samples (83.75mg/100g)[41]. The increased in total phenols in unblanched chili peppers might be due to inactive of enzymes at high temperature of drying that caused minimal degradation of total phenols by polyphenol oxidase to increase total phenols in unblanched chili peppers [42]. Total phenols protect the human body against diseases such as diabetes, cancer and cardiovascular diseases. It has anti – bacterial and anti – fungal properties which reduce the activities of bacteria and fungi [5].

#### **Total flavonoids of chili pepper fruits**

There was significant effect ( $p < 0.05$ ) on total flavonoids of chili pepper fruits. Unblanched chili pepper fruits had significantly ( $p < 0.05$ ) highest total flavonoids (51.97mg/100g) which continued same trend as total phenols (386.64mg/100g) followed by blanched (39.60mg/100g) chili pepper fruits while the least was recorded in controlled (22.91mg/100g) of chili pepper samples[43].

The increased in total flavonoids could be attributed to removal of moisture content from the unblanched chili pepper fruits which caused the deactivation of degradative enzymes to increase total flavonoids [44]. Again, the increment in flavonoids in unblanched chili peppers might be due to the release of flavonoids from the cell wall by disrupting hydrogen bonds [45]. Flavonoids function as detoxifying agent, antimicrobial defensive compound, capable of neutralizing free radicals, protect the human body against cancer, alzheimer, antherosclerosis, cardiovascular diseases such as heart attacks, ulcer, inflammation, osteoporosis, diarrhoea and arthritis [46].

### Total capsaicinoids of chili pepper fruits

There was significant difference ( $p < 0.05$ ) in total capsaicinoids where unblanched chili pepper samples had the best total capsaicinoids content (260.72mg/100g) which continued similar trend as total phenols (386.64mg/100g) and total flavonoids (51.97mg/100g) followed by blanched – solar dried (207.67mg/100g) but the least totalcapsaicinoids was recorded in controlled chili pepper samples (164.39mg/100g)[47].

Increased in total capsaicinoid content could be as a result of high temperature been harvested by solar dryer that induced the capsaicinods through their **dissociation** from the cell wall (placenta) and quicker removal of water that increased the concentration of total capsaicinoid content [48]. Capsaicinoidsare used for flavouring and preserving food due to high pungency as well as used as medicinal purposes. It is used as food additive and also used for treating various pathogenic infections [35]. Again, heat in pepper fruits is added to a wide range of alcoholic beverages to give better taste and flavour [8], [36]. Capsaicin is extracted from chili peppers to use as active ingredient in preparation of tear gas or pepper spray and organic insecticides or pesticides [9].

**Table 2: Effect of solar drying on **phytochemical** characteristics of chili pepper fruits**

Solar drying	Carotenoids (ug/100g)	Beta Carotene	Phenol	Flavonoids (mg/100g)	Capsaicinoids
<b>Controlled</b>	8035.4a	812.34a	83.75c	22.91c	164.39c
<b>Blanched</b>	6647.0b	670.78b	327.11b	39.60b	207.67b
<b>Unblanched</b>	5595.5c	562.18c	386.64a	51.97a	260.72a
HSD (0.05)	111.61	11.48	2.39	0.61	1.12
CV	2.02	2.06	1.10	1.95	0.65

Means with the same letters (s) in the columnare not significantly different from each other ( $p > 0.05$ , according to Tukeys HSD)

### Effect of Zeer Pot Refrigeration Lining Media on **Phytochemical** Characteristics of Chili Pepper fruits

**Phytochemical** analyses were conducted to determine the effect of zeer pot refrigeration lining media on **phytochemical** characteristics of chili pepper fruits. Table 3 revealed the results of the **phytochemical** analyses and the results indicated that, there were significant difference ( $p < 0.05$ ) in the phytochemical characteristics of chili pepper fruits.

### Total carotenoids of chili pepper fruits

Significantly, the best total carotenoids were shown in styrofoam lining material of zeer pot refrigeration (8506.1ug/100g) which was followed by sand lining media of zeer pot refrigeration (6327.7ug/100g) and the lowest indicated in wood shavings lining media of zeer pot (5444.0ug/100g)[49].

The increased in total carotenoids in styrofoam lining media might be due to styrofoam containing more air (98%) to aid in easy movement of air for effective evaporative cooling to increase total carotenoid content in chili peppers [50]. Carotenoids act as antioxidants which protect the human body against diseases by improving the human immune system [26].

### **Beta carotene of chili pepper fruits**

The trend continued as similar to total carotenoids where styrofoam lining media of zeer pot refrigeration had significantly ( $p < 0.05$ ) higher beta carotene of chili pepper fruits (861.64ug/100g) followed by sand lining media of zeer pot refrigeration (636.88ug/100g) and the least was recorded in wood shavings lining media of zeer pot refrigeration (546.79ug/100g)[51].

The high beta carotene in styrofoam lining media could be due to styrofoam containing more air (98%) to aid in easy movement of air for effective evaporative cooling to reduce oxidation of beta carotene in order to increase beta carotene content in chili peppers [50]. Beta carotene is a precursor of vitamin A which is good for vision or eye health, healthy skin, mucous membrane, lower the risk of heart diseases, cancer prevention and stronger immune system[6].

### **Total phenols of chili pepper fruits**

Sand used as lining material in zeer pot storage had significantly ( $p < 0.05$ ) the best effect (304.30mg/100g) on total phenols of chili pepper fruits followed by styrofoam (268.55mg/100g) and the least was determined in wood shavings lining media of zeer pot refrigeration (221.61mg/100g)[52].

The high total phenols in chili peppers stored in sand lining media may be attributed to sand been able to warm up and **transfer heat** quickly and large air spaces between **sand** particles **aided** easy movement of air to increase evaporative cooling and declined the temperature of chili peppers stored in the zeer pot to increase total phenols in chili peppers [53]. Total phenols protect the human body against diseases such as diabetics, cancer and cardiovascular diseases. It has anti – bacterial and anti – fungal properties which reduce the activities of bacteria and fungi [5 ].

### **Total flavonoids of chili pepper fruits**

Same trend continued with total phenols where sand used as lining media of zeer pot storage had significantly ( $p < 0.05$ ) higher total flavonoids (49.99mg/100g). There were no significant difference ( $p > 0.05$ ) between styrofoam (32.06mg/100g) and wood shavings (32.43mg/100g) thus they were same. Comparatively, styrofoam had the least effect on total flavonoids (32.06mg/100g) of chili pepper fruits[54].

The high total flavonoids in chili peppers stored in sand lining media may be due to sand been able to conduct heat quickly to **reduce temperature in the inner pots** and high porosity of sand for free circulation of air to increase evaporative cooling to reduce the temperature of chili peppers stored in the zeer pot to increase total flavonoids in chili peppers[53 ].

Flavonoids function as detoxifying agent, antimicrobial defensive compound, capable of neutralizing free radicals, protect the human body against cancer, alzheimer, antherosclerosis, cardiovascular diseases such as heart attacks, ulcer, inflammation, osteoporosis, diarrhoea and arthritis [46].

### Total capsaicinoids of chili pepper fruits

Significantly, higher total capsaicinoids were recorded in sand (242.17mg/100g) used as lining media of zeer pot refrigeration which had similar trend as total phenols (304.30mg/100g) and total flavonoids (49.99mg/100g) while wood shavings used as zeer pot lining media had the least total capsaicinoids (185.72mg/100g)[55 ].

The high total capsaicinoids in chili peppers stored in sand lining media may be attributed to sand been able to warm up and transfer heat quickly to remove heat from the inner pots and large air spaces between particles of sand for easy movement of air to increase evaporative cooling and decreased the temperature of chili peppers kept in the zeer pot to increase total capsaicinoids in chili peppers [53]. Capsaicinoids are used for flavouring and preserving food due to high pungency as well as used as medicinal purposes. It is used as food additive and also used for treating various pathogenic infections [35]. Again, heat in pepper fruits is added to a wide range of alcoholic beverages to give better taste and flavour [8], [36]. Capsaicin is extracted from chili peppers to use as active ingredient in preparation of tear gas or pepper spray and organic insecticides or pesticides [26].

**Table 3: Effect of zeer pot refrigeration lining media on phytochemical characteristics of chili pepper fruits**

Lining media	Carotenoids (ug/100g)	Beta Carotene	Phenol	Flavonoids (mg/100g)	Capsaicinoids
Sand	6327.7b	636.88b	307.34a	49.99a	242.17a
Styrofoam	8506.1a	861.64a	268.55b	32.06b	204.89b
Wood shavings	5444.0c	546.79c	221.61c	32.43b	185.72c
HSD (0.05)	111.61	11.48	2.39	0.61	1.12
CV	2.02	2.06	1.10	1.95	0.65

Means with the same letters (s) in the column are not significantly different from each other ( $p > 0.05$ , according to Tukeys HSD)

### CONCLUSION

Phytochemical analyses were conducted to determine effect of varieties, solar drying processes and zeer pot refrigeration lining media on phytochemical characteristics of chili peppers to reduce postharvest losses of chili peppers in the study area. Results from phytochemical analyses showed that, there were significant effect ( $p < 0.05$ ) on phytochemical characteristics of chili peppers. On effect of varieties on phytochemical properties of chili peppers, red cayenne was able to retain more total carotenoids (9222.4 ug/100g), beta carotene (934.23ug/100g) but scotch bonnet was high in total phenols (301.74mg/100g). Again, total flavonoids (50.04 mg/100g) and total capsaicinoids (247.30mg/100g) were higher in red cayenne. On effect of solar drying processes on phytochemical characteristics of chili peppers; controlled chili peppers were high in total carotenoids (8035.4 ug/100g) and beta carotene (812.334ug/100g). However, total phenols (386.64ug/100g), total flavonoids (51.97ug/100g) and total capsaicinoids (260.72ug/100g) were greater in unblanched chili peppers. On effect of zeer pot refrigeration lining media on phytochemical properties of chili peppers; styrofoam lining media of zeer pot refrigeration was able to preserve total carotenoids (8506.1ug/100g) and beta carotene (861.64ug/100g) better while sand used as lining media of zeer pot was able to conserve total phenols

(307.34mg/100g), total flavonoids (49.99mg/100g) and total capsinoids (242.17mg/100g) well.

## REFERENCES

1. Arnason, A. 2023. Chili peppers 101: Nutrition facts and health effects. <https://www.healthline.com>. 29/03/24
2. FAOSTAT 2024. Database Updates. <https://www.fao.org>. 2/04/24
3. GIRSAL 2023. Reviving Ghana Fresh Chili Pepper Export Industry: Commercial Chili Production Trial Net Housed by GIRSSAL and Partners. <https://www.girsal.com>. 4/04/24
4. Azlan, A., Sultana, S., Huei, C. S., and Razman, M. R. (2022). Antioxidants, Anti-Obesity, Nutritional and other Beneficial Effects of Different Chili Peppers. *Molecules*, 27(3): 898. <https://doi.org/10.3390/molecules27030898>
5. Pratyusha, S. 2022. Phenolic Compounds in Plant Development and Defence: An Overview. <https://www.intechopen.com>. 26/6/23. <https://doi.org/10.5772/intechopen102873>
6. Moreb, N., O'Dwyer, C., Jaiswal, S., Jaiswal, A. A. (2020). Chapter 13 – Pepper. *Nutritional Composition and Antioxidant Properties of Fruits and Vegetables*. 223 – 238. <https://doi.org/10.1016/B978-0-12-812780-3.00013-1>
7. Streit, L. 2019. Is black Pepper Good for You or Bad? Nutrition, Uses and more. <https://www.healthline.com/nutrition>. 6/1/2023
8. Apex Flavours 2020. Pure Red Pepper Extract, Natural (Medium Heat). <https://www.apexflavors.com/Beverage-indus>. 6/2023
9. Jalil, A. A., Taib, A. P., Abdilla, M. T., and Arimbay, I. I. (2023). “Efficacy of Mixture of Neem Leaves (*Azadirachta indica*), Garlic *Allium sativum* L.), and Hot Chili Pepper (*Capsicum annum*) Extract as Organic Insecticides in Okra (*Abelmoschus esculentus*)”. *International Journal of Multidisciplinary Research and Publications (IJMRAP)*, 5 (10). 35 – 38.
10. Kamran, M. (2022). Energy sources and technologies. *Fundamentals of Smart Grid Systems*. Pp 23 – 69. <https://doi.org/10.16/B978.0-323-99560-3.00010-7>

11. Udomkum, P., Romuli, S., Schock, S., Mahayothee, B., Sartas, M., Wossen, T., Njukwe, E., Vanlauwe, B., and Muller, J. (2020). Review of solar dryers for agricultural products in Asia and Africa: An innovation landscape approach. *Journal of Environmental Management* 268,110730. <https://doi.org/10.1016/j.jenvman.2020.110730>
12. Technology Exchange Lab 2019. Zeer Pot Fridge. <https://www.techxlab.org/solution/practical-action.08/03/2020>
13. Megan S. 2020. How to Make AZeer Pot (“Fridge” without Electricity). <https://www.survivalsullivan.com>. 31/03/2020
14. Anand, S., and Barua, M. K. (2022). Modeling the key factors leading to postharvest loss and waste of fruits and vegetables in the agri – fresh produce supply chain. *Computers and Electronics in Agriculture* 198, 106936. <http://doi.org/10.1016/j.compag.2022.106936>.
15. Rubagumya, I., Komakech, A. J., Menya, E., Kizito, S., Zziwa, A., and Kabenge, I. (2023). Postharvest losses of fruits and vegetables along their urban supply chain in eastern Africa: A case study of Uganda towards sustainable management, 48, 105 – 114. <https://doi.org/10.1007/s42853-023-00177-x>
16. Ministry of Food and Agriculture 2024. Dormaa East: Physical and natural environment, location and size. <https://www.mofa.com>. 5/4/24
17. Udensi, J., Loskutova, E., Loughman, J., and Byrne, H. J. (2022). Quantitative Raman Analysis of Carotenoid Protein Complexes in Aqueous Solution. *Molecules*, 4724. <https://doi.10.3390/molecule27154724>
18. Febrianti, A., Aina, G. Q., Farpina, E., Kaltim, P. K. (2022). Determination of vitamin C and Beta – carotene levels in several types of chili (*Capsicum sp*) using UV – Vis Spectrophotometry. *Formosa Journal of Science and Technology (FJST)*, 1 (8), 1129 – 1142. <https://doi.org/10.55927/fjst.v1i8.1949>.
19. Kupina, S., Fields, C., Roman, M. C., and Bruneele, S. L. (2019). Determination of Total Phenolic Content using the Folin – C Assay: Single – Laboratory validation, *First Action* 2017.13. *Journal of AOAC International*, 102 (1). <https://doi.org/10.5740/jaoacint.2017.13>

20. Nur, L., Handayani, V., and Rasyid, F. A., (2021). Spectrophotometric Determination of Total Flavonoid Content in *BiancheaeaSappan* (*Caesalpiniasappan* L.) Leaves. *JurnalFitofarmakaIndonesia*, 8(3) 1- 4. <https://doi.org/10.33096/jffi.v8i3.712>
21. Lalic, M., Solidic, A., Lalic, A., Lalic, Z., and Sertic, M. (2022).Development and Validation of an HPLC Method for Simultaneous Determination of Capsaicinoids and Camphor in Over – the – Counter Medication for Tropical Use. *Molecules*. 27(4). 1261. <https://doi.org/10.3390/molecules27041261>
22. Sharma, S., Kushwaha, R. S., Naman, S., Patil, U. K., Baldi, A. (2021). Optimized Extraction of Oleoresin *Capsicum* and Analytical Method Validation for Capsaicin using HPLC. *Indian J. Nat. Prod.* 35 (1), 38 – 45. <https://www.ijnponline.com>.
23. Duah, S. A., Souza, C. S., Daood, H. G., Pek, Z., Nemenyi, A., and Helyes, L. (2021). Content and response of  $\gamma$  - irradiation before over – ripening of capsaicinoid, carotenoid, and tocopherol in new hybrids of spice chili peppers. *LWT*, 147, 111555. <https://doi.org/10.1016/j.lwt.2021.111555>.
24. Souza, C. S., Daood, H. G., Dua, S. A., Vinogradov, S., Palotas, G., Nemenyi, A., Helyes, L., Pek, Z. (2022). Stability of carotenoids, carotenoid esters, tocopherols and capsaicinoids in new chili pepper hybrids during natural and thermal drying. *LWT*, 163, 113520. <https://doi.org/10.1016/j.lwt.2022.113520>
25. Duan, X., Zou, C., Jiang, Y., Yu, X., and Ye, X. (2023), Effects of Reduced Phosphate Fertilizer and Increased Trichoderma Application on the Growth, Yield, and Quality of Pepper. *Plants* 12, 2998. <https://doi.org/10.3390/plants12162998>
26. Maoka, T. (2020). Carotenoids as natural functional pigments. *Journal of Natural Medicine*, 24. 1 – 16. <https://doi.org/10.1007/s11418-019-01364-x>.
27. Soyong, M., Guevarra, P. R., Mateo, J. M. C., and Galvez, H. F. (2021). Evaluation of tomatoes fruits flesh colour, beta carotene and lycopene content. *International Journal of Agricultural Technology*, 12 (2). 727-736. <https://www.ijat-aatsea.com>.
28. Liu, Y., Chen, Y., Wang, Y., Chen, J., Huang, Y., Yan, Y., Li, L., Li, Z., Ren, Y., and Xiao, Y. (2000). Total phenolics, capsaicinoids, antioxidant activity, and  $\alpha$  – glucosidase inhibitory activity of three varieties of pepper seeds. *International Journal of food Properties*, 23 (1). <https://doi.org/10.1080/10942912.2020.1775646>
29. Dobon – Suarez, A., Gimenez, M. J., Castillo, S., Garcia – Pastor, G., and Zapata, P. J. (2021). Influence of the Phenological Stage and Harvest Date on the Bioactive Compounds

Content of Green Pepper fruits. *Molecules*, 26 (11), 3099.

<https://doi.org/10.3390/molecules26113099>

30. Mi, S., Zhang, X., Wang, Y., Zheng, M., Zhao, J., Gong, H., Wang, X. (2022). Effect of different genotypes on the fruit volatile profiles, flavonoid composition and antioxidant activities of chili peppers. *Food Chemistry*, 374, 131751.

<https://doi.org/10.1016/j.foodchem.2021.131751>

31. Ribes – Moya, A.M., Adalid, A.M., Raigon, M.D., Hellin, P., Fita, A., Rodriguez – Burrezo, A. (2020). Variation in flavonoids in a collection of peppers (*Capsicum sp.*) under organic and conventional cultivation: effect of the genotype, ripening stage and growing system. *J. Sci. Food Agric*, 100 (5). 2208 – 2223. <https://doi.org/10.1002/ssfa.10245>

32. Ullah, A., Munir, S., Badshah, S. L., Khan, N., Ghani, L., Poulson, B. G., Emwas, A., and Jaremko, M. (2020). Important Flavonoids and their Role as a Therapeutic Agent.

*Molecules*, 25 (22), 5243. <https://doi.org/10.3390/molecules25225243>

33. Vazquez – Espinosa, M., Fayos, O., Gonzalez – de – Peredo, A. V., Espada – Bellido, E., Ferreira – Gonzalez, M., Palma, M., Garces – Claver, A., Barbero, G. F. (2020). Content of Capsaicinoids and Capsiate in “Filius” Pepper Varieties as Affected by Ripening. *Plants*, 9 (9), 1222. <https://doi.org/10.3390/plants9091222>

34. Uarrota, V. G., Maraschin, M., De Bairros, A, De F., and Pedreschi, R. (2021). Factors affecting the capsaicinoid profile of hot peppers and biological activity of their non – pungent analogs (Capsinoids) present in sweet peppers. *Critical Review in Food Science and Nutrition*, 61(4). <https://doi.org/10.1080/10408398.2020.1743642>

35. Rezazadeh, A., Hamishehkar, H., Ehsani, A., Ghasempour and Kia, E. M. (2021). Applications of capsaicin in food industry: functionality, utilization and stabilization. *Critical Reviews in Food Science and Nutrition*. <https://doi.org/10.1080/10408398.1997904>

36. Hultquist, M. 2020. Chili Pepper Infused Tequila. <https://www.chiipeppermadness.com>. 13/02/2023

37. Babar, O. A., Arora, V. K., Nema, P. K., Kasara, A. F., and Tarafdar, A. (2021). Effect of PCM assisted flat plate collector solar drying of green chili on retention of bioactive compounds and control of aflatoxins development. *Solar Energy* 229, 102 – 111.

<https://doi.org/10.1016/j.solener.2021.07.077>

38. Tripodi, P., Lo Scalzo, R., and Ficcadenti, N. (2020). Dissection of heterotic, genotypic and environmental factors influencing the variation of yield components and health – related

compounds in chili pepper (*Capsicum annuum*). *Euphytica* 216, 112.

<https://doi.org/10.1007/s10681-020-02648-0>.

39. Romauli, N. D. M., Ambarital, H., Qadry, A., Sihombing, H. S., (2021). Effect of Drying Whole and Half Chili Pods Using a Solar Dryer with CaCl<sub>2</sub> Desiccant on Quality of Powder Chili. *International Journal of Food Science*, 2021. <https://doi.org/10.1155/2021/9731727>

40. Adewoyin, O., Famaye, A., Ipinmoroti, R., Ibidapo, A., and Fayose, F. (2023). Postharvest Handling Methods, Processes and Practices for Pepper. *Capsicum – Current Trends and Perspectives*. <https://doi.org/10.5772/intechopen.106592>

41. Lakshmi, D. V. N., Muthukumar, P., and Nayak, P. K. (2021). Experimental investigations on active solar dryers integrated with thermal storage for drying of black peppers. *Renewable Energy*, 167, 728 – 739. <https://doi.org/10.1016/j.renene.2020.11.144>.

42. Zhai, R., Hu, J., and Jin, M. (2022). Towards efficient enzymatic saccharification of pretreated lignin – derived phenolics and recent trends in mitigation strategies. *Biotechnology Advances*, 61, 108044. <https://doi.org/10.1016/j.biotechadv.2022.108044>

43. Al Maiman, S. A., Albadr, N. A., Almusallam, I. A., Al – Saad, M. J., Alsuliam, S., Osman, M. A., and Hassan, A. B. (2021). The potential of exploiting economical solar dryer in food preservation: Storability, physicochemical properties, and antioxidant capacity of solar –dried tomato (*Solanum lycopersicum*) fruits. *Food*, 10 (4), 734. <https://doi.org/10.3390/food10040734>

44. Quaabou, R., Nabil, B., Ouhammou, M., Idlimam, A., Lamharrar, A., Ennahli, S., Hanine, H., and Mahrouz, M. (2020). Impact of solar drying process on drying kinetics, and on bioactive profile of Moroccan sweet cherry. *Renewable Energy*, 151, 908 – 918. <https://doi.org/10.1016/j.renene.2019.11.078>

45. Cao, D., Liu, Q., Jing, W., Tian, H., Yan, H., Bi, W., Jiang, Y., and Chen, D. D. Y. (2020). *ACS Sustainable Chem. Eng.* 8 (51) 19169 – 19177. <https://doi.org/10.1021/acsuschemeng.0c08146>

46. Ullah, A., Munir, S., Badshah, S. L., Khan, N., Ghani, L., Poulson, B. G., Abdul – Hamid, E., and Jaremko, M. (2020). *Molecules*, 25 (22), 5243. <https://doi.org/10.3390/molecules25225243>

47. Grimaldi, M., Cavazza, A., Pitrollo, O., Xoccali, M., Mondello, L., and Giuffrida, D. (2022). Analytical evaluation of carotenoids, apocarotenoids, capsaicinoids, and phenolics to assess the effect of a protective treatment on chili peppers dried at different temperatures. *European Food Research and Technology* 248, 2339 – 2349. <https://doi.org/10.1007/s00217-022-04049-0>.

48. Boateng, I. D. (2022). Recent processing of fruits and vegetables using emerging thermal and non – thermal technologies. A critical review of their potentialities and limitations on bioactive, structure, and drying performance. *Critical Reviews in Food Science and Nutrition*, 1 – 35. <https://doi.org/10.1080/10408398.2022.2140121>

49. Ngamwonglumlert, L., Devahastin, S., Chiewchan, N., and Raghavan, V. (2020). Plant carotenoids evolution during cultivation, postharvest storage, and food processing: A review.

Comprehensive Reviews in Food Science and Food Safety 19 (4), 1561 – 1604.

<https://doi.org/10.1111/1541-4337.12564>

50.FOAMEX 2021. Properties of Expanded Polystyrene. <https://www.foamex.com>. 24/8/23

51.Balogun, A. A., Ariahu, C. C. Ojo, M. O. (2020). Quality Evaluation of Oranges Stored in Evaporative Coolers. Asian Food Science Journal.

<https://repository.futminna.edu.ng.8080/jspui/handle/1234567889/15756>

52. Kumar, N., Ojha, A., Upadhyay, A., Singh, R., and Kumar, S. (2021). Effect of active chitosan – pullulan composite edible coating enrich with pomegranate peel extract on the storage quality of green bell pepper. LWT, 138, 110435.

<https://doi.org/10.1016/j.lwt.2020.110435>

53. Benneth, A. 2023. Characteristics of Different soils. <https://www.ahdb.org.uk>. 24/08/23

54.Malik, F., Nadeem, M., Ainee, A., Kanwal, R., Sultan, M., Iqbal, A., Mahmoud, S., Alshehry, A., AL – Jumayi, H. A., and Algarni, H. A. (2022). Quality evaluation of lemon cordal stored at different times and microwave heating (Pasteurization). Sustainability, 14 (4), 1953. <https://doi.org/10.3390/su14041953>

55. Garra, A., Alkalai – Tuvia, S., Telerman, A., Paran, I., Fallik, E., and Elmann, A. (2020). Anti – proliferative activities, phytochemical levels and fruit quality of pepper (*Capsicum spp*) following prolonged storage. International Journal of Food Science and Technology, 55 (12), 3574 – 3584.<https://doi.org/10.1111/ijfs.14691>

