

PIGEONPEA PRODUCTIVITY AND SOIL HEALTH AS INFLUENCED BY PHOSPHORUS LEVELS AND AM FUNGI UNDER DIFFERENT PLANTING METHODS

Abstract

A field experiment was conducted at ARS, Bheemarayarnagudi, Karnataka to study the effect of phosphorus graded levels and inoculation of arbuscular mycorrhizal (AM) fungi under different establishment techniques in pigeonpea during *kharif* season of three years (2010-13) in Upper Krishna Project command area of Karnataka. The treatment consists of phosphorus graded levels (0, 12.5, 25, 37.5 and 50kg ha⁻¹) with inoculation VAM fungi under two establishment techniques (Transplanted and dibbled) with randomized block design. Three-year pooled data indicated that, Application of 50 kg P + VAM under transplanting technique recorded significantly higher pigeonpea yield (1510 kg ha⁻¹) and it was on par with application 37.5 kg P + VAM under transplanting method over rest of the treatment. The economics and nutrient uptake also noticed similar trend. It proved that pigeonpea could be successfully transplanted in the UKP command area of Karnataka by applying 37.5 kg of phosphorus ha⁻¹ in conjunction with VAM fungus inoculation. This boosted crop productivity, nutrient availability, and net returns.

Key words: Phosphorus, VAM fungi, transplanting, Pigeonpea

Introduction:

Pigeonpea (*Cajanus cajan* (L.) Millsp.) is one of the major grain legume (pulse) crops of the tropics and subtropics, endowed with several unique characteristics. It finds an important place in the cropping system adopted by small farmers in most of the developing countries. Globally pigeonpea ranks sixth in area and production in comparison to other grain legumes such as beans, peas and chickpea (Jagadale, 2011). Pigeon pea is the second most important pulse crop in India after chickpea in terms of area and production. It is cultivated over an area of 4.43 million hectares, with production accounting for 4.25 million tonnes, and an average productivity of 937 kg ha⁻¹ (Ministry of Agri & FW, 2018). It is largely grown in the northern parts of the state especially in Kalaburgi, which is called “Pulse bowl of Karnataka”. In this district, it occupies an area of 3.5 lakh ha and production of 1.37 lakh tonnes, but productivity is 359 kg ha⁻¹ which is very low compared to the state average productivity of 458 kg ha⁻¹ (Anon., 2008). Abiotic stresses associated to soil fertility and moisture content, particularly in the early stages of crop growth, are mostly responsible for the crop's low productivity (<1 t/ha).

The primary constraint to increased pigeonpea productivity in the subtropics of India is soil moisture-related limitation, which is made worse by the crop's experience with climate aberrations that are unfavorable to its growth and development (Praharaj et al., 2015 and Singh and Srivastava, 2018). The yield of pigeonpea is greatly influenced by a number of factors such as agronomic, pathogenic, entomological, genetic and their interaction with environment. Among the different agronomic practices, inadequate and imbalanced nutrient application is one of the important factors, which is limiting the yield (Praharaj et. al., 2015 and Mahapatra et al., 2020).

As pigeonpea is a legume crop, can meet its nitrogen demand through biological nitrogen fixation. However, the supply of phosphorus becomes crucial for yield maximization.

Phosphorus is an important mineral element for grain legumes as it helps in root development, participates in synthesis of phosphates and phosphor-proteins and takes part in energy fixing and releasing process in plants. Significant response of pigeon pea to phosphate nutrition (Singh and Yadav, 2008). In tropical and subtropical locations, phosphorus is typically the most limiting nutrient for the growth of leguminous crops (Noriharu et. al., 1990). Using Arbuscular Mycorrhizal fungi and plant growth-promoting rhizo-bacteria, often known as bio-fertilizers, can increase the efficiency of phosphorus consumption (Mathimaran, et.al 2020).

In dry farming areas of Northern Karnataka, where the rainfall is not only scanty but also erratic, soil moisture becomes the most limiting factor in production of pigeonpea. To ensure timely sowing due to late onset of monsoon and late release of water to the canal in Upper Krishna Project (UKP) command area of Karnataka, transplanting of pigeonpea seedlings is one of the climate resilient practices to overcome delayed sowing. This technique involves raising seedlings in the polythene bags in the nursery for a period of one month and then transplanting those seedlings in the main field, immediately after soil wetting rains (Anon., 2015). There is hardly any work done on the use of VAM in pigeonpea in this region. The technique of raising the pigeonpea seedlings in poly bags and transplanting is catching up in northern Karnataka region, due to manifold benefits. Use of AM fungi under transplanted condition (in poly bags containing germination media) would be advantageous over applying directly to the field under dibbled condition. Keeping all these aspects in view, the present investigation carried out to know effect of planting methods and phosphorus levels with AM fungi in northeastern dry zone of Karnataka.

Materials and methods

A field experiment was conducted at the College of Agriculture Farm, Bheemarayanagudi, University Agricultural Sciences, Raichur, Karnataka, during the *kharif* season of three years (2010-11 to 2012-13) to evaluate the performance of transplanted pigeonpea (Var. TS-3R) as influenced by mycorrhizal application at varied phosphorus levels. The treatment consisted of 5 phosphorus levels (0, 12.5, 25, 37.5, and 50 Kg P₂O₅/ha) with and without Arbuscular mycorrhizal fungi (AM) and two planting methods (transplanted and dibbling) replicated three times. The station comes under the North-Eastern dry zone of Karnataka, located between 16°43'N and 76°51'E longitude at an elevation of 411.75 meters above MSL, characterized by a dry climate with an average annual rainfall of 774.1 mm, and it lies in the heart of the UKP, a prestigious and largest irrigation command of the country with a projected irrigation potentiality of 10 lakh ha. The soil of the experimental site was medium black soil with soil pH 7.32, EC 0.18dS/m organic carbon 0.17%, available nitrogen 214 kg/ha, available phosphorus 31.5 kg/ha and available potassium is 352 kg/ha.

Two to three bold and healthy seeds of pigeonpea were sown during May in all the years in polythene bags having $\frac{3}{4}$ th of soil, vermicompost and the AM fungus *Glomus fasciculatum* was inoculated at 10 g per polythene bag. Polythene bags were watered regularly. After one month, the seedlings were transplanted in the main field, and seeds were also sown on the day of transplanting by dibbling two to three seeds up to 4 to 5 cm deep in the rows at a 90 x 60 cm spacing. The recommended dose of nitrogen and phosphorus (25: 50 kg NP kg/ha) as per the treatments was applied as basal dose at the time of sowing in the form of urea and SSP. weed growth suppressed by three hand weeding operations to keep the plots free from weeds during the cropping period. To control spotted pod borer (*Maruca vitrata* (G)), Pod borer and Pod fly,

Curacron (2 ml/l) – one spray, Acephate (2 ml/l) – one spray, were sprayed during flowering to grain filling stage.

Soil samples were collected from 0-30 cm depth before sowing and after harvest of the crop from each treatment in all the three replications. The collected samples were air-dried and processed in the laboratory. The soil samples were analyzed for available nitrogen, phosphorus and potassium content. Available soil nitrogen was estimated by alkaline permanganate method as outlined by Subbaiah and Asija (1956). Available phosphorus was determined by Olsen's method as outlined by Jackson (1973) using spectrophotometer. Available potassium was extracted with neutral normal ammonium acetate and the content was estimated by flame photometer (Jackson, 1973). Nitrogen, phosphorus and potassium contents in plant samples of pigeonpea at harvest were estimated by modified micro-kjeldhal method, Vanado-molybdate yellow colour method and flame photometer method, respectively as outlined by Jackson (1973). Nutrient uptake was calculated by using the following formula.

$$\text{Uptake (kg/ha)} = \frac{\text{Nutrient concentration (\%)} \times \text{Biomass (kg/ha)}}{100}$$

The analysis and interpretation of data was done using the Fisher's method of analysis of variance technique as described by Gomez and Gomez (1984). The level of significance used in "F" and "t" test was $p=0.05$. Critical difference values were calculated whenever the "F" was significant.

3. Result and Discussion:

3.1 Pigeonpea yield and growth performance:

The production of economic yield of a crop is an outcome of interaction among the crop, soil, environmental factors and the agronomic manipulations (Mahapatra et.al, 2020). The agronomic practices can modify its surrounding environment (microclimate) to a certain extent and thereby help the crop exploit the available resources more efficiently and achieve higher production (Yadahalli and Palled, 2004). Thus, maximum yields are obtained when optimum conditions are provided for a crop, which, in precise terms, is the object of optimum plant nutrition and a better method of establishment for increasing the productivity and production of the food legumes (Chaudhary and Thakur, 2005 and Jagdale, 2011).

Applying 50 kg P₂O₅/ha inoculating with AM fungi and transplanting of pigeonpea produced significantly higher seed yield (1510 kg/ha) compared to the other treatments (Table 1). However, it is on par to application of 37.5 P₂O₅/ha and inoculating with AM fungi and transplanting method (1478 kg/ha). The dibbling method of pigeon pea without phosphorus and the AM fungi showed significantly lower seed yield (785 kg/ha). Crop being photosensitive, proper time of planting is critical; planting early in the season with the onset of monsoon is more paying as it ensures adequate soil moisture throughout life cycle and accumulate required growing degree days (GDD) besides escaping from pod borer (Chittapur et.al, 2022 and Yadahalli et.al 2016). The early-planted crop's yield advantage was mostly attributable to higher growth, yield characteristics, and physiological traits. Climate and temperature during the periods of growth, development, and maturity also supported it. Higher growth and yield features were the result of better utilization of moisture, nutrients, and light interception (Jayaraja et.al 2022).

The seed yield per plant is governed by yield components like number of pods per plant, 100 seed weight and seed weight per plant. Application of 50 kg P₂O₅/ha inoculating with AM fungi and transplanting method recorded significantly higher seed yield per plant, 100 seed weight and seed weight per plant (83.00 g, 11.30g and 210 pods per plant, respectively) as over other treatments and it was on par transplanting of pigeonpea with application of 37.5kg P₂O₅/ha inoculating AM fungi (81.83g, 11.14 g and 207 pods per plant, respectively) (Table 1). Conspicuously higher yields obtained with application of 50 kg P₂O₅/ha inoculating AM fungi under transplanting technique may be attributed to satisfactory improvement in plant height (158.90cm) and number of branches per plant (17.57) which might have facilitated the crop to undergo increased photosynthetic activity which in turn might helped the crop to accumulate significantly higher dry matter production (113 g/plant) at harvest as compared with rest of the treatments and it was on par with application of 37.5 kg P₂O₅/ha inoculating AM fungi under transplanting technique (156.93cm and 17.37, respectively). All these parameters might have more crops to give significantly higher yield over dibbling method and application of phosphorus at lower graded levels (Table 1). Transplanting is a novel and clever agronomic technique that has recently gained popularity in the Northeastern dry zone (Pavan et al., 2011), the Eastern Gangetic plains of India (Praharaj et al., 2015), and the Northeastern transition zone (Chittapur et al., 2016). It is intended to overcome yield reduction caused by late sowing. Phosphorus plays a pivotal role in the higher yield, by stimulation of root development, energy transformation and metabolic processes in the plants, which turn, resulted in greater translocation of photosynthates towards the sink development (Ade et. al., 2018). Use of Arbuscular Mycorrhiza along with graded levels of phosphorus under transplanting technique enhanced growth and yield of pigeonpea (Jagadale., 2011).

3.2. Root length and Diameter

Mycorrhiza inoculated seedlings recorded comparatively higher root length (26 cm) and diameter (0.92 cm) over the non-mycorrhized seedlings (22 and 0.89 cm respectively), which may be due to optimum supply of phosphorus to the seedlings in the juvenile stage of growth compared to the seedlings grown without mycorrhizal inoculation (Fig.2). Use of Arbuscular Mycorrhiza increases the production of growth promoters, increases tolerance to diseases by plant and improves the synergistic interaction with Beneficial N-fixer and P-solubilizes microorganism in chickpea (Joshi et. al., 2021). This is achieved by increasing plant P absorption—a crucial step for biological nitrogen fixation by utilizing VAM. Additionally, mycorrhiza strengthens the plant's resilience to various diseases caused by soil, enhances its water intake, making it more drought-tolerant, and protects it from weed species like striga. Furthermore, mycorrhizal fungi produce glomalin, a glycoprotein that binds soil particles and improves soil structure and additionally, by binding heavy metals, it increases plants' tolerance to their detrimental effects (Njira et.al. 2017). Arbuscular Mycorrhizal Fungi (AMF) play important roles in agroecosystems, including the involvement of the extraradical mycelium in providing soil aggregation (Rillig, 2004).

3.3. Nutrient Uptake (kg ha^{-1})

Nutrient uptake of pigeonpea differed significantly due to graded phosphorus levels and mycorrhizal inoculation. Higher uptake of nitrogen, phosphorus and potassium ($116.33 \text{ kg ha}^{-1}$) was noticed with application of 50 kg P + AM fungi under transplanting technique and which was on par with 37.50 kg P + AM fungi with Transplanting method ($115.33 \text{ kg ha}^{-1}$) over other treatments (Table 2). A significantly higher uptake of nitrogen, phosphorus, and potassium was recorded in transplanted conditions when compared to the dibbled method. It was mainly due to

the strong and deep root system of pigeonpea transplanted rather than the dibbled method. Such plants had robust, well-developed root systems, and it's because mycorrhizal treatment enhanced the amount of available N, P, and K in the rhizosphere, as well as its solubility. The results agree with a study that shows an increase in mycorrhizal colonization with fertilization by N or P in nutrient limited soils, varying with species, with incidences of *Glomus spp.* increasing in relatively fertile soils (Treseder and Allen, 2002). Similarly, VAM fungal inoculation promotes rhizosphere colonization, leads to in higher nutrient uptake and improves BNF, crop growth, and yields in legume-based cropping systems (Njira, et al., 2017).

3.5 Available nutrients after the harvest of crop

Significantly higher availability of nitrogen (198kg ha^{-1}) was registered with application of $50\text{ kg P} + \text{VAM}$ under transplanting method which was on par with all other treatments except without 0 kg P - Dibbling (170kg ha^{-1}) and $0\text{ kg P} + \text{AM}$ with dibbling method (172kg ha^{-1}) as compared to other treatments (Table. 2). Significantly higher availability of phosphorus (60 kg ha^{-1}) was noticed in 50 kg -P with dibbling method compared to rest of the treatments. It might be due to the higher amount of phosphorus applied and lower amount of phosphorus uptake by the crop due to weak root system, P- fixation and absence of mycorrhiza in soil. Availability of potassium did not differ significantly due to graded levels of phosphorus application.

Higher availability of nitrogen and phosphorus was registered in transplanted pigeonpea when compared to dibbled one. This was because, in transplanted conditions crops have produced better root system, a greater number of root nodules per plant and higher number of root hairs as observed during harvest (Chittapur et al., 2016). It helped to leave more residual nitrogen in soil after crop harvest, which was the product of biological nitrogen fixation (Njira, et al., 2017 and Joshi et. al., 2021).

5.5 Economics

Application of 50 kg P + AM fungi under transplanting technique has recorded the higher Gross returns (Rs.58, 545 ha⁻¹) and net returns (Rs. 37,449 ha⁻¹) over the rest of the treatment. However, it was on par with the application of 37.5 kg P + AMfungi under transplanting method. This is mainly due to high seed yield being produced. The higher cost of cultivation recorded in with application of 50kg P + AM fungi under transplanting method (Rs. 21, 096 ha⁻¹) as compared to other treatments. This clearly indicates that the cost of cultivation in transplanted pigeonpea was higher than in dibbling, which was due to extra cost incurred on raising seedlings, polythene bags, nursery maintenance and more labour requirement for transplanting. Transplanted pigeonpea has recorded comparatively higher gross returns compared to dibbling (Table. 1), it was due to higher seed yield over the dibbled method. Similar results were analyzed by Priyanka et al. (2013) in transplanted and dibbled establishment techniques in pigeonpea. Application of 50 kg P + AMfungi under transplanting method recorded significantly higher B:C(2.88) as compared to other treatments, and it was on par with application of 50 kg P + AM fungi under transplanting method (2.87). This is mainly due to transplanted pigeonpea has given significantly higher net returns under same set of management compared to dibbling, which was because of significantly higher seed yield and cost saved on phosphorus supply. Similarly higher remunerative returns were observed in transplanted pigeonpea over dibbled method by Yadahalli et.al (2016), Jayaraja et al (2022) and Priyanka et al. (2013).

Conclusion: From the results of this agronomic investigation, it may conclude that the application of 37.5 kg phosphorus ha⁻¹ with mycorrhizal inoculation was found to be beneficial for pigeonpea because of higher seed yield and net returns. Which will save phosphorus to the

tune of 12.5 kg ha⁻¹. Among the different methods of crop establishment transplanting was found to be better because of higher seed yield and net returns. This method is very well suited to Upper Krishna Project command area of Karnataka because of unusual/late onset of monsoon.

REFERENCE:

1. Ade, U. K. Dambale, A. S. and Jadhav, D. B., 2018, Growth and Yield of pigeonpea (*Cajanus cajan* L. Millsp) as influenced by phosphorus and biofertilizer. *International Journal of Current Microbiology and Applied Science*, 6: 1427-1434.
2. Anonymous, 2008, All India area, production and yield of total pulses 2007-08. www.google.com/Department of Agriculture.
3. Anonymous, 2015, Redgram transplanting technology, KVK, Bidar, Karnataka, India. http://sri.ciifad.cornell.edu/aboutsri/othercrops/otherSCI/InKarnSCI_Redgram15.pdf.
4. Chaudhary, S.K. and Thakur S. K., 2005. Productivity of pigeonpea (*Cajanus cajan*)-based intercrops. *Indian Journal of Agricultural Sciences* 75: 496-497.
5. Chittapur B M, 2016, Climate smart agriculture: Lessons learnt, technological advances made and research priorities in semi-arid tropics. In, *Climate Smart Agriculture: Status and Strategies*, (Ed.) B. M. Chittapur, A. S. Halepyati, , M. R. Umesh, and B. K. Desai, Published by University of Agricultural Sciences, Raichur, Karnataka: 30-39.
6. Chittapur, B M., Anand N. and Potdar, M. P., 2022, Improved production technologies for harnessing genetic potential in semi-arid tropics of India. *Journal of Farm Science*. 35(1): 1-13.

7. Gomez, K.A. and Gomez, A. A.1984. Statistical procedure for Agriculture for Agriculture Research. *A Willey- Inter science Publication*, John Willey and Sons. New York, Pp. 108-137.
8. Jackson, M.L., 1973, Soil Chemical Analysis. Prentice Hall of India, Pvt. Ltd., New Delhi, pp. 478.
9. Jagadale, CL., 2011, Studies on phosphorus management in transplanted pigeonpea [*Cajanus cajan* (L.) millsp.] through VAM. *M.Sc(Agri) Thesis*, University of Agricultural Sciences, Raichur,
10. Jayaraja, P., Umesh M.R., Ajayakumar, M. Y., Ananda, N. and Chavan S. S., 2022, Climate Resilient Practices on Performance and Economics of Transplanted Pigeonpea and Sunflower Intercropping in Vertisol of Northern Karnataka. *Indian Journal of Ecology*, 49(3): 727-732
11. Joshi, G.H., Thalkar, M. G., Lanje, S. N., Pagore, G. K. and Kadam, A. D., 2021, Effect of PSB, VAM and phosphorus levels on plant height, shoot and root growth in chickpea (*Cicer arietinum* L.). *The Pharma Innovation Journal*, 10(4): 550-553.
12. K. K.Treseder and M. F.Allen, 2002, Direct nitrogen and phosphorus limitation of arbuscular mycorrhizal fungi: a model and field test. *New Phytologist*, 155(3), pp. 507–515
13. Mahapatra, S. S., Bhushan, C., Anil Shukla, Singh, V. K. and Navneet Pareek, 2020. Feasibility and Possibility of Transplanted Pigeonpea (*Cajanus cajan* L.) under different spacing and nutrient management practices. *International Journal of Current Microbiology and Applied Science*, 9(12): 3053-3067.

14. Mathimaran N., Jegan S., Thimmegowda M.N., Prabavathy V.R., Yuvaraj P., Kathiravan R, Sivakumar M.N., Manjunatha B.N., Bhavitha N.C., Sathish A., Shashidhar G.C., Bagyaraj D.J., Ashok E.G., Singh Dt., Kahmen A., Boller T., and Mäder P., 2020, Intercropping transplanted pigeonpea with finger millet: Arbuscular mycorrhizal fungi and plant growth promoting rhizobacteria boost yield while reducing fertilizer Input. *Frontiers in sustainable Food systems*. 4:88.
15. Ministry of Agriculture and Cooperation. 2018. Agricultural Statistics at a Glance, agricoop.nic.in.
16. Njira K, O W., Ernest Semu, Mrema, J. P. and Nalivata, 2017, Pigeonpea and cowpea-based cropping system improve vesicular arbuscular mycorrhizal fungal colonisation of subsequent maize on the alfisols in Central Malawi. *International Journal of Microbiology*, 7:9.
17. Noriharu A. E., Joji Arihara, Kensuke Okada, Teruhiko Yoshihara, Chris Johansen., 1990., Phosphorus uptake by pigeon pea and its role in cropping systems of the Indian subcontinent. *Science*, 248(4954):477-480.
18. Pavan, A. S., Nagalikar, V. P., Halepyati, A.S. and Pujari, B.T., 2009, Effect of planting on the yield, yield components and economics of transplanted pigeonpea. *Karnataka Journal of Agriculture Science* 22: 433-434.
19. Praharaj, C. S., Kumar, n., Ummed Singh, Singh, S. S. and Jagdish Singh, 2015, Transplanting in pigeonpea - A contingency measure for realizing higher productivity in Eastern Plains of India. *Journal of Food Legumes* 28(1): 34-39, 2015.

20. Priyanka, K., Patil, S. S., Hiremath, G. M., Joshi, A. T. and Kulkarni, S. A. 2013. Comparative analysis of transplanted and dibbled method of redgram cultivation in Bidar district of Karnataka. *Karnataka Journal of Agricultural Sciences*. 26(2): 238-242.
21. Rillig, M. C., 2004, Arbuscular mycorrhizae, glomalin, and soil aggregation, *CandianJournal of Soil Science*, 355 – 363.
22. Singh, R.S. and Yadav, M.K, 2008. Effect of phosphorus and biofertilizers on growth, yield and nutrient uptake of long duration pigeon pea under rainfed condition. *J. of Food Legumes* 21(1): 46-48.
23. Subbaiah, B. Y. and Asija, G. L., 1956, A rapid procedure for the estimation of available nitrogen in soils. *Curr. Sci.*, 25: 259-260.
24. Yadahalli, G. S., and Palled, Y. B., 2004, Response of black gram (*Vigna mungo* L.) genotypes to dates of sowing and phosphorus levels in Northern Transitional tract of Karnataka. *Karnataka Journal of Agricultural Sciences*, 17(2):215-219.
25. Yadahalli, G. S., Vidyavathi G.Y., Doddamani, B.M. and K.A. Hiremath, 2016, Phosphorus management in transplanted pigeon pea through VAM fungi. Published in *4th International Agronomy Congress extended summaries*, Vol 2, 741-742, during Nov. 22–26, New Delhi, India.







Fig:1. Pigeonpea yield (kg/ha) as influenced by planting methods and phosphorus levels with AM fungi inoculation from 2010-11 to 2012-13.

 Symbol indicates standard error bars at 5%)

UNDER




Fig 1. Root length and diameter (cm) of pigeonpea seedlings at the time of transplanting as influenced by mycorrhizal inoculation.

UNDER PEER REVIEW

Table: 1 Effect of phosphorus levels with AM fungi inoculation and planting methods on growth parameters, yield parameters and economics of pigeonpea(pooled over 3 year's data).

Treatment Details	Plant height (cm)	No of branches per plant	Dry weight (g/plant)	No of pods per plant	Seed weight (g/plant)	100 Seed wt. (g)	Pigeonpea yield (kg/ha)	Gross returns (Rs./ha)	Cost of cultivation (Rs./ha)	Net returns (Rs./ha)	B:C ratio
Transplanted with 0 kg P ₂ O ₅ /ha	132.07	14.97	141.30	169.57	67.03	9.13	997	38569	16818	21751	2.30
Transplanted with 12.5 kg P ₂ O ₅ /ha	137.57	15.17	148.10	177.77	70.27	9.57	1071	41473	17548	23925	2.38
Transplanted with 25 kg P ₂ O ₅ /ha	138.87	15.60	149.77	179.70	71.03	9.67	1132	43896	18230	25666	2.43
Transplanted with 37.5 kg P ₂ O ₅ /ha	147.07	16.43	160.10	192.10	75.93	10.34	1231	47744	19087	28657	2.54
Transplanted with 50 kg P ₂ O ₅ /ha	150.67	16.87	164.97	197.53	78.10	10.63	1375	53333	20112	33221	2.71
Transplanted with 0 kg P ₂ O ₅ /ha + VAM	140.03	15.77	151.23	181.30	71.73	9.77	1079	41860	17554	24306	2.40
Transplanted with 12.5 kg P ₂ O ₅ /ha +VAM	142.23	15.93	154.00	184.77	73.03	9.94	1151	44584	18234	26350	2.47
Transplanted with 25 kg P ₂ O ₅ /ha + VAM	149.97	16.70	163.77	196.50	77.67	10.55	1258	48764	19131	29632	2.59
Transplanted with 37.50 kg P ₂ O ₅ /ha + VAM	156.93	17.37	172.53	207.03	81.83	11.14	1478	57478	20708	36770	2.87
Transplanted with 50 kg P ₂ O ₅ /ha + VAM	158.90	17.57	175.00	210.00	83.00	11.30	1510	58545	21096	37449	2.88
Dibbling 0 kg P ₂ O ₅ /ha	109.70	12.77	113.00	135.57	53.60	7.30	785	30398	12004	18395	2.56
Dibbling 0 kg P ₂ O ₅ /ha + VAM	114.70	13.20	118.53	142.23	56.23	7.66	850	32926	12608	20318	2.65
Dibbling 50 kg P ₂ O ₅ /ha	115.77	13.03	120.67	144.77	57.23	7.79	928	35984	14044	21941	2.61
Dibbling 50 kg P ₂ O ₅ /ha + VAM	119.00	13.70	124.80	150.67	59.63	8.12	1008	39144	14750	24394	2.72
SEM ±	5.89	0.55	7.40	8.34	3.51	0.47	47	2131	-	2131	0.11
CD(0.05)	17.11	1.59	21.49	24.23	10.18	1.37	134	6193	-	6193	0.28

Table 2. Total uptake of nutrients and Available nutrients (kg/ha) in soil after harvest of pigeon pea as influenced by phosphorus levels with AM fungi inoculation and planting methods (Pooled over 3 year).

Details	Nutrients Uptake (kg/ha)			Available nutrients (kg/ha)		
	Nitrogen	Phosphorus	Potash	Nitrogen	Phosphorus	Potash
Transplanted with 0 kg P ₂ O ₅ /ha	114	27	32	186	33	380
Transplanted with 12.5 kg P ₂ O ₅ /ha	116	28	33	187	37	376
Transplanted with 25 kg P ₂ O ₅ /ha	117	31	34	189	48	376
Transplanted with 37.5 kg P ₂ O ₅ /ha	120	34	38	193	51	377
Transplanted with 50 kg P ₂ O ₅ /ha	122	39	39	197	53	375
Transplanted with 0 kg P ₂ O ₅ /ha + VAM	117	36	33	186	30	376
Transplanted with 12.5 kg P ₂ O ₅ /ha + VAM	121	42	34	188	36	376
Transplanted with 25 kg P ₂ O ₅ /ha + VAM	125	46	37	190	46	375
Transplanted with 37.50 kg P ₂ O ₅ /ha + VAM	128	51	42	195	38	374
Transplanted with 50 kg P ₂ O ₅ /ha + VAM	129	53	43	198	40	373
Dibbling 0 kg P ₂ O ₅ /ha	103	26	29	170	36	383
Dibbling 0 kg P ₂ O ₅ /ha + VAM	104	36	32	172	39	382
Dibbling 50 kg P ₂ O ₅ /ha	108	37	36	189	60	380
Dibbling 50 kg P ₂ O ₅ /ha + VAM	110	47	38	192	41	378
SEM ±	3.3	1.3	1.2	5.44	1.29	6.15
CD(0.05)	9.4	3.7	3.4	15.81	3.74	NS