

Original Research Article

Duration of illness and resilience as determinants of life satisfaction amongst patients with eye pathologies

ABSTRACT

Aim: To determine how duration of disease and resilience as determinants correlate with life satisfaction amongst eye pathology patients

Methods: This was a prospective cross-sectional survey amongst eye patients at Guinness Eye Centre, Onitsha, South East, Nigeria using pretested interviewer-administered structured questionnaire. Information such as the socio-demographics, duration of disease and questions aimed at assessing the resilience and life satisfaction scores of these patients were collected.

Results: A total of 137 patients in the age range 15 to 88 years, and with disease duration ranging from 1 to 23 years were enrolled. Gender and age of the participants were controlled as covariate. There was significant correlation between duration of eye pathology diagnosis and life satisfaction $r = -0.22$, $p = 0.01$, and resilience and life satisfaction $r = 0.30$, $p = 0.01$. In other words as the duration of eye pathology diagnosis increases, the more likely that the patient will report lesser life satisfaction. In contrast, patients who report higher resilience were more likely to also report higher life satisfaction.

Conclusion: Life satisfaction amongst eye pathology patients is significantly determined by duration of eye disease and resilience.

INTRODUCTION

Life Satisfaction is the central aspect of human welfare. It is an ultimate goal and every human being strives to achieve this goal throughout their life. According to Ed Diener it refers to an individual's personal judgement of wellbeing and quality of life based on his or her own chosen criteria.¹

It is subjective, but measurable. Life satisfaction is based on the variables that an individual finds personally important in their own life. A person's life satisfaction will not be determined based on a factor that he/she does not actually find personally meaningful.² Quality of Life (QOL), a term related to life satisfaction, is a measure of well-being. It is associated with living conditions like the amount and quality of food, the state of one's health, and the quality of one's shelter.³ However, the difference between this related variable and life satisfaction is that life satisfaction is subjective and more inherently emotional. Someone who is homeless or terminally ill may well have a higher life satisfaction than a wealthy person in good health, because they may place importance on a very different set of variables than those involved in quality of life. Amongst the factors listed by experts income and self assessed level of health are the most important predictors of Life satisfaction.⁴

Resilience, although with several definitions, fundamentally, refers to positive adaptation, or the ability to maintain or regain mental health, despite experiencing adversity.⁵ More broadly, it is "the protective factors and processes or mechanisms that contribute to a good outcome, despite experiences with stressors shown to carry significant risk for developing psychopathology."⁶ The various definitions together acknowledge 2 points: various factors and systems contribute as an interactive dynamic process that increases resilience relative to adversity; and resilience may be context and time specific and may not be present across all life domains.⁷ Resilience is multidimensional and can be affected by multiple factors. There are four main themes of factors affecting resilience: (1) the influence of individual factors (e.g. individual traits, temperament, having a higher purpose, being self-determined), (2) environmental and organizational factors (e.g. workplace culture), (3) approaches that an individual takes when interacting with her/his professional circumstances (e.g. professional shielding and self-reflection), and (4) effective educational interventions (e.g. resilience workshops).⁸

Ocular health has a unique place in the overall health and functioning of an individual. Reasonably, most people dread having eye problems in their lifetime let alone getting blind. A survey of 2044 Americans showed that Loss of sight was rated the worst health affliction for an individual and 87.5% believed that for a person to be said to be in overall good health, good vision must be present. Blindness was rated worse than, or equal to other serious losses like loss

of limb, memory, speech and hearing. The participants' major reason was degradation of quality of life associated with loss of vision.⁹

A person lives his life, and sees life from personal perspective. In other words, an individual ultimately determines the metrics of his satisfaction. A low QOL may co-exist with good life satisfaction. In this light, a life satisfaction study of persons with eye pathologies might reveal the situation from the patient's point of view. A number of authors have carried out vision related QOL studies worldwide with outstanding results, however, as life satisfaction is the ultimate goal the authors have set out to investigate the effects of eye pathologies on the life satisfaction of adults in Nigeria; looking out for any moderating roles of age and gender. The results of this study will be useful in the wholesome management of the patients with eye pathology.

MATERIALS AND METHOD

This is a cross-sectional descriptive study in which patients diagnosed with eye pathologies were recruited consecutively at the Guinness Eye Centre, Onitsha, South east Nigeria.

A sample size of participants with visual impairment was determined using the formula

$$N = z^2pq/d^2$$

Where,

N = Minimum sample size.

Z = The standard normal deviate, usually set at 1.96 corresponding to 95% confidence interval.

p = Assumed prevalence taken from the estimated prevalence of visual impairment (mild, moderate and severe) which is 10.1% (0.101).[10]

$$q = 1.0 - p (1.0 - 0.101) = 0.899.$$

d = Precision level acceptable = 5% (0.05).

This gives a minimum sample size of approximately 140.

Informed verbal consent was obtained from each respondent. Oral informed consent was considered since the data was collected by using an interview administered structured questionnaire (see appendix) and also there was no invasive examination procedure conducted on the patients for the sake of this research. Patient information was obtained with no identifier and confidentiality was maintained.

Data was analyzed using the Statistical Package for the Social Science version 22 (IBM Software Group, Chicago, IL, USA). Means of continuous variables was compared using Student's *t*-test and ANOVA. Statistical significance was set at $P < 0.05$ for all analyses.

RESULTS

One hundred and thirty seven patients with ocular pathologies were interviewed. The age range was from 15 to 88years while the duration of illness ranged from 1 to 23years.

Table 1. Descriptive (mean and standard deviation) and correlation of the study variables

		mean	SD	1	2	3	4	5
1	Age	54.61	16.47	1				
2	Gender	-	-	-	1			
3	Life satisfaction	24.25	5.85	.15	.03	1		
4	Duration of Illness	4.32	4.22	.14	-.13	-.22**	1	
5	Resilience	32.29	3.78	.02	.02	.30**	-.10	1

** $p=.01$

The study explores the duration of eye pathology diagnosis, and resilience as determinants of Life satisfaction among patients living with some eye pathologies. Gender and age of the participant were controlled as covariate.

The initial correlations showed that there was significant correlation between duration of eye pathology diagnosis and life satisfaction $r= -.22$, $p=. ; 01$ resilience and life satisfaction $r= .30$, $p=.01$ (see Table 1).

Table 2: Multiple regression for duration of illness and resilience as determinants of life satisfaction

	R^2	$Df1(df2)$	F	$Beta$	T
Model 1	.01	2(134)	1.78		
Age				.16	1.85
Gender				.05	.56
Model 2	.13	2(132)	6.17**		

Duration of Illness				-.21**	-2.56
TResilience				.28**	3.47

**p=.01

Further analysis to test the study hypotheses utilizing hierarchical multiple regression shows that duration of eye pathology diagnosis, and resilience contributed to 13.2% ($R^2 = .132$) of life satisfaction among the participants. The ANOVA summary shows that the model was significant at $F(4,136)=6.18$, $P=.01$ (see Table 2).

Specifically, the beta coefficient shows that duration of eye pathology diagnosis negatively and significantly predicted Life satisfaction. $B=-.21$, $P=.01$. In other words as the duration of eye pathology diagnosis increases, the more likely that the patient will report lesser life satisfaction.

The second hypothesis was accepted, resilience positively and significantly predicted life satisfaction among patients living with some eye pathologies at $\beta = .28$, $P=.01$ (see Table 2). In other words patients who report higher resilience were more likely to also report higher life satisfaction.

DISCUSSION

There are several quality of life (QoL) studies on patients with eye pathologies, however, none was found dwelling on the aspect of life satisfaction and its determinants in patients with ocular diagnoses. Nonetheless, few studies have suggested that 'life satisfaction assessment could be considered a measure of quality of life'.^{11,12} Yildirim et al reported that a significant positive correlation was found between life satisfaction and QOL among nursing students in Turkey.¹² This correlation is true for the various domains of QoL. In corroborating the choice of life satisfaction assessment over QoL as in this study, Diener and co-workers asserted that the judgement of how satisfied people are with the present state of affairs is based on a comparison with a standard which each individual sets for him or herself. They say that it is not externally imposed. For this reason researchers believe that self-report is the most direct and most accurate way to measure life satisfaction.^{13,14}

From this study it was found that as the duration of the eye diseases since diagnosis increased the patients were less satisfied with life ($r = -0.22$, $p = 0.01$). This significant negative correlation may be accounted for by the fact that most long lasting eye diseases tend to cause more burden of visual impairment hence more impact on life satisfaction. In fact, the Blue Mountain Study on the impact of bilateral visual impairment on health-related quality of life (HRQOL) reported that the impact appeared to be directly related to the severity of visual impairment but not to the underlying eye condition. The impact of visual impairment was comparable with that of major medical conditions and affected mental more than physical domains.¹⁵ A different report from study of Parkinson disease patients indicated that increasing disease duration correlated with lower HRQOL when assessed as a global construct. However, when subscales were evaluated, difficulties with bodily discomfort and cognitive complaints were comparable in individuals in the 1-5 years and 6-10 years duration groups.¹⁶ Furthermore, in contrast to the finding of this study an assessment of the satisfaction with life in a group of psoriasis patients in Poland revealed a surprise when the duration of the disease was considered. A longer duration of the disease was associated with a higher satisfaction with life. This phenomenon was most evident in women. In contrast, in men the disease lasting longer than 40 years was reflected by a marked decrease in the satisfaction levels, despite men having an increased satisfaction with life proportionally to the disease duration up to this cut off value.¹⁷ Some patients who experience disease remission following improved modern care over time may expectedly gain in their satisfaction with life. It is possible this aspect of positive correlation between duration of disease and life satisfaction resulted from build-up of resilience to the disease over considerable number of years.

This study found that the more resilient the eye pathology patients were the better their satisfaction with life ($r = 0.30$, $p = 0.01$). This was consistent with most other previous studies.

Peng et al in China, while studying glaucoma patients, reported a significant positive correlation between resilience and QoL with sleep disturbance as a mediating factor(18). They found that this relationship between resilience and QoL is mitigated in patients with sleep disturbance. Accordingly, glaucoma patients with severe sleep disturbance have lowered QoL. In a similar study, resilience was found to mediate the relationship between social support and quality of life in primary glaucoma patients. (19) This finding suggested that increasing resilience and social support can improve the quality of life of primary glaucoma patients in clinical practice. Studies done in other chronic medical conditions showed similar trend: inflammatory bowel disease,(20) recurrent coronary artery disease,(21) Parkinson's disease,(22) rheumatoid arthritis,(23) epilepsy,(24) diabetes(25) and colon cancer (26).This study shows that in combination resilience and duration of disease significantly accounted for 13.6% of total life satisfaction of the patients although further analysis was not done to determine the other contributors to life satisfaction. This may include level of income/employment, educational level, social support and environmental factors.(12) Similarly, a study has shown that the extent of resilience exhibited may be determined by the type of coping strategy adopted by the patients especially the strategies of "positive refocusing", "positive reevaluation" and "positive thinking" .(27) In contrast, few studies show that the physical component of quality of life is also related to resilience, but in a negative way.(27, 28) Patients undergoing oncological treatment could be suffering from numerous physical severe symptoms that directly influence their physical condition, such as bodily pain, fatigue, sleep difficulties, gastrointestinal or endocrine disorders, among others. (28) This relation could be derived from the influence of bodily pain, which is also negatively and significantly associated with resilience.

This study is limited by the non- categorization of the specific eye pathologies in relation to the findings as most eye pathologies do not carry the same weight of visual morbidity. In addition, the comparison of QoL studies with this present study on life satisfaction may pose some analytical errors. However, the findings of significant correlations of disease duration and resilience with life satisfaction will open a new horizon of further studies on the validity of interchangeability of usage of both tools in assessing patients.

CONCLUSION

The life satisfaction of eye disease patients is negatively affected as the duration of the disease prolongs. However, patients with enhanced resilience have improved satisfaction with life. Psychotherapies aimed at enhancing resilience will positively impact on life satisfaction of eye pathology patients.

CONSENT

The authors report no conflict of interest concerning this work.

REFERENCE

1. Diener, Ed (1984). "Subjective well-being". *Psychological Bulletin*. **95** (3): 542–575. [doi:10.1037/0033-2909.95.3.542](https://doi.org/10.1037/0033-2909.95.3.542). [PMID 6399758](https://pubmed.ncbi.nlm.nih.gov/6399758/)
2. Ackerman CE. PositivePsychology.com <https://positivepsychology.com/life-satisfaction/> Accessed on 16/08/2021.
3. Veenhoven, R. (1996). The study of life satisfaction. In W. E. Saris, R. Veenhoven, A. C. Scherpenzeel, & B. Bunting (Eds.) *A Comparative Study of Satisfaction with Life in Europe* (pp. 11-48). Budapest, Hungary: Eötvös University Press.

4. UKEssays. Definition Of Life Satisfaction Psychology Essay [Internet]. November 2018. [Accessed 15 August 2021]; Available from: <https://www.ukessays.com/essays/psychology/definition-of-life-satisfaction-psychology-essay.php?vref=1>
5. Wald J, Taylor S, Asmundson GJG, et al. Literature review of concepts: psychological resiliency. Toronto (ON): Defence R&D Canada; 2006.
6. Hjemdal O, Friborg O, Stiles TC, et al. A new scale for adolescent resilience: grasping the central protective resources behind healthy development. *Measurement and Evaluation in Counseling and Development*. 2006;39:84–96.
7. Herrman H, Stewart DE, Diaz-Granados N, Berger EL, Jackson B, Yuen T. What is Resilience? *The Canadian Journal of Psychiatry*. 2011;56(5):258-265.
8. Huey CWT, Palaganas JC. What are the factors affecting resilience in health professionals? A synthesis of systematic reviews. *Med Teach*. 2020 May;42(5):550-560.
9. Scott AW, Bressler NM, Ffolkes S, Wittenborn JS, Jorkasky J. Public Attitudes About Eye and Vision Health. *AMA Ophthalmol*. 2016; 134(10) : 1111-1118.
10. Fatima Kyari, Murthy V. S. Gudlavalleti, Selvaraj Sivsubramaniam, Clare E. Gilbert, Mohammed M. Abdull, Gabriel Entekume, Allen Foster, the Nigeria National Blindness and Visual Impairment Study Group; Prevalence of Blindness and Visual Impairment in Nigeria: The National Blindness and Visual Impairment Survey. *Invest. Ophthalmol. Vis. Sci*. 2009;50(5):2033-2039. doi: <https://doi.org/10.1167/iovs.08-3133>

11. Ferrans, C. E., & Powers, M. J. (1992). Psychometric assessment of the Quality of Life Index. *Research in nursing & health*, 15(1), 29-38.
12. Yildirim, Y., Kilic, S. P., & Akyol, A. D. (2013). Relationship between life satisfaction and quality of life in Turkish nursing school students. *Nursing & health sciences*, 15(4), 415-422.
13. Diener, E., Emmons, R.A., Larsen, R.J. & Griffin, S. (1985). The Satisfaction with Life Scale. *Journal of Personality Assessment* 49 (1), 71-75.
14. Worell, J. (2002). *Encyclopedia of Women and Gender: Sex Similarities and Differences and the Impact of Society on Gender*. San Diego: Academic Press.
15. E. Chia, P. Mitchell, W. Smith, E. Rochtchina, J. Wang, Impact of Bilateral Visual Impairment on Health-Related Quality of Life: The Blue Mountains Eye Study [Investigative Ophthalmology & Visual Science](#) 2003 45(1):71-6
16. Benge JF, Kekecs Z, Encarnacion E, Ainslie M, Herff C, Elkins G, Herath P. Duration of disease does not equally influence all aspects of quality of life in Parkinson's disease. *J Clin Neurosci*. 2016 Jun;28:102-6.
17. Jankowiak B, Sekmistrz S, Kowalewska B, Niczyporuk W, Krajewska-Kułak E. Satisfaction with life in a group of psoriasis patients. *Postepy Dermatol Alergol* 2013; 30: 85–90.
18. Peng, Q., Qu, B., Sznajder, K. K., Chen, Q., Fu, J., He, S., & Yang, X. (2022). Exploring the Association Between Resilience and Quality of Life Among Glaucoma Patients: Sleep Disturbance as a Mediating Factor. *Frontiers in Medicine*, 9, [842864]. <https://doi.org/10.3389/fmed.2022.842864>
19. Wang Y, Zhao Y, Xie S, Wang X, Chen Q, Xia X. Resilience Mediates the Relationship Between Social Support and Quality of Life in Patients With Primary Glaucoma. *Front Psychiatry*. 2019 Jan 31;10:22.
20. Dai W, Zeng Y, Liang E, Zhou Q, Zhang L, Peng J. The actuality of resilience, social support and quality of life among patients with inflammatory bowel disease in China. *Nurs Open*. (2021) 2021:946. doi: 10.1002/nop2.946

21. Jo E, Kim SR, Kim HY. Predictive model for quality of life in patients with recurrent coronary artery disease. *Eur J Cardiovasc Nurs.* (2019) 18:501–11. doi: 10.1177/1474515119847544
22. 36. Kwok JYY, Choi EPH, Chau PH, Wong JYH, Fong DYT, Auyeung M. Effects of spiritual resilience on psychological distress and health-related quality of life in Chinese people with Parkinson's disease. *Qual Life Res.* (2020) 29:3065–73. doi: 10.1007/s11136-020-02562-x
23. Liu, L., Xu, X., Xu, N. *et al.* Disease activity, resilience and health-related quality of life in Chinese patients with rheumatoid arthritis: a multi-center, cross-sectional study. *Health Qual Life Outcomes* **15**, 149 (2017). <https://doi.org/10.1186/s12955-017-0725-6>
24. Glória Maria Almeida Souza Tedrus, Jose Mario Limongi, Jose Victor RoveroniZuntini, Resilience, quality of life, and clinical aspects of patients with epilepsy, *Epilepsy& Behavior* Volume 103, Part A, 2020,
25. Nawaz A, Malik JA, Batool A. Relationship between resilience and quality of life in diabetics. *J Coll Physicians Surg Pak.* 2014 Sep;24(9):670-5. PMID: 25233974.
26. Franjić D, Babić D, Marijanović I, Martinac M. Association between Resilience and Quality of Life in Patients with Colon Cancer. *PsychiatrDanub.* 2021 Dec;33(Suppl 13):297-303. PMID: 35150498.
27. Macía P, Barranco M, Gorbeña S, Iraurgi I. Expression of resilience, coping and quality of life in people with cancer. *PLoS One.* 2020 Jul 29;15(7):e0236572. doi: 10.1371/journal.pone.0236572. PMID: 32726344; PMCID: PMC7390401.
28. Deimling GT, Albitz C, Monnin K, RenzhoferPappada HT, Nalepa E, Boehm ML, Mitchell C. Personality and psychological distress among older adult, long-term cancer survivors. *J Psychosoc Oncol.* 2017 Jan-Feb;35(1):17-31. doi: 10.1080/07347332.2016.1225145. Epub 2016 Aug 19. PMID: 27541961.



UNDER PEER REVIEW