

UNIFORM ESTIMATES ON LENGTH OF PROGRAMS AND COMPUTING ALGORITHMIC COMPLEXITIES FOR QUANTITATIVE INFORMATION MEASURES

Abstract

Shannon entropy and Kolmogorov complexity are two conceptually distinct information metrics since the latter is based on probability distributions while the former is based on program size. All recursive probability distributions, however, are known to have an expected Up to a constant that solely depends on the distribution, the Kolmogorov complexity value is equal to its Shannon entropy. We investigate if a comparable correlation exists between Renyi and Havrda-Charvat Entropy entropies order α , indicating that it is consistent solely with Renyi and Havrda-Charvat entropies of order 1.

Kolmogorov noted that the characteristics of Shannon entropy and algorithmic complexity are comparable. We examine a single facet of this resemblance. Specifically, linear inequalities that hold true for Shannon entropy and for Kolmogorov complexity. As it happens, the following are true: (1) all linear inequalities that hold true for Shannon entropy and vice versa for Kolmogorov complexity; (2) all linear inequalities that hold true for ranks of finite subsets of linear spaces for Shannon entropy; and (3) the reverse is untrue.

Keywords and phrases: Kolmogorov complexity, Shannon entropy, non-Shannon entropy etc.

1. Introduction

The size of the smallest program that can construct an object x is used to assess its exact information content. This is known as Kolmogorov complexity, or $C(x)$. Assigning a probability of $2^{-C(x)}$ to any given string x , it constructs a probability distribution over Σ^* naturally. μ stands for the universal probability distribution, which is the name given to this probability distribution introduced in [3].

A random variable X 's average uncertainty is expressed as its Shannon entropy, or $S(X)$. In average, it is the least amount of bits needed to express x , the random variable X 's output. Shannon entropy and Kolmogorov complexity are conceptually distinct from one another since the former is dependent on program length. Three separate generalizations of Shannon entropy are of importance to us and that was provided in [6], [8], [9], [12] and [16].

Shannon Entropy : Let X be a finite or infinitely countable set and let X be a random variable taking values in X with distribution P introduced in [12]. The Shannon entropy of the random variable X is

$$S(X) = - \sum_{x \in \chi} p(x) \log p(x)$$

Renyi Entropy :Let X be a finite or infinitely countable set and let X be a random variable taking values in X with distribution P introduced in [13] , [14] and let $\alpha \neq 1$ be a non -negative real number. The Renyi 'entropy of order α of the random variable X is defined as:

$$R_{\alpha}(X) = \frac{1}{1-\alpha} \log \left(\sum_{x \in X} p(x)^{\alpha} \right)$$

Havrda and Charvat Entropy: Let X be a finite or infinitely countable set and Let X be a random variable taking values in X with distribution P introduced in [15],[17] and let $\alpha \neq 1$ be a non negative real number .TheHarveda –Charvat entropy of order α of the random variable X is defined as:

$$HC_{\alpha}(X) = \frac{1}{1-\alpha} \sum_{x \in X} p(x)^{\alpha} \frac{1}{1-\alpha}$$

It is easy to prove that $\lim_{\alpha \rightarrow 1} R_{\alpha}(X) = \lim_{\alpha \rightarrow 1} HC_{\alpha}(X) = S(X)$.

Note that we also use the notation $R_{\alpha}(P)$, $HC_{\alpha}(P)$ and $S(P)$ to denote the Renyi,Harveda-Charvat and Shannon entropies of distribution P , respectively.

2. Our Results:

Kolmogorov Complexity and Non-Shannon entropy

Considering the conceptual distinctions between Shannon entropy and Kolmogorov complexity, it's noteworthy to note that under certain loose constraints on the string distribution, They share a relationship. Up to a constant term that solely depends on the distribution, the Shannon entropy value actually matches the expected value of Kolmogorov complexity that we provided in [4],[5],[9] and [16].

Proposition: There are recursive probability distribution P such that:

- i) $\sum_x P(x) K(x) - R_{\alpha}(P) > K(P)$, where $\alpha > 1$
- ii) $\sum_x P(x) K(x) - R_{\alpha}(P) < 0$, where $\alpha < 1$
- iii) $\sum_x P(x) K(x) - HC_{\alpha}(P) > K(P)$, where $\alpha > 1$
- iv) $\sum_x P(x) K(x) - HC_{\alpha}(P) < 0$, where $\alpha < 1$

Proof: consider the following probability distribution:

$$D_n(x) = \begin{cases} \frac{1}{2} & \text{if } x = 0^n \\ 2^{-n} & \text{if } x = 1x', x' \in \{0,1\}^{n-1} \\ 0 & \text{otherwise} \end{cases}$$

i) First we observe that

$$\begin{aligned} S(D_n) &= - \sum_x D_n(x) \log D_n(x) \\ &= - \left(\frac{1}{2} \log \frac{1}{2} + \frac{1}{2^n} 2^{n-1} \log \frac{1}{2^n} \right) \end{aligned}$$

$$= -\left(-\frac{1}{2} - \frac{1}{2^n} 2^{n-1} n\right)$$

$$= \frac{n+1}{2}$$

By theorem we have

Let $P(x)$ be a recursive probability distribution ,then:

$$0 \leq \sum_x D_n(x) K(x) - S(D_n) \leq K(D_n) "$$

Which implies that

$$0 \leq \sum_x D_n(x) K(x) - S(D_n)$$

$$\sum_x D_n(x) K(x) \geq S(D_n)$$

$$\sum_x D_n(x) K(x) \geq \frac{n+1}{2}$$

On the other hand,by definition

$$R_\alpha(D_n) = \frac{1}{1-\alpha} \log \left(\sum_{x \in \mathcal{X}} D_n(x)^\alpha \right)$$

$$= \frac{1}{1-\alpha} \log \left(\frac{1}{2^n} + 2^{n-1} \times \frac{1}{2^{n\alpha}} \right)$$

$$= \frac{1}{1-\alpha} (\log(2^{(n-1)\alpha} + 2^{n-1}) - n\alpha)$$

To prove that $\sum_x D_n(x) K(x) - R_\alpha(D_n) > K(D_n)$, it is sufficiently to prove that

$$\lim_n \left(\sum_x D_n(x) K(x) - R_\alpha(D_n) > K(D_n) \right) > 0$$

$$\text{i.e.} \lim_n \left(\frac{n+1}{2} - \frac{1}{1-\alpha} (\log(2^{(n-1)\alpha} + 2^{n-1}) - n\alpha) - c \log n \right) > 0$$

$$\text{But, } \lim_n \left(\frac{n+1}{2} - \frac{\log((2^{(n-1)\alpha} + 2^{n-1}))}{1-\alpha} + \frac{n\alpha}{1-\alpha} - c \log n \right)$$

$$\geq \lim_n \left(\frac{n+1}{2} - \frac{\log((2^{(n-1)\alpha}))}{\alpha-1} + \frac{n\alpha}{\alpha-1} - c \log n \right)$$

$$= \lim_n \left(\frac{n+1}{2} + \frac{(n-1)\alpha}{\alpha-1} - \frac{n\alpha}{\alpha-1} - c \log n \right)$$

$$= \lim_n \left(\frac{n+1}{2} - \frac{\alpha}{\alpha-1} - c \log n \right)$$

$= +\infty$

ii) To prove this item we use the other inequality

$$\sum_x D_n(x) K(x) - S(D_n) \leq K(D_n)$$

Which implies that

$$\sum_x D_n(x) K(x) \leq K(D_n) + S(D_n)$$

$$\leq \frac{n+1}{2} + c \log n$$

So, $\sum_x D_n(x) K(x) - R_\alpha(D_n)$

$$\leq \frac{n+1}{2} + c \log n - \frac{1}{1-\alpha} (\log(2^{(n-1)\alpha} + 2^{n-1}) - n\alpha)$$

$$\leq \frac{n+1}{2} + c \log n - \frac{\log(2^{n-1})}{1-\alpha} + \frac{n\alpha}{1-\alpha}$$

$$= \frac{n+1}{2} + c \log n - \frac{1}{1-\alpha} (n-1 - n\alpha)$$

$$= -\frac{n}{2} + \frac{1}{2} + c \log n + \frac{1}{1-\alpha}$$

Thus, taking n sufficiently large.

iii) The Havrda and Charvat Entropy of order α of distribution P_n is :

$$HC_\alpha(D_n) = \frac{1}{1-\alpha} \sum_{x \in X} D_n(x)^\alpha - 1$$

$$= \frac{1}{1-\alpha} \left(\frac{1}{2^\alpha} + 2^{n-1} + \frac{1}{2^{n\alpha}} \right) - 1$$

$$= \frac{1}{1-\alpha} (2^{(n-1)\alpha} + 2^{(n-1)}) - 1$$

$$= \frac{2^{(n-1)\alpha}}{1-\alpha} + \frac{2^{(n-1)}}{1-\alpha} - 1$$

Using the inequality, we get

$$\sum_x D_n(x) K(x) - HC_\alpha(D_n) = \sum_x D_n(x) K(x) - \frac{2^{(n-1)\alpha}}{1-\alpha} - \frac{2^{(n-1)}}{1-\alpha} + 1$$

$$\geq \frac{n+1}{2} - \frac{2^{(n-1)\alpha}}{1-\alpha} - \frac{2^{(n-1)}}{1-\alpha} + 1$$

$$\geq \frac{n+3}{2} - \frac{2^{(n-1)\alpha}}{1-\alpha} - \frac{2^{(n-1)}}{1-\alpha}$$

For n sufficiently large

$$\begin{aligned}
&\geq \frac{n+3}{2} - \frac{2^{(n-1)\alpha}}{1-\alpha} - \frac{2^{(n-1)}}{1-\alpha} \\
&> c \log n \\
&= kD_n
\end{aligned}$$

iv) Using the inequality, we get:

$$\sum_x D_n(x) K(x) - HC_\alpha(D_n) \leq \frac{n+1}{2} + \log n - \frac{2^{(n-1)\alpha}}{1-\alpha} - \frac{2^{(n-1)}}{1-\alpha} + 1$$

Since n is sufficiently large, thus we conclude that:

$$\begin{aligned}
&< \frac{n+1}{2} + \log n - \frac{2^{(n-1)\alpha}}{1-\alpha} - \frac{2^{(n-1)}}{1-\alpha} + 1 \\
&< 0
\end{aligned}$$

Uniform Continuity of the Shannon and Non Shannon entropies

i) Shannon entropy

In order to prove the uniform continuity of the Shannon entropy, we need some technical lemmas and theorems.

Theorem: let X and Y be two probability distribution over $\sum n$, let S(X) and S(Y) be the Shannon entropy of distribution X and Y respectively. then,

$$\forall \varepsilon > 0, \exists \delta > 0, \forall X, Y: \max |X(u) - Y(u)| \leq \delta \Rightarrow |S(X) - S(Y)| \leq \varepsilon$$

Proof. By lemma the function $u \log u$ is uniformly continuous in $[0,1]$, we have that:

$$\forall \gamma > 0, \exists \beta > 0, \forall u, v: \max |u - v| \leq \beta \Rightarrow |u \log u - v \log v| \leq \gamma$$

So,

$$\begin{aligned}
|S(X) - S(Y)| &= \left| -\sum_u X(u) \log X(u) + \sum_u Y(u) \log Y(u) \right| \\
&= \left| \sum_u X(u) \log X(u) - \sum_u Y(u) \log Y(u) \right| \\
&\leq \sum_u |X(u) \log X(u) - Y(u) \log Y(u)| \\
&\leq \sum_u \gamma
\end{aligned}$$

$$= 2^n \gamma$$

It is sufficient to consider $\gamma = \frac{\varepsilon}{2^n}$ and consider $\delta = \beta$

ii) Non-Shannon Entropy

a) Renyi Entropy

Let X and Y be two probability distribution over $\sum n$, let $R_\alpha(X)$ and $R_\alpha(Y)$ be the Renyi entropy of distribution X and Y respectively. then ,

$$\forall \epsilon > 0, \exists \delta > 0, \forall X, Y: \max |X(u) - Y(u)| \leq \delta \Rightarrow |R_\alpha(X) - R_\alpha(Y)| \leq \epsilon$$

Proof: By Lemma, the function x^α is uniformly continuous in $[0,1]$. Thus,

$$\forall \gamma > 0, \exists \beta > 0, \forall x, y: \max |x - y| \leq \beta \Rightarrow |x^\alpha - y^\alpha| \leq \gamma \quad (2.1)$$

We have to show that

$$\text{If } \max |X(u) - Y(u)| \leq \delta \Rightarrow |R_\alpha(X) - R_\alpha(Y)| \leq \epsilon$$

$$\begin{aligned} |R_\alpha(X) - R_\alpha(Y)| &= \left| \frac{1}{1-\alpha} \log \left(\sum_{x \in \mathcal{X}} X(x)^\alpha \right) - \frac{1}{1-\alpha} \log \left(\sum_{x \in \mathcal{X}} Y(x)^\alpha \right) \right| \\ &= \frac{1}{1-\alpha} \left| \log \left(\sum_{x \in \mathcal{X}} X(x)^\alpha \right) - \log \left(\sum_{x \in \mathcal{X}} Y(x)^\alpha \right) \right| \end{aligned}$$

$$\leq \frac{2}{1-\alpha} \left| \sum_{x \in \mathcal{X}} X(x)^\alpha - \sum_{x \in \mathcal{X}} Y(x)^\alpha \right|,$$

by Lemma we have" $|\log a - \log b| \leq 2|a - b|$ "

$$\leq \frac{2}{1-\alpha} \sum_x |X(x)^\alpha - Y(x)^\alpha|$$

From (2.1) we consider $\gamma = \frac{\epsilon(1-\alpha)}{2^{n+1}}$

$$\begin{aligned} &\leq \frac{2}{1-\alpha} \sum_x \frac{\epsilon(1-\alpha)}{2^{n+1}} \\ &\leq \frac{2^{n+1} \epsilon(1-\alpha)}{1-\alpha} \frac{1}{2^{n+1}} \\ &\leq \epsilon \end{aligned}$$

(b) Havrda and Charvat Entropy

Let X and Y be two probability distribution over $\sum n$, let $HC_\alpha(X)$ and $HC_\alpha(Y)$ be the **Havrda-Charvat** entropy of distribution X and Y respectively. then ,

$$\forall \epsilon > 0, \exists \delta > 0, \forall X, Y: \max |X(u) - Y(u)| \leq \delta \Rightarrow |HC_\alpha(X) - HC_\alpha(Y)| \leq \epsilon$$

Proof: We have,

$$\begin{aligned}
|HC_\alpha(X) - HC_\alpha(Y)| &= \left| \frac{1}{1-\alpha} \sum_{x \in X} X(x)^\alpha - \frac{1}{1-\alpha} - \frac{1}{1-\alpha} \sum_{x \in X} Y(x)^\alpha + \frac{1}{1-\alpha} \right| \\
&= \left| \frac{1}{1-\alpha} \sum_{x \in X} X(x)^\alpha - \frac{1}{1-\alpha} \sum_{x \in X} Y(x)^\alpha \right| \\
&= \frac{1}{1-\alpha} \left| \sum_{x \in X} X(x)^\alpha - \sum_{x \in X} Y(x)^\alpha \right| \\
&\leq \frac{1}{1-\alpha} \sum_{x \in X} |X(x)^\alpha - Y(x)^\alpha| \\
&\leq \frac{1}{1-\alpha} \gamma
\end{aligned}$$

We consider $\gamma = \epsilon (1 - \alpha) \leq \epsilon$

Conclusion

Among the three entropies that we have examined, we have demonstrated that only the Shannon entropy meets the conditions for the relationship with the expected value of Kolmogorov complexity stated in by presenting a probability distribution for which the relationship breaks down for certain values of α of the Havrda - Charvat and Renyi κ entropies. This relationship also applies for the Shannon entropy in a time-bounded manner, assuming that the cumulative probability distribution can be computed in the given amount of time. We investigated the convergence of this distribution under the Havrda- Charvat and Renyi' entropies , which are two generalizations of the Shannon entropy. Additionally, we demonstrated the uniform continuity of the three entropies taken into consideration in this work.

References:

- 1) **Cachin, C. (1997):**“Entropy measures and unconditional security in cryptography”, 1st ed.; In ETH Series in Information Security and Cryptography; Hartung-Gorre Verlag: Konstanz, Germany.
- 2) **Chaitin, G.J. (1966):**“On the length of programs for computing finite binary sequences”. J. ACM , 13, 547–569.
- 3) **Cover, T.; Gacs, P.; Gray, M. (1989):**“Kolmogorov’s contributions to information theory and algorithmic complexity”. Ann, 17, 840–865.
- 4) **Grunwald, P.; Vitanyi, P. (2004):**“Shannon information and Kolmogorov complexity”.
- 5) **Hammer, D. and A. Shen, A.,(1998):**“A strange application of Kolmogorov complexity, Theory Comput”
- 6) **Kolmogorov, A.N. (1965):**“Three approaches to the quantitative definition of information”,Probl. Inform. Transm. , 1, 1–17.
- 7) **Kolmogorov, N. (1965):**“Three approaches to the quantitative definition of information, Problems Inform”, Transmission 1, 17.

- 8) **Kolmogorov, N. (1968):** “Logical basis for information theory and probability theory” IEEE Trans. Inform.Theory14 , 662664.
- 9) **Li, M. and Vitanyi, P. (1997):** “An Introduction to Kolmogorov Complexity and Its Applications”, second edition, Springer-Verlag, Berlin, New York.
- 10) **Li, M.; Vitanyi, P. (2008):** “An Introduction to Kolmogorov Complexity and Its Applications”, 3rd ed.; Springer Publishing Company, Incorporated: New York, NY, USA.
- 11) **Ramshaw, J. (1995):** “Thermodynamic stability conditions for the Tsallis and Renyi entropies”, Phys. Lett. A, 198, 119 – 121.
- 12) **Shannon, C.E. (1948):**“A mathematical theory of communication”, Bell Syst. Tech. J. , 27 379–423,623–656.
- 13) **Solomonoff, R. A. (1964):** “Formal theory of inductive inference”; Part I. Inform. Contr., 7, 1–22.
- 14) **Solovay, R.M.,(1975):** “Draft of a paper (or series of papers) on Chaitin’s work”. IBM Thomas J. Watson Research Center, Yorktown Heights, New York, NY, USA. Unpublished manuscript.
- 15) **Tadaki, K. (2008):**“The Tsallis entropy and the Shannon entropy of a universal probability”. In Proceedings of the International Symposium on Information Theory, Toronto, Canada, 6–11; abs/0805.0154, pp. 2111–2115.
- 16) **Uspensky, V.A. and A. Shen,A.(1996):**“Relation between varieties of Kolmogorov complexities”, Math. Systems Theory 29, 271292.
- 17) **Zhang, Z.,(2007):**“Uniform estimates on the Tsallis entropies”. Lett. Math. Phys, 80, 171–181.