

Studies on effect of Weed Management on Growth and Yield of Okra (*Abelmoschus esculentus* L.) cv. Arka Anamika

ABSTRACT

The present investigation entitled “**Studies on effect of Weed Management on Growth and Yield of Okra (*Abelmoschus esculentus* L.) cv. Arka Anamika.**” was conducted during Kharif season of 2022-23 at student instructional farm of the S.C. Bose P.G. College, Kahli, Terwa, Gaushganj, Hardoi (U.P.). The experiment was conducted in Randomized Block Design with 6 treatments combinations and 3 replications. Viz. T1- Weed check (Control), T2- Weed free check (3 hand weeding), T3-Pre emergence application of pendimethalin @ 6ml/l, T4-Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding, T5-Pre emergence application of pendimethalin @ 6ml/l. + quazalofop ethyl 40-50 gm./h. at 20 DAS, T6- Post emergence application of metribuzin @ 525 g/h. at 20 DAS.

Keywords-Treatments, Parameters, Pre-emergence, Pendimethalin, Attributes, Replications, Management, Experiments.

INTRODUCTION:

Okra (*Abelmoschus esculentus* (L). Moench) belongs to family malvaceae, known as Bhendi or Lady's finger, is one of the most important traditional Vegetable grown in tropical and sub-tropical parts of the world. Okra is also one of the most nutritious vegetables which contains on an average 1.9g of protein, 0.2g fat, 6.4g carbohydrate, 0.7g minerals and 1.2g fibres per 100g of edible portion. (Gopalan et al., 1989).

In the year 2022-23, in India the area of vegetable cultivation was 11.34(000'ha) and production 212.55 (000'MT). India is the largest producer of okra with an area of 513.6(000' ha) and production of 6170 (000'MT) with the productivity of 11.33 MT, respectively (NHB, 2018-19). Percent share of vegetable production is 59.83% of total horticulture Production.

Its tender green fruits are used as a vegetable and are generally marketed as fresh state, but sometimes as canned or in dehydrated form. Mature fruits and stem containing crude fibre are used in the paper industry. In some places the plants are also soaked in water and the result and solution is used as a clarifier in the manufacture of jaggery.

The daily balance diet for adult human in our condition should constitute 125g leafy vegetables, 75g other vegetables and 100g of root and tuber vegetables along with 475g cereals, 80g pulses, 92g fruits, 200ml milk, 40g sugar or jaggary (Choudhary, 1992).

Hence, the importance of vegetables in a daily diet increasing day by day in many developing countries. National Agricultural Development Programme is focused on the need to increase the production of vegetables in order to increase their proportion in the daily diet (Choudhary, 1992).

MATERIAL & METHODS-

The experiment was carried out at student instructional farm of the S.C. Bose P.G. College, Kahli, Terwa, Gaushganj, Hardoi (U.P.). Total Six treatments are taken and divided in three blocks. Effect of different treatments can be seen easily on various growth and yield parameters. Treatment T₁ has Weed check (Control), In treatment T₂, Weedfree check (3 hand weeding), In treatment T₃, Pre emergence application of pendimethalin @ 6ml/l. are occur, In treatment T₄, Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding are occur, In treatment T₅, pre emergence application of pendimethalin @ 6ml/l. + quizalofop ethyl 40-50 gm/h. at 20 DAS are occur and last one, Treatment T₆, post emergence application of metribuzin @ 525 g/h. at 20 DAS are occur.

RESULT AND DISCUSSION-

Effect of different treatments on different days on Growth parameters & Yield attributes parameters are as under following-

Plant height (cm.)-

The effect of weed control was found significant on plant The highest plant height was recorded under T₄- Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding (87.88 cm) and the lowest plant height were recorded in T₁- weed check (control) is (81.22).

No. of fruits per plant-

Weed management treatments registered significant effect on number of fruits per plant. T₄: Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding recorded highest number of fruit per plant (24.99) In weed (control) plot the number fruit /plant were found lowest T₁(19.99).

Length of Pod-

The lowest fruit length was observed under treatment T₁ (weed check) is (10.33 cm.). and highest fruit length was observed in T₄ (Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding) are (14.66 cm.).

No. of branches-

Number of branches per plant was significantly influenced by various treatments of weed management and Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding (T₄) recorded maximum number of branches (5.16). and minimum no. of branches are in (T₁)- Weed Control (2.83).

No. of Nodes-

Number of nodes per main stem was significantly influenced by various treatments of weed management and Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding (T₄) recorded maximum nodes per main stem (17.16) and minimum no. of nodes are recorded in (T₁) Weed check (Control) are (13.66).

Fresh weight of fruit (gm.)-

Treatment T₁ [Weed check (Control)] has lowest fresh weight of fruit is (10.33 gm.) and highest are recorded in T₄ (Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding) are (13.99).

Fruit yield/ plant(gm.)-

The fruit yield per plant ranged from 184.16 to 233.49. Among the treatments maximum fruit yield per plant was observed in T₄ (233.49) and Minimum fruit yield per plant are in T₁ (184.16).

Total Pod yield (t/ha.)-

Among different weed control treatments T₄ (Pre emergence application of pendimethalin @ 6ml/l.+ one hand weeding) recorded maximum okra fruit yield (15.99 t/ha) and being at par with treatment T₃ and T₄. Significantly the lowest okra green fruit yield (12.49 t/ha) was recorded under treatment weedy check (T₁).

Table1. showing different yield attributing characters and growth parameters.

Treatments	Plant height (cm.)	No. of fruits/ plant	Length of Pod (cm.)	No. of branches
T1- Weed check (Control)	81.220	19.995	10.330	2.830
T2-Weed free check (3 hand weeding)	83.993	22.995	13.160	4.160
T3-Pre emergence application of	85.553	23.495	12.830	3.830


pendimethalin @ 6ml/l				
T4-Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding	87.883	24.995	14.665	5.160
T5-Pre emergence application of pendimethalin @ 6ml/l. + quazalofop ethyl 40-50 gm./h. at 20 DAS	84.443	20.995	12.660	3.500
T6- Post emergence application of metribuzin @ 525 g/h. at 20 DAS.	85.997	22.495	12.165	3.160
C.D.(5%)	1.837	1.919	1.698	0.002
S.E.(m)	0.576	0.516	0.457	0.000
S.E.(d)	0.814	0.730	0.646	0.001

Table2. showing different yield attributing characters and growth parameters.


Treatments	No. of Nodes	Fresh weight of Fruit (gm)	Fruit yield/ plant (gm.)	Total Pod yield (t/ha.)
T1- Weed check (Control)	13.665	10.330	184.165	12.495
T2-Weed free check (3 hand weeding)	16.165	12.995	198.160	14.495
T3-Pre emergence application of	15.165	12.160	206.830	13.495

pendimethalin @ 6ml/l				
T4-Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding	17.160	13.995	233.495	15.995
T5-Pre emergence application of pendimethalin @ 6ml/l. + quazalofop ethyl 40-50 gm./h. at 20 DAS	14.660	12.995	219.665	13.330
T6- Post emergence application of metribuzin @ 525 g/h. at 20 DAS.	14.995	12.330	224.660	14.995
C.D.(5%)	1.157	1.745	6.146	1.197
S.E.(m)	0.311	0.470	1.654	0.322
S.E.(d)	0.440	0.664	2.339	0.456

Graph 1. showing different yield attributing character and growth parameters-



Graph 2. showing different yield attributing character and growth parameters-



Conclusion:-

Plant height affected by the treatments had manifested the maximum height of the plant(87.88cm) at all the stages of observation indicated that the height of plant mainly

governed by weed free check followed by pre- emergence application of pendimethalin @ 6ml/l+one hand weeding and lowest (81.22cm) in weedy check(control).

Maximum number of fruits per plant clearly indicated that in (T₄) Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding (24.99) followed by Pre emergence application of pendimethalin @ 6ml/l. produced number of fruits (23.49) and lowest (19.99) in weed check in control.

Length of Pod clearly indicated that (T₄) Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding produced highest length of fruit (14.66 cm.) followed by (T₂) Weed free check (3 hand weeding). It was also clear that the lowest in T₁ (10.33 cm.) fruit length per plant recorded in weedy check (control).

No. of Branches clearly indicated that (T₄) Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding produced highest no. of branches (5.16) followed by (T₂) Weed free check (3 hand weeding). It was also clear that the lowest in T₁ (2.83) no. of branches per plant recorded in weedy check (control).

No. of Nodes clearly indicated that (T₄) Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding produced highest no. of nodes (17.16) followed by (T₂) Weed free check (3 hand weeding). It was also clear that the lowest in T₁ (13.66) no. of nodes per plant recorded in weedy check (control).

Average fruit weight (g) of ten fruits showed highest in (T₄) Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding followed by (T₅) Pre emergence application of pendimethalin @ 6ml/l. + quazalofop ethyl 40-50 gm./h. at 20 DAS and lowest in (T₁) weedy check(control).

Fruit yield per plant clearly indicated that (T₄) Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding produced heights fruits yield (233.49g) followed by (T₆) Post emergence application of metribuzin @ 525 g/h. at 20 DAS. It was also clear that the lowest in T₁ (184.16g) fruit yield per plant recorded in weedy check (control).

Total Pod yield/plant (t./ha.) clearly indicated that (T₄) Pre emergence application of pendimethalin @ 6ml/l. + one hand weeding produced highest pod yield/plant (15.99) followed by (T₆) Post emergence application of metribuzin @ 525 g/h. at 20 DAS. It was also clear that the lowest in T₁ (12.49) pod yield/plant recorded in weedy check (control).

References-

Bhooshan, Brij and Singh, V. K. (2014) Effect of planting method, Irrigation schedule and weed management practice on the Performance of fieldpea (*Pisum sativum* L. arvense). **Journal of Food Legumes**27 (2): 112-116

Bhalla, P. L. (1978) Weed competition, crop losses and chemical weed control in onion (*Allium cepa* L.)- **A review Pestology**2 (11): 35-39.

Chadha, Rameshwar; Singh, Sanjay; Sharma, G. D. and Rana Surinder (2002) Evaluation of herbicides for weed control and economics in onion. **Indian Journal of weed Science**34 (1-2): 68-71.

Chopra, Nisha and Chopra, N. K. (2007) Production of weed- free mother bulb of onion (*Allium cepa*) through integration of herbicides and weeding. **52(1):80-82. Indian Journal of Agronomy.**

Choudhary, B (1992) Vegetables. New Delhi, **National Book Trust:52-59.**

Dalavai, Kandasam; Hanumanthappa, B.L and Arasumallaiah, O.S. A. (2008) Effect of herbicides and their application Techniques on weed flora in onion **Environment and Ecology** 26 (4): 2157-2160.

Gopalan, C; Sastri,B.V.Rama and Balasubramanian, S. C. (1989) Nutritive value of Indian foods, **National Institute of Nutrition. ICMR, Hyderabad, India. 156.**

Gupta, O. P and Lamba, P. S. (1978) Modern Weed Science. **Today and Tomorrow's Printers and Publishers, New Delhi.**

Kumar, Abhay and Choudhary, B. M (2004) Weed management of okra (*Abelmoschus esculentus* (L.) Moench). **Orissa Journal of Horticulture**32(1): 73-74.

Kumar, Naresh and Mourya, I. B. (2006) Effect of different herbicides on growth and efficiency for weed control in onion. **Annals of Agricultural Research. 27 (3):245-249.**

Kumari, Usha; ReddyK; ReddyI.P; ReddyM. D and Neeraja G. (2001) Effect of spacing and weed control methods on growth and yield of onion. **Indian Journal of Weed Science** 33 (3&4):222-224.

Mehta, Ahsok; Rangaswamy,V; Kader, P. and Mohideen, M. (1979) Effect of herbicides on weed control and yield in okra (*Abelmoschus esculentus* (L.) Moench). **South Indian Horticulture**27(2): 88-93

Patel, R. B; Patel, B. D; Meisuriya, M. I. and Patel, V. J. (2004) Effect of methods of herbicide application on weeds and okra (*Abelmoschus esculentus* (L.) Moench). **Indian Journal of Weed Science**36 (3/4): 304-305.

Sharma, R. C. and Khandwe, Rupendra (2008) Response of weed control measures in kharif onion. **Research on Crops**9 (2): 348-349.

Sharma, Shweta and Patel, B.D. (2011): Weed management in okra grown in kharif season under middle Gujarat conditions. **Indian Journal of Weed Science**43 (3&4):226-227.

Singh, Moolchand, Prabhukumar, S. and Sairam, C.V. (2010) Integrated Weed management in okra (*Abelmoschus esculentus* (L.) Moench). **Annals of Plant Protection Sciences** 18 (2):481-483.

UNDER PEER REVIEW