

Seed Health Testing of Recombinant Inbred Lines Derived from Cross between Kalonunia and Pusa Basmati 1637

ABSTRACT

Detection seed born pathogen is important as it disseminate to the next crop using seed as medium. Identification of the hidden pathogen with seed will offer the way to take precautionary measured for seed treatment. Study was carried out to evaluate the seed health, such as germination, pre-emergence death, post-emergence test, infection (%) and presence of different seed-born pathogen of 11 inbred lines developed from a cross between Kalonunia and Pusa Basmati 1637 along with these two parents. Four different genera of pathogens were identified, *Aspergillus*, *Penicillium*, *Curvularia* and *Magnaporthe*. *Aspergillus* was predominant and it was found in all the seed lots. As the duration of seed in storage increased the germination reduced. No notable increase or decrease was noted with the duration of seed storage.

Keywords: Seed born pathogen, rice, inbred lines

1. INTRODUCTION

Seeds are the cornerstone of agriculture, cradling the potential for bountiful harvests and a thriving food system. However, seeds can harbour unseen threats, the pathogens like fungi, bacteria, and viruses that can wreak havoc on crops. This hidden danger underscores the critical importance of seed health, a concept encompassing the absence of seed-borne diseases and pests. Diseased seeds can significantly reduce yields by estimates ranging from 15% to a staggering 90%, and in few cases 100% impacting food security and economic stability.

The spread of these pathogens through infected seeds can introduce new diseases to previously unaffected areas, jeopardizing entire agricultural ecosystems. Ensuring seed health is paramount not only for maximizing crop yields but also for safeguarding food security and environmental sustainability. In recent years seed has become an international commodity used to exchange germplasm around the world. Seed is, however, also an efficient means of introducing plant pathogens into a new area as well as providing a means of their survival from one cropping season to another [33].

Seed health testing is thus routinely carried out in most countries for domestic seed certification, quality assessment and plant quarantine [7]. Seed health testing is an integral for all seed companies in disease risk management [10]. The test used depends on the organism being tested for and the purpose of the test quality assurance

or phytosanitary purposes when seed is exported [11]. Maintaining seed health is not merely about protecting individual plants; it safeguards entire agricultural ecosystems.

By prioritizing seed health, we pave the way for a more resilient agricultural future. Seed-borne fungi are one of the most significant biotic constraints on seed production worldwide. They cause both pre-emergence and post emergence death of seeds affecting seedling vigour which in result causes morphological variance in plants and reduction in germination [31,16]. Fungi outnumber all other types of pathogens that attack plants and cause a very serious economic impact on agricultural production due to their ability to induce diseases of cultivated crops that result in important yield losses [18]. The degree of seed infection is contingent upon various environmental factors, including elevated relative humidity, optimal temperature, and a high seed moisture content. Considering importance of presence of seed-borne pathogen in stored seed, the present study was performed to detect the presence seed associated pathogens in different seed lots of recombinant inbred lines of rice.

2. MATERIALS AND METHODS

The experiment was conducted during August, 2023 and November, 2023 at the Department of Seed Science and Technology and the Department of Plant Pathology, Uttar Banga Krishi Viswavidyalaya, Pundibari, Cooch Behar, West Bengal.

The materials used were obtained from Niche Area of Excellence's ongoing project focused on developing rice lines with enhanced blast resistance. Eleven inbred lines were used along with one blast resistant parent and one blast susceptible (Kalonunia) parent. Agar plate is the most common method used for identification of seedborne fungi [20]. The Agar Plate Method was used to assess the presence and incidence of seed-borne fungi. Forty seeds were collected from a random sample and tested for Agar Plate Method. Water agar at 8g/L concentration was prepared following standard protocols. The prepared water agar solution was autoclaved for sterilization at 121°C for 30 minutes at 15 psi. The sterilized agar solution was then poured into pre-sterilized 90mm (90*17mm) diameter plastic petri dishes at 30ml/plate to create a solidified growth medium in a laminar airflow cabinet to avoid contamination.

A random sample of 40 seeds were collected and each sterilized petri dish containing solidified water agar medium received 10 seeds. Four petri dishes were used for each sample, resulting in a total of 26 petri dishes. The petri dishes containing seeds were sealed and incubated at a constant temperature of 20°C ± 2°C under alternating cycles of 12 hours light and 12 hours darkness for 7 days. After incubation, the seeds were examined under a stereo binocular microscope to record the incidence of different seed borne fungi. Different

fungal growth was identified based on their morphological characteristics. The number of seeds colonized by each fungal type was recorded for each petri dish. Germination of these incubated seeds were also recorded. Infection percentage was calculated as:

$$\text{Infection (\%)} = \frac{\text{Infected number of seeds (pre – emergence + post – emergence)}}{\text{Total seeds}} \times 100$$

3. RESULTS AND DISCUSSION

Rice suffers from several biotic and abiotic stresses during cultivation and seed storage resulting heavy losses to farmers and reduction seed quality during storage. Among the several constrains to rice production, diseases caused by fungi, nematodes and bacteria cause major economic loses [25,26,21]. Fungi play significant roles in reducing the quality of rice seed during infection [14].

3.1. Analysis of Variance

The analysis of variance test is the initial step in analysing factor that effect a given data set. The statistical analysis of variance of seed tested during August, 2023 and November, 2023 showed highly significant variation among the rice genotypes for seed germination, pre-emergence death, post emergence death and infection (Table 1). The statistical analysis of pooled data also showed high signification variation among the rice genotypes for all the four characters.

Table 1. ANOVA of different components For August month

Source of Variation	D.f.	Mean Sum of Square (August, 2023)			
		Germination %	% Pre emergence death	% Post Emergence Death	% Infection
Replication	1	0.000	0.000	61.538	61.538
Treatment	12	253.846***	253.846 ***	70.513 .	225.000 **
Error	12	33.333	33.333	28.205	36.538
Mean Sum of Square (November)					
Replication	1	138.462	246.154 *	61.538	553.85 **
Treatments	12	221.154 ***	213.462 **	59.615 NS	211.54 *
Error	12	30.128	37.821	69.872	53.85
Mean Sum of Square (Pooled ANOVA)					
Replication	1	15.385	34.615	3.846	15.385
Treatments	12	182.532 *	192.788 **	52.885 **	184.135 **
Error	12	54.968	40.865	12.179	34.135

Significant codes: 0 '****' 0.001 '***' 0.01 '**' 0.05 '*' .

3.2. Seed Germination (%)

The seed germination varied from 75.00-100.00% and 70.00-100.00% with a mean of 88.85 and 84.62% during August and November, respectively (Table 2A). Maximum seed germination was reported for NAE-6, NAE-10 and NAE-12 (100.00%) during when tested during August and lowest seed germination was recorded for KaloNunia.

The genotype NAE-10 and NAE-12 retained 100.00% germination when they were tested during November (Table 2A). The germination (%) of the breeding lines, namely 192, 207, NAE-11 and KaloNunia remained the same during August and November. However, the germination fallen down for all the breeding lines when tested during November.

The mean of August and November varied from 75.00 to 100.00% with a mean of 86.73% (Table 2A). Highest mean germination was noted for NAE-10 and NAE-12 (100.00%). Lowest mean germination was recorded for KaloNunia and NAE-4 (75.00%). Some of the fungal pathogens are known to cause seed rot leading to decrease in seed germination, pre-emergence death and post-emergence death [1].

3.3. Pre-emergence Death (%)

The death of a seedling before or shortly after emergence due to decomposition of the root and/or lower stem by pathogen is known as pre-emergence death (Fig. 1). With pre-emergence death, seeds and seedlings are affected during or after germination but before emergence resulting in poor stands.

The pre-emergence death varied from 0.00 to 30.00% in both of the testing time (Table 2A). With a mean of 12.31% and 14.62%, respectively during August and November. The mean of two time of testing of respective individual varied from 0.00 to 27.00%. It was found that the pre-emergence death increased from 12.31% to 14.62% with passes of storage duration. The breeding lines NAE-10 and NAE-12 did not show any pre-emergence death during entire period of testing. Maximum pre-emergence death (27.00%) was observed for the breeding line 205 and Kalonunia. Some contradictory result was also obtained for the breeding line 207. It showed pre-emergence death of 10.00% during August, whereas no pre-emergence death was recorded during November. Kalonunia had 30.00% pre-emergence death during August which decreased to 25.00% during November.

3.4. Post-emergence Death (%)

If seedlings emerge and then wilt and collapse, that is called post-emergence death (Fig. 1). Post-emergence death also deteriorates the crop stand. It varied from 0.00 to

15.00% and 0.00 to 20.00% during August and November testing (Table 2A). Four breeding lines, namely 192, 205, NAE-3 and NAE-11 showed no post-emergence death during entire period of testing. Highest mean of post-emergence death (15.00%) was recorded for the breeding line 207 followed by NAE-6 (12.00%), NAE-1 (10.00%) and NAE-10 (10.00%). Some contradictory results also reported for this character. The breeding lines 182 and NAE-6 had lower post-emergence death during November as compared to August.

Table 2A. Seed health parameters and associated pathogens

Entries	Germination (%)			Pre-emergence death (%)			Post-emergence death(%)		
	D1	D2	Mean	D1	D2	Mean	D1	D2	Mean
182	95.00	90.00	92.50	15.00	10.00	12.50	10.00	5.00	7.50
192	80.00	80.00	80.00	20.00	20.00	20.00	0.00	0.00	0.00
205	80.00	75.00	77.50	30.00	25.00	27.50	0.00	0.00	0.00
207	90.00	90.00	87.50	10.00	0.00	5.00	10.00	20.00	15.00
NAE-1	90.00	70.00	80.00	10.00	30.00	20.00	10.00	10.00	10.00
NAE-3	90.00	85.00	87.50	10.00	15.00	12.50	0.00	0.00	0.00
NAE-4	80.00	70.00	75.00	20.00	30.00	25.00	5.00	5.00	5.00
NAE-6	100.00	90.00	95.00	0.00	10.00	5.00	15.00	10.00	12.50
NAE-10	100.00	100.00	100.00	0.00	0.00	0.00	10.00	10.00	10.00
NAE-11	95.00	95.00	95.00	10.00	5.00	7.50	0.00	0.00	0.00
NAE-12	100.00	100.00	100.00	0.00	0.00	0.00	0.00	5.00	2.50
PB 1637	95.00	80.00	87.50	5.00	20.00	12.50	10.00	0.00	5.00
KaloNunia	75.00	75.00	75.00	30.00	25.00	27.50	10.00	5.00	7.50
Mean	88.85	84.62	86.73	12.31	14.62	13.46	6.15	5.38	5.77
Range	75.00-100.00	70.00-100.00	75.00-100.00	0.00-30.00	0.00-30.00	0.00-27.00	0.00-15.00	0.00-20.00	0.00-15.00
LSD (5%)	11.98	13.39	15.56	11.98	15.85	13.72	NS	NS	7.33
LSD (1%)	16.70	18.68	21.70	16.70	22.10	19.14	NS	NS	10.23




Fig. 1. Pre-emergence death shown in red arrows. Post-emergence death shown in green arrow

3.5. Infection (%)

Infection occurs when the pathogen invades the plant tissue and establishes a parasitic relationship between itself and the plant. These pathogens cause seed discolouration, seed rot, reduce seed germination, and vigour in seedlings, as well as making the plant weak during its early growth period.

The infection varied from 0.00 to 40.00% and 5.00 to 40.00% during August and November, respectively (Table 2B). No pathogen was reported for NAE-12 when tested during August, however it showed 5.00% infection when tested during November. Maximum infection mean of two testing times was observed for Kalonunia (35.00%) followed by NAE-1 (30.00%), NAE-4 (30.00%), 205 (27.00%), 182 (20.00%), 207 (30.00%) and 192 (30.00%).

3.6. Presence of Pathogens

Seed infection is dominated by fungal pathogen. Fungal pathogens that are seed-borne are comparatively challenging to manage because the fungal hyphae get established and becomes dormant. Fungi infects crops on the field and they persist and proliferate in storage resulting in increased fungal contamination with duration of seed storage [3]. Most crop diseases that are important economically are seed-borne and seed transmitted, including blast disease of rice. With the consideration that the pathogenic seed transmission plays an important role in the spread and development of epidemic diseases in rice, then the seed health tests need to be incorporated into the seed certification process. Seed health test is required to detect the presence of pathogen or seed health status.




Fig. 2. *Aspergillus niger*

In this study, the major fungal genera contaminating rice in order of decreasing predominance were *Aspergillus*, *Penicillium*, *Curvularia* and *Magnaporthe*. Only one seed lot had *Magnaporthe* pathogen. *Aspergillus* and *Penicillium* were the most common mycoflora [17,28,8,9,19] as found in this piece of work.

All the seed lots had the infection of seed-borne pathogen and *Aspergillus niger* (Fig. 2) was found in all the seed lots (Table 2B). *Aspergillus* is one of the dominant fungi in rice, contaminating in rice of Malaysia [23], India [22], Philippines [24] and Vietnam [29]. *Aspergillus* is most prevailing fungus in storage of rice seed reported by [8,12,19 and 27].

Penicillium was in seed lot of 182, NAE-3 and NAE-11 (Fig. 3A&B). Citrinin is considered to be the major causal agent of the yellow rice disease described for a long time in Japan. The characteristic colour in this product is due to the presence of citrinin-producing *Penicillium* species, mainly *Penicillium citreoviridae*, *Penicillium atrium* and *Penicillium islandicum*. *Penicillium* was categorized as one of the most predominant fungi in storage of rice [8].

Curvularia was reported in 207, NAE-10 and NAE-11 (Fig. 3C). It is also known as predominant fungus in rice seed storage. Species of the genus *Curvularia* has been reported to infect the embryo of the seed, therefore, reducing the percentage of germination of rice seeds [9,4,6, 30, 2].

Table 2B. Seed health parameters and associated pathogens

	Infection (%)			Pathogen identified		
	D1	D2	Mean	D1	D2	All Pathogen
182	25.00	15.00	20.00	<i>Aspergillus, Pencillium</i>	<i>Aspergillus</i>	<i>Aspergillus, Penicillium</i>
192	20.00	20.00	20.00	<i>Aspergillus</i>	<i>Penicillium, Aspergillus</i>	<i>Penicillium, Aspergillus</i>
205	30.00	25.00	27.50	<i>Aspergillus</i>	<i>Aspergillus</i>	<i>Aspergillus</i>
207	20.00	20.00	20.00	<i>Aspergillus, Curvularia</i>	<i>Magnaporthe</i>	<i>Aspergillus, CurvulariaMagnaporthe</i>
NAE-1	20.00	40.00	30.00	<i>Aspergillus</i>	<i>Aspergillus</i>	<i>Aspergillus</i>
NAE-3	10.00	15.00	12.50	<i>Aspergillus</i>	<i>Aspergillus, Penicillium</i>	<i>Aspergillus, Penicillium</i>
NAE-4	25.00	35.00	30.00	<i>Aspergillus</i>	<i>Aspergillus</i>	<i>Aspergillus</i>
NAE-6	15.00	20.00	17.50	<i>Aspergillus</i>	<i>Aspergillus</i>	<i>Aspergillus</i>
NAE-10	10.00	10.00	10.00	<i>Curvularia</i>	<i>Aspergillus</i>	<i>Curvularia, Aspergillus</i>
NAE-11	10.00	5.00	7.50	<i>Aspergillus, Curvularia</i>	<i>Aspergillus, Penicillium</i>	<i>Aspergillus, Curvularia, Penicillium</i>
NAE-12	0.00	5.00	2.50	<i>Aspergillus</i>	None	<i>Aspergillus</i>
PB 1637	15.00	20.00	17.50	<i>Aspergillus</i>	<i>Aspergillus</i>	<i>Aspergillus</i>
Kalonunia	40.00	30.00	35.00	<i>Aspergillus</i>	<i>Aspergillus</i>	<i>Aspergillus</i>
Mean	18.46	20.00	19.23	-	-	-
Range	0.00-40.00	5.00-40.00	2.50-35.00	-	-	-
LSD (1%)	13.39	NS	12.35	-	-	-
LSD (5%)	18.68	NS	17.23	-	-	-

Only one seed lot, namely 207 has been infect with *Magnaporthe*(Fig. 3D).*Magnaporthe (Pyriculariaoryzae)* is reported to cuase very low seed infection by [13]. Earlier

the pathogen of rice blast was known as *Pyriculariaoryzae*. It is a dynamic pathogen and can adapt quickly to the conditions of the host plant. This pathogen also has a high degree of genetic diversity and the ability to produce the new breeds [15].




Fig. 3. A&B) *Penicillium*; C) *Curvularia*; D) *Pyriculariaoryzae*

CONCLUSION

The experiment was conducted to appraise the presence of seed-born pathogen in 13 rice seed lots. Experiment was carried out in agar plate. Germination reduced with increase in storage duration. Pre-emergence and post-emergence death were also recorded. Four different pathogens were identified in this study, namely *Aspergillus*, *Penicillium*, *Curvularia* and *Magnaporthe*. Among these, *Aspergillus* was dominating and it was found in all the seed lots followed by *Penicillium* and *Curvularia*. This piece of study will provided the information about the presence of rice seed born pathogen under agro-climatic condition of Tarai Zone of West Bengal. However, this experiment could be carried out with a particular interval for one year on stored seed.

ACKNOWLEDGEMENT

We are thankful to Indian Council of Agricultural Research for providing the financial assistance for research on development of blast resistant genotypes under Niche Area of Excellence project (Sanction order No. Edn. 5(12)/2017-EP&HS, dated 18.11.2019).

REFERENCES

1. Al-Kassim M and Monawar M. Seed-borne fungi of some vegetable seeds in Gazan province and their chemical control. Saudi. Journal of Biological Science. 2000; 7:179-184.
2. Ashfaq M, Shaukat M, Akhter M, Haider M, Mubashar U and Hussain S. Comparison of fungal diversity of local and exotic rice (*Oryza sativa* L.) Germplasm for their seed health. Journal of Animal and Plant Science. 2015; 25:1349-1357.
3. Bainton SJ, Coker RD, Jones BD, Morley EM, Nagler MJ and Turner RI. Mycotoxin training Manual. Tropical Product Institute. London. 1980; 1:176.
4. Bautista E and Opina O. Isolation, identification and pathogenicity tests of seedborne fungi associated with cowpea seeds. Pest Management Council of the Philippines, Manila Philippines, 8-11 May 1991.
5. Bolanle Tolani Edun A, Yahuza Lurwanu A, Mustapha Sunusi B, Ali A and Sulaiman A. Seed Health, Quality Test, and Control of Seed-borne Fungi of Some Improved and Local Cultivars of Rice (*Oryza sativa* L.) in Kano, Northwestern Nigeria. Journal of Tropical Crop Science 2019; 6(3):145-152.
6. Butt AR, Yaseen SI and Javaid A. Seed borne mycoflora of stored rice grains and its chemical control. J. Animal Plant Sci. 2011; 21(2): 821-824.
7. FAO. Seeds in Emergencies. A technical handbook. Plant production and protection Paper. 2010:202.
8. Hussaini A Makun, Timothy A Gbodi, Olufunmilayo H Akanya, Ezekiel A Salako and Godwin H Ogbadu. Fungi and some mycotoxins contaminating rice (*Oryza sativa*) in Niger State, Nigeria. African Journal of Biotechnology. 2007; 6(2):099-108.
9. Imolehin E. The rice seed multiplication centres in relation to seed borne pathogens of rice. A case study of Ondo State Rice Multiplication Centres. Nigerian Journal of plant Protection. 1987; 11:37-42.
10. International Seed Federation (ISF). Guidelines on Seed Health Testing in the Vegetable Seed Industry. 2010
11. ISTA. International Rules for Seed Testing. A next Chapter 7 Seed Health Testing. Seed Health Testing Methods. International Seed Testing Association, Bassersdorf, Switzerland. 2009:(b).

12. Mansur Ahmed M, Mehbub Hossain, Kamrul Hassan and Chandra Kanta Dash. Seed Health and Quality Test of Three Rice Varieties for the Detection of Fungi Associated with Seed Sample. *Universal Journal of Plant Science*. 2013;1(2): 37-42.
13. Martini T, EvyPujiastuti, SitiNurhaeni, MekkyKusumaDewi, NurlIndrayatiPrabaNingrum, and danMansyur. Evaluation of Locally Rice Seeds Heal. *AGRIC*. 2022; 34(1):45-56.
14. Nanda JS and Sharma SD. Crops that feed the world 7: rice Food Security. 2012; 4(1):7-24.
15. Nasution A dan and Usyati N. Observasi ketahanan varietas padi lokal terhadap penyakit blas (*Pyricularia grisea*) di rumah kaca. *Balai Besar Penelitian Tanaman Padi*. 2015; 1:19-22.
16. Niaz I and Dawar S. Detection of seed borne mycoflora in maize (*Zea mays* L.). *Pak. J. Bot.* 2009; 41(1): 443-451.
17. Ominski KH, Marquardi RR, Sinha RN and Abramson D. Ecological aspects of growth and mycotoxin production by storage fungi. In: Miler JD and Trenholm HL. *Mycotoxins in grains: Compounds other than aflatoxins*. Eagan Press, St. Paul Minnesota, USA. 1994:287 -314.
18. Paplomatas EJ. Molecular Diagnostics of Fungal Pathogens. *Arab J. Pl. Prot.* 2006;24:147-158.
19. Phan LTK, Trang Minh Tran, Kris Audenaert, Liesbeth Jacxsens and Mia Eeckhout. Contamination of *Fusarium proliferatum* and *Aspergillus flavus* in the Rice Chain Linked to Crop Seasons, Cultivation Regions, and Traditional Agricultural Practices in Mekong Delta, Vietnam. *Foods*. 2021; 10: 2064.
20. Rao NK and Bramel PJ. Manual of genebank operations and procedures. Technical Manual no. 6. ICRISAT, Patancheru, India. 2000.
21. Reddy OR and Sathyanarayana N. Seed-borne fungi of rice and quarantine significance in Major Fungal Diseases of Rice. Springer, Berlin, Germany. 2001: 331–345.
22. Reddy K, Reddy, C and Muralidharan K. Detection of *Aspergillus* spp. and aflatoxin B1 in rice in India. *Food Microbiol.* 2009; 26: 27–31.
23. Reddy KR, Farhana NI and Salleh B. Occurrence of *Aspergillus* spp. and aflatoxin B1 in Malaysian foods used for human consumption. *J. Food Sci.* 2011;76: T99–T104.
24. Sales AC and Yoshizawa T. Mold counts and *Aspergillus* section *Flavi* populations in rice and its by-products from the Philippines. *J. Food Prot.* 2005;68:120–125.

25. Singh KM, A Jha, M Meena, and R Singh. Constraints of rainfed rice production in India: an overview in *Innovations in Rice Production*, Eds: P. K. Shetty, M. R. Hegde, and M. Mahadevappa. 2012: 71–84.
26. Strange RN and Scott PR. Plant disease: a threat to global food security. *Annual review of phytopathology*. 2005;43:83–116.
27. Suhendar MA, Koswanudin D, Hidayatun N, Diantina S, Manzila I, Zulchi PH, and Wawan. Detection of pathogens to test seed health of some food crops in storage. 2nd Agrifood System International Conference (ASIC-2022). IOP Conf. Series: Earth and Environmental Science. 2022: 1160.
28. Taligoola H, Ismail MA and Chebon SK. Mycobiota Associated with Rice Grains Marketed in Uganda. *J. Biol. Sci.* 2004; 4(1):271-278.
29. Trung TS, Bailly J, Querin A, Le Bars P and Guerre P. Fungal contamination of rice from south Vietnam, mycotoxinogenesis of selected strains and residues in rice. *Rev. De Medecine Veterinaire*, 2001;152: 555–560.
30. Utobo E, Ogbodo E and Nwogbaga A. Seedborne mycoflora associated with rice and their influence on growth at Abakaliki, Southeast Agro-Ecology, Nigeria. *Libyan Agricultural Research Center Journal International*. 2011;2: 79-84.
31. Van Du P, Loan LC, Cuong ND, Nghiep HV and Thach ND. Survey on seed borne fungi and its effects on grain quality of common rice cultivars in the Mekong Delta. *Omonrice*. 2001; 9:107-113.
32. Walcott RR. Detection of seedborne pathogens. *HortTechnology*. 2003;13:40–47.