

Preparation and evaluation of the Benefits of herbal compositions of Bacopa Extract and Bitterless Ebelin Lactone on various Biological Activities

ABSTRACT

Although Bacopa extract is well known for its cognitive and memory support, it has received minimal attention in pharmacy due to its terrible palatability and poor bioavailability. This study aimed to develop an improved, cost-effective, and eco-friendly process for the preparation of stable and highly bioavailable non-hygroscopic bacosides enriched fraction in such a manner that it contains a specific amount of Bacoside A3 and jujubogenin, and the aglycone derivative (Ebelin lactone). We prepared two compositions 1 and 2 having Bacoside A3 and jujubogenin: Ebelin lactone ratios 1:1 and 3:1 respectively. The efficacy of these herbal compositions was evaluated independently in comparison to the bioactivity of bacosides (20%), bacosides (50%), and Ebelin lactone. This study also aimed to find a wide range of potential activities of formulated composition, including acetylcholinesterase (AChE) inhibitor activity, anti-oxidant activity, cyclooxygenase (COX) inhibitor activity, and anti-cancer activity. We noted both that both composition worked better than parent Bacopa for biological activity assays but the herbal Bacopa Composition 1 is more potent than Bacopa Composition 2. Hence, further research on Composition 1 should be explored for its potential to treat conditions such as Alzheimer's and various cancer diseases.

Keywords: *Bacopa monnieri*, Ebelin lactone, anti-oxidant, anti-inflammatory, anti-cancer

1. Introduction

Medicinal plants have been used historically to treat a wide range of illnesses and are an important source of pharmacologically active compounds. Traditional medicinal plants are used every day by nearly 3.3 billion people all over the world (1). Many pharmacological components of medicinal plants are known that can be used in the creation of new drugs (2).

South and Southeast Asia are home to the *Bacopa monnieri* plant, often known as Bhrami, which is a member of the Scrophulariaceae family. It is well recognized for enhancing memory and reducing anxiety and has a long history in traditional medicine. Currently, the plant is promoted around the world as a blood sugar stabilizer and memory booster (3, 4).

Recently, researchers have focused on better understanding the processes and usefulness of *B. monnieri* in treating human illnesses. Bacopa extracts have been investigated as a traditional medicine and used for centuries for treating a variety of illnesses, such as providing relief to patients suffering from anxiety and skin ailments particular applications include the treatment of asthma, pain, inflammation, insanity, and epilepsy (5, 6). Perhaps the most prominent medicinal claim regarding its advantages has been that it enhances memory (7). The Bacopa plant, also known as a nootropic herb, aids in the healing of injured neurons, neuronal synthesis, and synaptic activity restoration, as well as improving brain function. Alkaloid brahmine, nicotine, herpestine, Bacosides A and B, saponins A, B, and C, and triterpenoid saponins (jujubogenin and pseudojujubogenin) are major phytochemicals found in *B. monnieri*(8). The primary bioactive components considered to be responsible for the cognitive effects of *B. monnieri* are believed to be characteristics of saponins known as "Bacosides," notably bacoside A (9, 10). Numerous *in vitro* and animal investigations have shown that the Bacosides present in *B. monnieri* strengthen the body's defenses against oxidative stress by reducing the buildup of free radicals in the brain (11). *In vitro* testing on human lymphocytes has found that extracts from *B. monnieri* are successful in exhibiting anti-mutagenic and free radical scavenging abilities with no evidence of genotoxic activity (12). These active constituents have been shown to have anticancer activity in addition to cytotoxic, anti-mutagenic, and anti-inflammatory properties (13). *In vitro*, the triterpene saponins, Bacopaside I (bac I) and Bacopaside II (bac II), derived from Bacopa leaf extract, demonstrated a synergistic effect by inhibiting breast cancer cell migration and invasion (14).

B. monnieri has neuropharmacological effects and can be administered orally. The active ingredients that provide pharmacological action must be ingested and CNS active, which means that the compound must cross the blood-brain barrier and be absorbed through the gut. Poor membrane permeability *via* the blood-brain barrier and gut is most likely caused by these adverse physicochemical characteristics of the parent Bacosides. However, by satisfying four of the requirements, the aglycones (jujubogenin, pseudojujubogenin, Ebelin lactone, and bacogenin A1) demonstrated superior CNS drug-like characteristics. The blood-brain barrier allows for extensive passive lipid-mediated transit of small compounds (16). Aglycones' tendency to permeate the brain also tends to increase as their lipophilicity rises. Orally administered substances for CNS activity is able to cross the blood-brain barrier (BBB) and be absorbed from the intestines (17, 18).

This study aims to develop a formulation with maximum effectiveness of bacopa extract to improve its membrane permeability and bioavailability (absorption capacity). The increased absorption is demonstrated by the herbal compositions containing Bacosides enriched with Bacoside A3, jujubogenin, and the aglycone derivative Ebelin lactone. To ensure the drug's safety, toxicity profiling was also performed. This study also aimed to explore its effectiveness as a potent acetylcholinesterase (AChE) inhibitor, anti-oxidant, cyclooxygenase inhibitor, and anti-cancer agent.

2. MATERIALS AND METHODS

Different compositions of bacopa extracts, which were a mixture of non-hygroscopic Bacosides enriched with Bacoside A3, jujubogenin, and aglycone derivative, i.e., Ebelin lactone, were assessed for their biological activities. Composition 1 was Mixture of (non hydroscopic bacosides enriched with Bacoside A3, jujubogenin)+ (aglycone derivative Ebelin lactone) combined in the ratio 1:1; and Composition 2 was the mixture of (non hydroscopic bacosides enriched with Bacoside A3, jujubogenin)+ (aglycone derivative Ebelin lactone) combined in the ratio 3:1 Their efficacy was evaluated independently against the bioactivity of Bacosides (20%), Bacoside (50%), and an aglycone derivative, Ebelin lactone. All the chemicals and kits used in the study were purchased from Thermofisher Scientific.

2.1 Determination of intestinal absorption:

The Caco-2 cell line was used as an *in vitro* model for studying the intestinal permeability of the herbal composition of Bacopa extract enriched with Bacoside A3, jujubogenin, and aglycone derivative Ebelin lactone (19–21). The absorption (permeability) tests were carried out using the maximum non-toxic concentration found in a preliminary set of cyto-toxicity testing. In three separate experiments, the permeability test was done at 37 °C for 2 hours in a shaking incubator with 100 RPM in the apical-basolateral (A-B) and basolateral-to-apical (B-A) directions. The samples were collected, and the amount of transported samples was determined using HPLC. To quantify the transported materials, reverse-phase chromatography using a C18 column and a UV-VIS detector was used.

Data for the permeability investigation was calculated using MS Excel. The apparent permeability coefficient (Papp) was determined using the equation:

$$(dQ / dt) / Papp (A \times C),$$

where dQ/dt ($\mu\text{g}\cdot\text{s}^{-1}$) is the rate of drug transport, A is the surface area of the cell monolayer (cm^2), and C is the initial drug concentration on the administered side ($\mu\text{g}\cdot\text{ml}^{-1}$).

$$\text{Efflux ratio} = Papp (BA)/Papp (AB)$$

where Papp (BA) is the apparent permeability coefficient for the test substance transported from the basal to the apical side (secretive direction) and Papp (AB) is the apparent permeability coefficient for the test substance transported from the apical to the basal side (secretive direction) (absorptive transport) (21, 22).

For drug absorption studies, Caco-2 cells were seeded on 35mm culture plates (80,000–100,000 cells) and allowed to adhere for 12 hours. Subsequently, the cells were treated with various herbal Bacopa compositions at 1 mg/ml dosage over different time intervals (0–360 min). After treatment, cell culture medium was collected, stored at -80 °C, and the cells were trypsinized,

washed, and lysed in buffer. The lysed cells underwent sonication, followed by centrifugation to collect the supernatant. Protein precipitation was carried out using acetone, and the resulting mixture was centrifuged. The supernatant was then concentrated, removing acetone through vacuum concentration at 40 °C for 40 minutes. The final cell lysate was stored at -80 °C.

The chromatogram profile was obtained by HPLC analysis of the medium and cell lysate samples to determine drug absorption by the cell. The HPLC analysis was carried out with Shimadzu's Lab Solutions software and a photo diode array detector to determine the qualitative analysis of the herbal bacopa compositions with the aid of an analytical standard. Bacosides and Ebelin lactone were separated using a C18-BDS column and a Restek ROC C18 column, respectively, with the appropriate solvent systems serving as the mobile phase.

2.2 Acetylcholinesterase Inhibition Assay

The amount of acetylthiocholine hydrolyzed by AChE to produce thiocholine was measured by the Acetylcholinesterase Assay Kit using DTNB. Acetylcholinesterase-generated thiocholine takes on a yellow hue when combined with 2, 5-nitrobenzoic acid. The product's color intensity at 412 nm is proportional to the amount of enzyme activity in the sample.

2.3 Determination of Radical scavenging activity

At various concentrations, the antioxidant activity of bacosides enriched with bacoside A3, jujubogenin, and an aglycone derivative, namely Ebelin lactone, was investigated. Ascorbic acid (1mg/100µl) was employed as a reference standard. The concentrations analyzed for Bacopa samples were 1 µg/µl, 2.5 µg/µl, 5 µg/µl and 10 µg/µl. After adding DPPH, the reaction mixture was stirred and incubated for 30 minutes at 25 °C. A semi-automatic analyzer was used to detect absorbance at 517 nm (24, 25). By comparing the absorbance readings of the test and control samples, the percentage inhibition was calculated.

2.4 Determination of Glutathione Reduction Assay

The herbal composition containing Bacosides enriched with Bacoside A3, jujubogenin, and the aglycone derivative Ebelin lactone, Bacosides 20%, Bacosides 50%, and Ebelin lactone was assessed for its ability to generate ROS. Ascorbic acid (1 mg/ml) was employed as a reference standard. The herbal compositions along with the Bacosides 20%, Bacosides 50% and Ebelin lactone were evaluated against 1 µg/µl concentrations of the reference standard (26). A semi-automatic analyzer was used to measure the absorbance at 412 nm.

2.5 Determination of Anti-inflammatory activity

A test that is quick, simple, sensitive, and dependable for screening COX-2 inhibitors is provided by the COX-2 inhibitor screening kit. This assay uses fluorometry to detect prostaglandin G₂, a COX enzyme intermediate (27). The sample was scanned for fluorescence with an excitation wavelength of 535 nm and an emission wavelength of 587 nm for 10 minutes (20 cycles) along with a time interval of 30 seconds.

2.6 In vitro Anti-cancer assay

Breast cancer cells (MDA-MB-468) and colorectal cancer cells (HCT116) were analysed for anti-cancer activity. DMSO was used as a negative control. As a positive control, doxorubicin was used. Mallick et al., (2015) method was used (28). A Tecan Multimode reader was used to record the data and plot a non-linear regression curve against the log concentration versus absorbance

2.7 In vitro Cytotoxicity assay

The Kidney Epithelial cells (Hek293T) and Skin Fibroblast cells (HFF-1) were used for cytotoxicity assay. 0.1 mg/ml, 0.25 mg/ml, 0.5 mg/ml, and 1 mg/ml of each sample were tested. DMSO was used as a negative control. Standard method explained by Mallick et al., (2015) was used for the study. At 24, 48, and 72 hours, spectrophotometric readings were taken at 570 nm (28).

3. RESULTS AND DISCUSSION

3.1 Determination of intestinal absorption

The transport of active constituents in an apical to basolateral (A-B) direction was investigated at a concentration of 0.125 mg/ml. Composition 1 (1:1) was found to have the highest apparent permeability coefficient (P_{app}) in absorptive transport at $141 \pm 18.7 \text{ cm} \cdot \text{s}^{-1}$ compared to secretive direction at $77.4 \pm 6.2 \text{ cm} \cdot \text{s}^{-1}$ and an efflux ratio of 0.54 (Figure 1). Whereas Composition 2 (3:1) was found to have an apparent permeability coefficient (P_{app}) in absorptive transport (A-B) as $133 \pm 26.3 \text{ cm} \cdot \text{s}^{-1}$ compared to secretive direction (B-A) as $77.4 \pm 10.2 \text{ cm} \cdot \text{s}^{-1}$ and with an efflux ratio of 0.57. Enriched aglycone derivative Ebelin lactone displayed an apparent permeability coefficient (P_{app}) in absorptive transport as $121 \pm 11.3 \text{ cm} \cdot \text{s}^{-1}$ compared to the secretive direction as $78.4 \pm 8.4 \text{ cm} \cdot \text{s}^{-1}$ and with an efflux ratio of 0.647. Whereas Bacoside 20% and 50% showed apparent permeability coefficient (P_{app}) in absorptive transport as $101 \pm 27.3 \text{ cm} \cdot \text{s}^{-1}$ and $113 \pm 13.5 \text{ cm} \cdot \text{s}^{-1}$ respectively and secretive direction apparent permeability coefficient (P_{app}) as 62.4 ± 5.3 and 74.4 ± 9.6 respectively and with an efflux ratio of 0.62 and 0.66 respectively (Table 1).

The Caco-2 cells completely absorbed Bacopa Composition 1 (1:1) from the media 15 minutes later, as demonstrated by HPLC qualitative determination. The Caco-2 *in vitro* absorption study revealed that Compositions 1, composed of Bacosides and Ebelin lactone in a 1:1 ratio, had better membrane permeability and absorption than Bacoside 20%, Bacoside 50%, enriched aglycone derivative Ebelin lactone, and Compositions 2 (3:1), which were evaluated independently.

3.2 Acetylcholinesterase Inhibition assay

An acetylcholinesterase inhibition assay showed 34% and 46% inhibition of bacoside 20% and bacoside 50% inhibited acetylcholinesterase respectively. Enriched aglycone derivative Ebelin lactone and Composition 2 were found to inhibit Acetylcholinesterase with 65% and 82% inhibition at ~ 1 mg/ml, respectively. Composition 1 (1:1), on the other hand, was found to potentially inhibit acetylcholinesterase with 85% inhibition at ~ 1 mg/ml (Figure 2).

3.3 Free Radical scavenging activity of the Bacopa herbal Compositions

A DPPH assay resulted as Both the Bacopa herbal compositions exhibited good anti-oxidant activity. The IC₅₀ values of Bacoside 20% and Bacoside 50% were found to ~ 96.12 µg/ml and ~ 84.65 µg/ml, respectively. The IC₅₀ values of the Enriched aglycone derivative Ebelin lactone and Composition 2 were found to ~ 45.34 µg/ml and ~ 52.12 µg/ml, respectively. Composition 1 (1:1), on the other hand, demonstrated potential anti-oxidant activity with an IC₅₀ value of ~ 58.32 µg/ml in comparison to standard ascorbic acid with an IC₅₀ value of ~ 7.12 µg/ml (Figure 3).

3.4 Glutathione Reduction Assay

Bacoside 20% and bacoside 50% inhibited glutathione with 34% and 25% inhibition, respectively. The enriched aglycone derivative Ebelin lactone and composition 2 (3:1) inhibit glutathione with 13% and 27% inhibition, respectively, at ~ 1 mg/ml. Composition 1 (1:1) was found to reduce glutathione with 22% inhibition at ~ 1 mg/ml in comparison to ascorbic acid (Positive control), which exhibited only 7% glutathione reduction (Figure 4).

3.5 Synergistic cyclooxygenase Inhibitory Activity

Bacoside 20% and Bacoside 50% was found to inhibit COX-2 with 76% and 83% inhibition at ~ 1 mg/ml, respectively. Enriched aglycone derivative Ebelin lactone and Composition 2 (3:1) was found to inhibit COX-2 with 89.5% and 90% inhibition at ~ 1 mg/ml, respectively. Whereas, Composition 1 (1:1) was found to potentially inhibit COX-2 with 91.1% inhibition at ~ 1 mg/ml in comparison to celecoxib (Positive control), which exhibited only 69% COX-2 inhibition (Figure 5).

3.6 In vitro Anti-cancer activity

Bacoside 20% (Figures 6A and 7A) was found to effectively inhibit MBA-MD-468 and HCT116 cells with an IC₅₀ value of ~ 0.2 mg/ml and ~ 0.06 mg/ml, respectively. Bacoside 50% (Figures 6B and 7B) was found to effectively inhibit MBA-MD-468 and HCT116 cells with an IC₅₀ value of ~ 0.03 mg/ml and ~ 0.09 mg/ml, respectively. Ebelin lactone, an enriched aglycone derivative (Figures 8C

and 9C), was found to effectively inhibit MBA-MD-468 and HCT116 cells with IC₅₀ values of 0.210 mg/ml and 0.055 mg/ml, respectively. Composition 1 (1:1) (Figures 6D and 7D) was found to effectively inhibit MBA-MD-468 and HCT116 cells with an IC₅₀ value of ~ 0.11 mg/ml and ~ 0.055 mg/ml, respectively. Whereas, Composition 2 (3:1) (Figures 6E and 7E) was found to effectively inhibit MBA-MD-468 and HCT116 cells with an IC₅₀ value of ~ 0.11 mg/ml and ~ 0.055 mg/ml, respectively.

3.7 In vitro Toxicity Studies

Bacoside 20% (Figures 8A and 9A) was found to be non-toxic against Hek293T and HFF-1 with an IC₅₀ value of ~ 0.6 mg/ml and ~ 1.026 mg/ml, respectively. Bacoside 50% (Figures 8B and 9B) was found to be non-toxic against Hek293T and HFF-1 with an IC₅₀ value of ~ >10 mg/ml and ~ 2.151 mg/ml, respectively. Ebelin lactone (Figures 8C and 9C) was discovered to be non-toxic against Hek293T and HFF-1, with IC₅₀ values of ~ 0.9 mg/ml and ~ 0.862 mg/ml, respectively. Composition 2 (3:1) (Figures 8E and 9E) was found to be safe and non-toxic against Hek293T and HFF-1 with an IC₅₀ value of ~ 0.8 mg/ml and ~ 1.052 mg/ml, respectively. Whereas, Composition 1 (1:1) (Figures 8C and 9C) was found to be safe and non-toxic against Hek293T and HFF-1 with an IC₅₀ value of ~ 0.9 mg/ml and ~ 1.583 mg/ml, respectively.

The major active constituents of *B. monnieri* are the steroidal saponins Bacosides A, which is a combination of Bacoside A3, Bacopaside II, Bacopaside X, and Bacopasaponin C. Bacoside A is thought to be a mediator of *B. monnieri*'s memory-enhancing and cognitive effects (29, 30). Nevertheless, there isn't sufficient information regarding the underlying mechanism for the bacoside components' activity. Previous studies have revealed that, the bacoside A which cater the pharmacological action in order to be CNS active is improbable to be absorbed in the gut so as to pass through the blood-brain barrier. As a result, the Bacosides must be transformed such that the sugar units are eliminated so as to mediate enhanced memory and cognitive activity (31, 32). In comparison to the parent Bacosides, aglycones (jujubogenin and pseudo-jujubogenin) along with their aglycone derivatives such as Ebelin lactone have projected higher binding affinity to all CNS receptors along with improved docking to AChE in insilico models. Higher affinity to CNS receptors results in increased CNS active properties. This offers excellent oral absorption and BBB penetration of *B. monnieri*. In comparison to other ligands, Ebelin lactone also displayed high CNS receptor binding and increased BBB penetration (32). The mixture of Bacosides with a high concentration of bacoside A3, jujubogenin, and an Ebelin lactone of an aglycone derivative has demonstrated acetylcholinesterase inhibition activity, in addition to anti-oxidant, anti-inflammatory, anti-cancer, and anti-cytotoxic characteristics.

4. CONCLUSION

Our findings contribute to the better scientific evidence demonstrating the bioactivity of Bacopa Composition 1 (1:1) which is a mixture of Bacosides with a high concentration of bacoside A3, jujubogenin, and an aglycone derivative, Ebelin lactone. In comparison to the parent Bacosides, our study is unique in the aspect that it focuses on making *B. monnieri* more accessible so that it may cross the blood-brain barrier (BBB) and be absorbed more readily from the intestines. Furthermore, Bacopa Composition 1 (1:1) has the additional benefit that the aglycone derivative components themselves are bioactive, as they have high free radical scavenging activity, acetylcholinesterase inhibition activity, anti-inflammatory activity, high absorption, and no toxicity. Thus, it synergistically enhances the bioactivity of Bacosides from *B. monnieri*.

The findings of such research will provide a better indication of *B. monnieri*'s potential for treating Alzheimer's disease. Furthermore, larger, longer-term trials comparing *B. monnieri* to currently available conventional medications are needed to evaluate whether *B. monnieri* is a viable alternative therapy in the treatment of various human illnesses.

REFERENCES


1. Davidson-Hunt I. Ecological ethnobotany: stumbling toward new practices and paradigms. *MASA J.* 2000;16(1):1-3.
2. Jeyasri R, Muthuramalingam P, Suba V, Ramesh M, Chen JT. Bacopa monnieri and their bioactive compounds inferred multi-target treatment strategy for neurological diseases: A cheminformatics and system pharmacology approach. *Biomolecules.* 2020;10(4):536.
3. Indian Herbal Pharmacopoeia. Indian Drug Manufacturers Association (IDMA). Food and Drug Administration of Maharashtra, Mumbai; 2002.
4. Brimson JM, Brimson S, Prasanth MI, Thitilertdecha P, Malar DS, Tencomnao T. The effectiveness of Bacopa monnieri (Linn.) Wettst. as a nootropic, neuroprotective, or antidepressant supplement: analysis of the available clinical data. *Scientific reports.* 2021;11(1):596.
5. Mukherjee GD, Dey CD. Clinical trial on Brahmi. I. *Journal of experimental medical sciences.* 1966;10(1):5-11.


6. Calabrese C, Gregory WL, Leo M, Kraemer D, Bone K, Oken B. Effects of a standardized Bacopa monnieri extract on cognitive performance, anxiety, and depression in the elderly: a randomized, double-blind, placebo-controlled trial. *The journal of alternative and complementary medicine*. 2008;14(6):707-713.
7. Stough C, Singh H, Zangara A. Mechanisms, efficacy, and safety of Bacopa monnieri (Brahmi) for cognitive and brain enhancement. *Evidence-Based Complementary and Alternative Medicine*. 2015;2015.
8. Devishree RA, Kumar S, Jain AR. Short term effect of Bacopa monnieri on memory—A brief review. *J. Pharm. Res.* 2017;11:1447-1450.
9. Dhawan BN, Singh HK. Pharmacological studies on Bacopa monniera, an Ayurvedic nootropic agent. *European Neuropsychopharmacology*. 1996;6:144.
10. Singh HK, Dhawan BN. Neuropsychopharmacological effects of the Ayurvedic nootropic Bacopa monniera Linn.(Brahmi). *Indian Journal of Pharmacology*. 1997;29(5):359.
11. Simpson T, Pase M, Stough C. Bacopa monnieri as an antioxidant therapy to reduce oxidative stress in the aging brain. *Evidence-based complementary and alternative medicine*. 2015;2015.
12. Deb DD, Kapoor P, Dighe RP, Padmaja R, Anand MS, D'souza P, Deepak M, Murali B, Agarwal A. In vitro safety evaluation and anticlastogenic effect of BacoMind™ on human lymphocytes. *Biomedical and Environmental Sciences*. 2008;21(1):7-23.
13. Ghosh S, Khanam R, Chowdhury AA. The evolving roles of Bacopa monnieri as potential anti-cancer agent: a review. *Nutrition and Cancer*. 2021;73(11-12):2166-2176.
14. Palethorpe HM, Smith E, Tomita Y, Nakhjavani M, Yool AJ, Price TJ, Young JP, Townsend AR, Hardingham JE. Bacopasides I and II act in synergy to inhibit the growth, migration and invasion of breast cancer cell lines. *Molecules*. 2019;24(19):3539.

15. Pajouhesh H, Lenz GR. Medicinal chemical properties of successful central nervous system drugs. *NeuroRx*. 2005;2:541-553.
16. Hansch C, Björkroth JP, Leo A. Hydrophobicity and central nervous system agents: on the principle of minimal hydrophobicity in drug design. *Journal of pharmaceutical sciences*. 1987;76(9):663-687.
17. Egan WJ, Merz KM, Baldwin JJ. Prediction of drug absorption using multivariate statistics. *Journal of medicinal chemistry*. 2000;43(21):3867-3877.
18. Egan WJ, Lauri G. Prediction of intestinal permeability. *Advanced drug delivery reviews*. 2002;54(3):273-289.
19. Hidalgo IJ, Raub TJ, Borchardt RT. Characterization of the human colon carcinoma cell line (Caco-2) as a model system for intestinal epithelial permeability. *Gastroenterology*. 1989;96(2):736-749.
20. Artursson P, Palm K, Luthman K. Caco-2 monolayers in experimental and theoretical predictions of drug transport. *Advanced drug delivery reviews*. 2001;46(1-3):27-43.
21. Van Breemen RB, Li Y. Caco-2 cell permeability assays to measure drug absorption. *Expert opinion on drug metabolism & toxicology*. 2005;1(2):175-185.
22. Hubatsch I, Ragnarsson EGE, Artursson P. Determination of drug permeability and prediction of drug absorption in Caco-2 monolayers. *Nature protocols*. 2007;2(9):2111-2119.
23. Miao Y, He N, Zhu JJ. History and new developments of assays for cholinesterase activity and inhibition. *Chemical reviews*. 2010;110(9):5216-5234.

24. Saija A, Tomaino A, Lo Cascio R, Rapisarda P, Dederen JC. In vitro antioxidant activity and in vivo photoprotective effect of a red orange extract. *International Journal of Cosmetic Science*. 1998;20(6):331-342.
25. Dontha S. A review on antioxidant methods. *Asian J. Pharm. Clin. Res*. 2016;9(2):14-32.
26. Giustarini D, Dalle-Donne I, Milzani A, Fanti P, Rossi R. Analysis of GSH and GSSG after derivatization with N-ethylmaleimide. *Nature protocols*. 2013;8(9):1660-1669.
27. Hawkey CJ. COX-1 and COX-2 inhibitors. *Best Practice & Research Clinical Gastroenterology*. 2001;15(5):801-820.
28. Mallick MN, Akhtar MS, Najm MZ, Tamboli ET, Ahmad S, Husain SA. Evaluation of anticancer potential of *Bacopa monnieri* L. against MCF-7 and MDA-MB 231 cell line. *Journal of pharmacy & bioallied sciences*. 2015;7(4):325.
29. Dhawan BN, Singh HK. Pharmacological studies on *Bacopa monniera*, an Ayurvedic nootropic agent. *European Neuropsychopharmacology*. 1996;6:144.
30. Singh HK, Dhawan BN. Neuropsychopharmacological effects of the Ayurvedic nootropic *Bacopa monniera* Linn. (Brahmi). *Indian Journal of Pharmacology*. 1997;29(5):359.
31. Hansch C, Björkroth JP, Leo A. Hydrophobicity and central nervous system agents: on the principle of minimal hydrophobicity in drug design. *Journal of pharmaceutical sciences*. 1987;76(9):663-687.
32. Ramasamy S, Chin SP, Sukumaran SD, Buckle MJC, Kiew LV, Chung LY. In silico and in vitro analysis of bacoside A aglycones and its derivatives as the constituents responsible for the cognitive effects of *Bacopa monnieri*. *PLoS One*. 2015;10(5):e0126565.

Figure 1: Graph representing the Intestinal absorption of various Bacopa Herbal Compositions (A) Bacoside 20%, (B) Bacoside 20%, (C) Enriched aglycone derivative Ebelin lactone, (D) Composition 1 (1:1), and (E) Composition 2 (3:1) were tested for intestinal absorption.





A-B- Apical to Basolateral transport,
 B-A- Basolateral to Apical transport

Figure 2: Graph representing the percentage of Acetylcholinesterase inhibition for various Bacopa Herbal Compositions




Figure 3: Graph representing the percentage of Radical scavenging activity for various




Figure 4: Graph representing the percentage of Glutathione Reduction for various Bacopa Herbal Compositions




Figure 5: Graph representing the percentage Cox-2 Inhibition for various Bacopa Herbal Compositions




Figure 6: Graph illustrating the percentage inhibition of proliferation of MBA-MB-468 cells





Figure 7: Graph illustrating the percentage inhibition of proliferation of HCT116 cells







Figure 8: Graph illustrating the percentage inhibition of proliferation of Hek293T cells






Figure 9: Graph illustrating the percentage inhibition of proliferation of HFF-1 cells




Table 1: Interpretation of the Intestinal absorption of Bacoside 20%, Bacoside 50%, Enriched aglycone derivative Ebelin lactone, Composition 1 (1:1), and Composition 2 (3:1) in Caco-2 cells

Papp X 10⁻⁶ (cm/s)			
Drugs	Concentration (mg/ml)	Apical to Basolateral transport	Basolateral to Apical transport
Bacoside 20%	0.125	101 ± 27.3	62.4 ± 5.3
Bacoside 50%	0.125	113 ± 13.5	74.4 ± 9.6
Enriched aglycone derivative Ebelin lactone	0.125	121 ± 11.3	78.4 ± 8.4
Composition 1 (1:1)	0.125	141 ± 18.7	77.4 ± 6.2
Composition 2 (3:1)	0.125	133 ± 26.3	74.4 ± 10.2

UNDER PEER REVIEW