

Comparative efficacy of bio active plant extracts and insecticides against cereal leaf beetle (*Oulema melanopus* L.) infesting oats in North Kashmir, India

ABSTRACT.

Experiment was carried out at the farmers field near the Faculty of Agriculture, Wadura, Kashmir, India during 2021-22 under free choice conditions to evaluate efficacy of various insecticides and botanicals against cereal leaf beetle (*Oulema melanopus*) infesting Rabi Oats (*Avena sativa* L.) in North Kashmir. Among plant extracts and insecticides evaluated against *O. melanopus* are concerned, Abamectin proved to be most effective (24, 21.3, 8.3, 1.6 and 1 %) after 2,4,6,8 and 10 days of spray, respectively during 2021 followed by Thiamethoxam (32.6, 15.3, 8, 0.3%) after 2, 4, 6, 8 and 10 days of spray. Same trend of results was observed in 2022 and the pooled data. Among tested botanicals viz. neem and *Artimesia*, *Artimesia* was effective (44, 42, 41, 39 and 37%) after 2, 4, 6,8 and 10 days after spray in managing the pest. Severe infestation has been observed at maturity of crop and keeping in view the use as fodder to cattle it is necessary to inculcate the use of locally available botanicals for better health of cattle during the dearth period of fodder in Kashmir valley. Overall, the marginal farmers might be saved from any untoward by utilization of the bioactive plant extracts in management of this notorious pest.

Key words:Oats, *Oulema melanopus*, mortality, plant extract

1.Introduction

The oats, *Avena sativa* Linnaeus occupies the 6th place among main crops, cropped throughout the world including wheat, maize, rice, barley and sorghum. The main purpose of its cultivation is to obtain source of food for livestock and also for seed because it is known to be nutritionally and medicinally valuable. The grain is now mostly preferred to use because of its rich source of nutrients and beneficial for health. The oats also occupy important place in the biochemical and cosmetic industry. Oats is being used as an important grain crop especially by the people of peripheral populations among the developing countries, and for specifically

important uses in developed world. Oats are grown as a source of grain as well as for source of food for livestock, straw for bedding, hay, haylage, silage and chaff in many parts of the world.

Oats are rated as rich source of protein, dietary soluble fibers and minerals. The oats are believed to contain protein content, equivalent to meat, milk and, egg protein. As food source oats are being most preferably used in breakfast, however it is being considered by consumers as healthiest food from natural source which adds to its global demand. Moreover, oats are preferred over other types of grain because it possess a distinctive nutritional composition and multifunctional benefits of select bioactive compounds, including protein, unsaturated fatty acids, soluble(beta-glucan) and insoluble fibre, micronutrients such as iron, potassium, copper and magnesium, thiamine, folate, zinc and phosphorus (Yu *et al.*, 2012). Oats is also provided with source of many compounds such as tocopherol, inositol hexaphosphate, phenolic compounds, avenanthramides and sterols that bears antioxidant activity. (Brindzova *et al.*,2012). Oats being rich in dietary fibers believed to produce various health benefits with preferably recommended to exhibit hypocholesterolemic and anticancerous properties. In addition to above mentioned properties, it is known to be effective against a number of diseases prevailing in mankind Prasad *et al.* (2015).

The total world production of oats was reported to be 23 million tonnes with major contributions from Russia and Canada, with both producing approximately 20% of the total global production. Other remarkable producers were, Australia and Finland, the United Kingdom and Poland each with over 1 million tonnes (Anonymous a, 2019). In India, the total fodder cultivation constitutes approximately 4.9 (%), which covers an area of approximately 8.6 million ha on individual crop basis. Whereas, oat cultivation occupies approximately about 1.0 million ha in the country with its production of 35-50 t/h (Anonymous, 2019). In India, it is cultivated in Punjab, Haryana, Jammu and Kashmir, Himachal Pradesh, Uttar Pradesh, Madhya Pradesh, Rajasthan, Maharashtra and West Bengal. Uttar Pradesh holds maximum share of oat production with - 34 per cent, followed by Punjab - 20 per cent, Bihar -16 per cent, Haryana - 9 per cent and Madhya Pradesh - 6 per cent (Panday and Roy, 2011).

Besides diseases, A number of insect pests has been reported so far on oats which lead to considerable losses in green fodder and oat meal as well. Fifteen (15) insect pests, belonging to 5 different orders were recorded in the oats (*Avena sativa* L.) Kashmir agro-ecosystem. Soil

borne insects viz. white grubs (*Holotrichialongipennis*, *Brahmina* spp, *Melolontha furcicauda* and *Oryctes* spp), wireworm (*Agriotes* spp) and cutworm (*Agrotis* spp); sucking pests including wheat aphid (*Schizaphisgraminum*), green stink bug (*Carpocoris* spp), aster leafhoppers (*Macrostesquadrillineatus*), broad-headed bug (*Camptopuslateralis*) and oat thrips (*Stenothripsgraminum*) and foliage pests including cereal leaf beetle (*Oulema melanopus* L.), surface grasshopper (*Chrotogonus trachypterus* Blanchard), armyworm (*Mythimnaseparata*) and gram pod borer (*Helicoverpa armigera*) appeared as pests in the oats field. Among these *M. separata*, *O. melanopus* and *H. armigera* were major pests (Kumar *et al.*, 2019). The initial appearance of cereal leaf beetle was recorded in Southern Michigan of the United States in the beginning of 1960s but now have occupied major area of North America. This beetle is characterized to produce one generation in a year. The Adult members of this beetle feed before winter and stay much of over wintering time in protected areas such as wind rows, crop stubble and tree bark crevices. Adults perform mating once the temperature reaches optimum levels of above 9-10 degrees Celsius (or 44.6 degrees F) and females have a comparably long laying period of approximately 45-60 days. They eggs are laid on the undersurface of leaves. The larvae make appearance in 7-15 days, and begin feeding on leaves with marked stripes along the leaf. The larvae took 12-20 days to attain maturity. Larvae poses serious threat to crops, as they consume upper surface of the leaves and make them hindered and unable to photosynthesize. Pupation takes place inside soil and took 20-25 days for adults to emerge, to repeat the cycle once again. Keeping in view the above facts it becomes imperative to study the “Seasonal Incidence of *Oulema melanopus* Infesting Rabi Oats (*Avena sativa* L.) in North Kashmir”.

2. Materials and methods

An experiment was carried out to assess the efficacy of different insecticides and botanicals against the *Oulema melanopus* on oats crop at farmers field near Faculty of Agriculture, Wadura, SKUAST- Kashmir during 2021-22. Variety Sabzar was sown on 12th of

November 2020 for this purpose. The experiment was laid out in a Randomized Complete Block Design with seven treatments and three replications, in a field measuring about plot size was 3x4m² (Total=24 plot) and spacing 22.5cm.

2.1 Green leaf Extraction of *Artemisia annua*

Leaves of *Artemisia annua* were collected, washed with water to remove dirt and chopped into small pieces, then grinded using clean water in the ratio of 50 gm plant material in one litre of water to get 5% solution.

The treatments were imposed using a Knapsack sprayer. The death of larvae population per m² was recorded after spray at 2, 4, 6, 8, and 10 days after treatment. Per cent mortality of larvae over negative control was calculated. Data collected from the experimental site was subjected to standard statistical procedure using standard statistical procedures. Insecticides and botanicals were applied in the evening to escape the effect of sunlight on chemicals. Per cent mortality was calculated as:

$$\text{Per cent mortality} = \frac{\text{Pre count} - \text{Post count}}{\text{Post count}} \times 100$$

Whereas,

Pre count = Number of live larvae before treatments

Post count = Number of live larvae after treatment

Similarly, increase mortality over control was calculated as:

$$\text{Percentage Increase over control} = \frac{\text{Treated} - \text{control}}{\text{Treated}} \times 100$$

3. Results and Discussion

Field efficacy of different plant extracts and insecticides was evaluated against *O. melanopus* infesting oats under field conditions during 2021 and 2022. During 2021 it was evident from the data that among different insecticides evaluated after two days of spray,

Abamectin 1.8EC was found effective (24%) in managing this pest population. Abamectin was followed by Chlorpyrifos 20EC (30.6%), Dimethoate 30EC (32%). After 4 days of spray Thiamethoxam 25 WG proved to be most effective in managing *O. melanopus* by (15.3%) followed by Abamectin 1.8 EC (21.3%). After 6 days of spray Thiamethoxam 25 WG significantly proved to be most effective with (8.0%) in rapid declining the pest population with closely followed by Abamectin 1.8EC (8.3%). After 8 days of spray both Thiamethoxam 25 WG and Abamectin 1.8EC resulted equally effective against pest population with (1.6%) followed by Chlorpyrifos 20EC with (4.6%). After 10 days of spray Thiamethoxam 25 WG dominate the efficacy level with (0.3%) followed by Abamectin 1.8EC with (1.3%).

However, among the bioactive plant extracts evaluated against *O. melanopus* infesting oats, Artemisia GLE @5% was found most effective in overall mean per cent efficiency in causing mortality with (30.42%, 31.61%, 32.80%, 35.19% and 37.57%) after 2, 4, 6, 8 and 10 days of spray followed by Neem oil @5%.

During 2022 it is evident from the studies that among different insecticides evaluated after 2 days of spray, Abamectin 1.8EC was significantly found effective (25.3%) in managing this pest population. Abamectin was followed by Chlorpyrifos 20EC (30.6) and Thiamethoxam 25 WG (32.3%). After 4 days of spray Thiamethoxam 25 WG proved to be most effective in managing *O. melanopus* by (14.3%) followed by Abamectin 1.8 EC (22.0%). After 6 days of spray Thiamethoxam 25 WG significantly proved to be most effective with (8.0%) in rapid declining the pest population with closely followed by Abamectin 1.8EC (8.3%). After 8 days of spray both Thiamethoxam 25 WG and Abamectin 1.8EC resulted equally effective against pest population with (1.6%) followed by Chlorpyrifos 20EC with (3.6%). After 10 days of spray Thiamethoxam 25 WG dominate the efficacy level with (0.3%) followed by Abamectin 1.8EC with (0.6%).

Among the botanicals evaluated against *O. melanopus* infesting oats, Artemisia GLE @5% was found most effective with (34.59%, 39.99%, 41.15%, 43.47 and 45.78%) after 2, 4, 6, 8 and 10 days of spray followed closely by Neem oil @5%.

In pooled data of 2021 and 2022 Abamectin 1.8EC proved most effective (24.6, 21.6, 8.3, 1.6 and 1 %) after 2, 4, 6, 8 and 10 days after spray, respectively. It was followed by

Thiamethoxam 25WG by (32.5, 14.8, 8, 1.6 and 1%) after 2, 4, 6, 8 and 10 days after spray, respectively.


Among two botanicals tested *Artemisia* was found more effective with (44.83, 42.00, 41.00, 39.00, 37.00%) after 2, 4, 6, 8 and 10 days of spray respectively. Although, Laznik *et al.* (2010) described the activity of entomopathogenic nematodes (*Rhabditida*) against cereal leaf beetle (*O. melanopus* [L.], Coleoptera, Chrysomelidae). In this course of study, the three Slovenian strains of entomopathogenic nematodes) and commercial product Entonem (active ingredient of *S. feltiae*), were brought in use to measure their activity against cereal leaf beetle. The most effective strain and good alternative to chemical insecticides with greater tendency to control overwintered cereal leaf beetles was *S. carpocapsae* C101. The efficiency of entomopathogenic nematode was found greatly dependent on temperature. Similarly, Skuhrovec *et al.* (2018) tested the efficiency of plant-derived essential oils and their encapsulations for protection of wheat from adults and larvae of cereal leaf beetle, *O. melanopus*. The two formulations extracted from plants were *Rosmarinus officinalis* with *Cymbopogon citratus* and *Pelargonium graveolens* with *Thymus vulgaris*. These two formulations of essential oil and encapsulation showed 100 per cent mortality within 24 hrs.

Furthermore, flag leaf defoliation causes more damage than does injury to lower leaves. These results are consistent with current knowledge of wheat physiology that the 3-wk period before anthesis is the most critical phase for determining grain yield, in the form of number of seeds per acre (Frederick and Bauer 1999). Cereal leaf beetle eggs hatch over an extended period, which once led to recommendations to wait until most of the cereal leaf beetle eggs had hatched before applying an insecticide (Wilson *et al.* 1969, McPherson 1983). However, considerable defoliation can occur before most of the eggs have hatched (Ihrig *et al.* 2001, Herbert and van Duyn 1999). Detailed studies in Virginia and North Carolina show a close relationship between egg populations and number of fourth-stage larvae per stem (Ihrig *et al.* 2001). Phosphate insecticides generally were not effective when applied before most eggs had hatched. Lambda Cyhalothrin provides the option of using a threshold based on cereal leaf beetle eggs per stem, and applying the insecticide early before or near 50% egg hatch.

Table 1: Efficacy of various botanicals and insecticides against cereal leaf beetle (*Oulema melanopus*)

Treatment	Pre-count Year		2DAS			4DAS			6DAS			8DAS			10DAS		
	2021	2022	2021	2022	Pooled	2021	2022	Pooled	2021	2022	Pooled	2021	2022	Pooled	2021	2022	Pooled
Chlorpyrifos 20EC @ 0.02%	85.00	88.66	30.66 (33.60)	30.66 (33.61)	30.66	21.66 (27.72)	22.33 (28.18)	22.00	9.33 (17.77)	9.66 (18.10)	9.50	4.66 (12.45)	3.66 (11.01)	4.16	2.00 (7.94)	1.66 (7.33)	1.83
Dimethoate 30EC @ 0.03%	82.66	85.33	32.00 (34.43)	33.66 (35.45)	32.83	24.66 (29.75)	25.00 (29.98)	24.83	17.00 (24.33)	17.33 (24.59)	17.16	5.00 (12.87)	5.00 (12.87)	5.00	2.33 (8.74)	2.00 (7.94)	2.16
Thiamethoxam 25WG @ 0.01%	84.00	87.00	32.66 (34.84)	32.33 (34.64)	32.50	15.33 (23.02)	14.33 (22.23)	14.83	8.00 (16.40)	8.00 (16.40)	8.00	1.66 (7.33)	1.66 (7.33)	1.66	0.33 (1.91)	0.33 (1.91)	0.33
Abamectin 1.8EC @0.09%	83.33	86.00	24.00 (29.31)	25.33 (30.20)	24.66	21.33 (27.48)	22.00 (27.95)	21.66	8.33 (16.76)	8.33 (16.76)	8.33	1.66 (7.33)	1.66 (7.33)	1.66	1.33 (6.53)	0.66 (3.82)	1.00
Artemisia GLE @ 5%	84.00	86.33	45.00 (42.11)	44.66 (41.92)	44.83	44.00 (41.53)	40.00 (39.21)	42.00	43.00 (40.95)	39.00 (38.62)	41.00	41.00 (39.79)	37.00 (37.44)	39.00	39.00 (38.62)	35.00 (36.25)	37.00
Neem oil @ 5%	84.66	86.00	47.33 (43.45)	46.33 (42.88)	46.83	45.66 (42.49)	40.33 (39.40)	42.99	45.33 (42.30)	38.66 (38.41)	41.99	43.00 (40.95)	37.00 (37.44)	40.00	40.00 (39.21)	36.00 (36.85)	38.00
Control	84.33	87.66	83.33 (65.93)	82.66 (65.39)	83.00	80.00 (63.42)	80.00 (63.41)	80.00	76.33 (60.87)	76.00 (60.64)	76.16	73.66 (59.10)	74.66 (59.75)	74.16	67.66 (55.33)	68.33 (55.73)	68.00
C.D.	N/A	N/A	2.03	1.71		2.15	1.66		1.67	2.04		2.37	2.24		3.37	3.62	
SE(m)	1.09	1.63	0.65	0.55		0.69	0.53		0.53	0.65		0.76	0.72		1.08	1.16	
C.V.	2.86	4.28	2.78	2.35		3.27	2.59		2.96	3.71		5.13	5.04		8.28	9.40	

Fig:1 -Efficacy of various botanicals and insecticides against cereal leaf beetle,*Oulema melanopus*



Percentage of larvae mortality by different insecticides and plant bio extracts

Comparative efficacy of different plant extracts and chemical insecticides was evaluated to assess the mortality of *Oulema melanopus* larvae infesting oats under field conditions during 2021 and 2022. During 2021 it was evident from the data that among different insecticides evaluated after two days of spray, Abamectin 1.8EC was found effective and caused (54.52%) mortality. Abamectin was followed by Chlorpyrifos 20EC resulted in causing mortality upto (48.92%), Thiamethoxam 25 WG with (45.11%). After 4 days of spray Thiamethoxam 25 WG proved to be most effective in causing huge mortality of *O. melanopus* larvae by (65.75%) followed by Chlorpyrifos 20EC (59.51%). After 6 days of spray Thiamethoxam 25 WG significantly proved to be most effective with (74.47%) in rapid mortality of the pest population with closely followed by Chlorpyrifos 20EC (74.02%). After 8 days of spray Thiamethoxam 25 WG showed (82.02%) mortality followed by Abamectin 1.8EC with (81.33%) mortality. After 10 days of spray Thiamethoxam 25 WG dominate the efficacy level with (83.60%) followed by Chlorpyrifos 20EC with (82.64%).

However, among the bioactive plant extracts evaluated against *O. melanopus* infesting oats, Artemisia GLE @5% was found most effective in overall mean per cent efficiency in causing mortality with (30.42%, 31.61%, 32.80%, 35.19% and 37.57%) after 2, 4, 6, 8 and 10 days of spray followed by Neem oil @5% as mentioned in the table below.


During 2022 it is evident from the studies that among different insecticides evaluated after 2 days of spray, Abamectin 1.8EC was significantly found effective in causing (56.61%) mortality. Abamectin was followed by Chlorpyrifos 20EC (54.07%) and Thiamethoxam 25 WG (49.08%). After 4 days of spray Thiamethoxam 25 WG proved to be most effective in causing most mortality of *O. melanopus* by (70.52%) followed by Chlorpyrifos 20EC (63.47%). After 6 days of spray Thiamethoxam 25 WG significantly proved to be most effective with (77.80%) mortality with closely followed by Chlorpyrifos 20EC (77.76%). After 8 days of spray Thiamethoxam 25 WG resulted effective with (85.09%) mortality followed by Chlorpyrifos 20EC with (84.53%). After 10 days of spray Chlorpyrifos 20EC dominate the efficacy level with (86.78%) with closely followed by Thiamethoxam 25 WG with (0.6%) larval mortality.

Among the botanicals evaluated against *O. melanopus* infesting oats, Artemisia GLE @5% was found most effective with (34.59%, 39.99%, 41.15%, 43.47 and 45.78%) after 2, 4,6,8 and 10 days of spray followed closely by Neem oil @5% as mentioned in the table below.

Table:2 -Percentage of larvae mortality count

Treatment	2DAS		4DAS		6DAS		8DAS		10DAS	
	2021	2022	2021	2022	2021	2022	2021	2022	2021	2022
Chlorpyriphos 20EC @ 0.02%	48.92	54.07	59.51	63.47	74.02	77.76	79.51	84.53	82.64	86.78
Dimethoate 30EC @ 0.03%	43.94	45.88	52.82	56.03	62.09	65.02	76.61	79.47	79.84	82.98
Thiamethoxam 25WG @ 0.01%	45.11	49.83	65.75	70.52	74.47	77.80	82.02	85.09	83.60	86.62
Abamectin 1.8EC @0.09%	54.52	56.64	57.73	60.41	73.33	76.31	81.33	84.06	81.73	85.23
Artemisia GLE @ 5%	30.42	34.59	31.61	39.99	32.80	41.15	35.19	43.47	37.57	45.78
Neem oil @ 5%	28.75	32.12	30.72	39.10	31.11	41.04	33.86	42.97	37.41	44.13

Fig:2 -Percentage of larvae mortality count



4. Conclusion

In reference to field efficacy of different plant extracts and insecticides evaluated against *O. melanopus* Abamectin proved to be most effective (24, 21.3, 8.3, 1.6 and 1 %) after 2,4,6,8 and 10 days of spray, respectively during 2021 followed by Thiamethoxam (32.6, 15.3, 8, 0.3%) after 2, 4, 6, 8 ad 10 days of spray. Same trend of results was observed in 2022 and the pooled data.

Among two tested botanicals viz. neem and *Artimesia*, *Artimesia* was effective with (44.83%, 42.00%, 41.00%, 39.00% and 37.00%) after 2, 4, 6,8 and 10 days after spray in managing the pest.

Infestation of cereal leaf beetle has been at alarm in the area, hence keen attention is to be taken to devise best possible management strategy. Local botanical *Artimesia* along with other organic approaches can be utilized in formulating the integrated pest management programme to get rid of cereal leaf beetle for boosting the farmers income, minimizing environmental hazards especially for the cattle whom oats act as a feed.

5. References

1. Anonymous. Ministry of Agriculture, Govt. of India. 2019 www.indiastat.com.
2. Brindzova L, Certik M, Rapta P, Zalibera M, Mikulajova A, Takacssova M. Antioxidant activity, Beta-glucan and lipid contents of Oats Variety. *Journal of Food Science*. 2012; 26(3): 163-173.
3. Frederick JR, Bauer PJ. Physiological and numerical components of wheat yield, pp. 45-65. In G. A. Slafer and E. H. Satorre (eds.), *Wheat ecology and physiology of yield determination*. Food Products Press, An Imprint of The Haworth Press, New York. 1999.
4. Herbert DA Jr, Van Duyn JW. *Cereal leaf beetle biology and management*. Virginia Cooperative Extension Publication No. 1999; 444-350, Blacksburg, VA.
5. Ihrig RA, Herbert DA, Van Duyn JW, Bradley JR. Relationship between cereal leaf beetle (Coleoptera: Chrysomelidae) egg and fourth-instar populations and impact of fourth-instar defoliation of winter wheat yields in North Carolina and Virginia. *Journal of Economic Entomology* 2001; 94: 634-639
6. Kumar KK, Sridhar J, Ramasamy K, Murali-Baskaran, Sengottayan. Microbial biopesticides for insect pest management in India: Current status and future prospectus. *Journal of invertebrate Pathology*. 2019; 165: 74-81.
7. McPherson RM. Seasonal abundance of cereal leaf beetles (Coleoptera: Chrysomelidae) in Virginia small grains and corn. *Journal of Economic Entomology*. 1983; 76(6): 1269-1272.
8. Pandey KC, Roy AK. *Forage Crops Varieties*. 2011. IGFRI Jhansi (India)
9. Prasad R, Alok J, Latha S, Arvind K, Unnikrishnan VS. Nutritional advantages of oats and opportunities for its processing as a value-added food—a review. *Journal of food science and technology*. 2015; 52: 662-675.
10. Skuhrovec J, Douda O, Zouhar M, Manasova M, Novak P, Bozik M. Insecticidal activity of two formulations of essential oils against the cereal leaf beetle. *Acta Agriculturae Scandinavica, Section B- Soil and Plant Science*. 2018; 68(6): 489-495.
11. Wilson MC, Treece RE, Shade RE, Day KM, Stivers KM. Impact of cereal leaf beetle larvae on yields of oats. *Journal of Economic Entomology*. 1969; 62: 699-702.
12. Yu L, Ront T, Shahidi. *Cereals and Pulses Nutraceutical Properties and Health Benefit*. 1st edition, USA, Wiley-Blackwell. 2012; 50-54.