

## Original Research Article

### Effect of fungicides on Mycelium growth of *Rhizoctonia solani* of rice under *in-vitro*.

#### ABSTRACT

The experiment was conducted at Department of plant Pathology, College of Agriculture, Rewa (M.P.) during 2018-19. The experiment was laid out in a complete randomized design (CRD) with eight treatments including untreated (control) and replicated four. Seven fungicides viz., Flusilazole 12.5%+carbendazim 25% SC, Azoxystrobin 18.2% w/w+difenoconazole 11.4 w/w SC, Azoxystrobin 11%+Tebuconazole 18.3% w/w SC, Tricyclazole 18%+ mancozeb 62% WP, Zineb 68% + hexaconazole 4% WP, Trifloxystrobin 25%+Tebuconazole 50% WG, Mancozeb 50%+carbendazim 25% WS, were evaluate against *R. solani* of rice at different concentrations under *in-vitro* by poisoned food technique. The present study it could be concluded that among the tested fungicides, Azoxystrobin 11%+Tebuconazole 18.3% w/w SC at 200 ppm and 100 ppm was found significantly superior in inhibiting the mycelial growth of *R. solani* over untreated check at 96 hrs. after incubation.

**Key words:** *Oryza sativa*, Sheath blight, Fungicides and Mycelial growth

#### INTRODUCTION

Rice (*Oryza sativa* L.) is one of the most important cereal crops and a major food grain contributor to the total world food grain basket. It is one of the most important staple food crop grown in different ecology zones and contributes 40% of total food grain production in India. Productivity of rice can be increased by adopting the hybrid rice, integrated nutrient and pest management for combating the economic losses due to biotic stresses. In India, area 46.38 Mha with production 130.29 Mt and productivity 2809 kg/ha (Anon, 2022). Sheath Blight disease of rice caused by *Rhizoctonia solani* Kühn is one of the most important and widely distributed diseases in all the rice growing regions of the world and causing considerable loss in grain yield of 54.3 per cent. According to Chahal *et al.* (2003). the estimation of losses due to sheath blight of rice in India has been reported up to 54.3 per cent. The natural infection of the sheath blight disease occurs at the seedling, tillering and booting stages of rice. Infection usually starts near the water line of rice plants in paddy fields. Lesions develop upward to the upper leaf sheaths and leaf blades. The centre of lesion become grayishwhite with brown margin, later several spots coalesce and show blight symptoms (Ou,

**Comment [MT1]:** This experiment not the experiment. Give an opening sentence and also the aim of the study before methodology;

**Comment [MT2]:**

**Comment [MT3]:** rephrase this sentence

**Comment [MT4]:** Your language is not clear. write the sentence clearly

**Comment [MT5]:** Rephrase this sentence and also conclude your findings

**Comment [MT6]:** Write clearly what you want to say or advised to removed this sentence

**Comment [MT7]:** I suggest this sentence can be rephrase by adding this'

**Comment [MT8]:** removed the

**Comment [MT9]:** sentence not clear rephrase

**Comment [MT10]:** add this

**Comment [MT11]:** correct

**Comment [MT12]:** correct

**Comment [MT13]:** correct

**Comment [MT14]:** remove this

1985). When humidity exceeds 95 % and temperature ranges from 29 to 32<sup>0</sup>c, infection spreads rapidly which appear on plant parts, including leaf blades, causing irregularly shaped lesions with brown borders as bands. This symptom generally referred as “banded blight”. *Rhizoctonia solani* possess pale to dark brown rapidly growing mycelium with septum in the branch near the point of origin. It produces large number of globose, sclerotia which initially turn white, later turn brown to purplish brown. Fungicide application is the most common approach among the farmers for the management of sheath blight throughout the world. The complex genetic nature of resistance to sheath blight and genetic variability of the pathogen increases the difficulty in developing resistant host genotypes, as well as in effectively deploying available tolerant cultivars. Unfortunately, at present there is no known rice varieties which is either immune or possess high degree of resistance to sheath blight disease. In the absence of suitable resistant donors, fungicides are the main answer to check these diseases. The present study was undertaken to evaluate the different fungicides for control of sheath blight of rice.

#### MATERIALS AND METHODS

The experiment was laid out in a complete randomized design (CRD) with eight treatments including untreated control and replicated fours. Seven fungicides viz., Flusilazole 12.5%+carbendazim 25% SC, Azoxystrobin 18.2% w/w+difenoconazole 11.4 w/w SC, Azoxystrobin 11%+Tebuconazole 18.3% w/w SC, Tricyclazole 18%+ mancozeb 62% WP, Zineb 68% + hexaconazole 4% WP, Trifloxystrobin 25%+Tebuconazole 50% WG, Mancozeb 50%+carbendazim 25% WS, were evaluate against *R. solani* of rice at different concentrations. The present study was undertaken in the laboratory conditions to find out their relative efficacy to inhibit the mycelial growth on PDA medium by poisoned food technique (Nene and Thapliyal, 1979). The required quantity of fungicides were added to the PDA medium at luke warm stage and thoroughly mixed before pouring into petri plate so as to get the desired concentration of active ingredient of each fungicides separately. Twenty ml of fungicide- amended medium was poured in each of 90 mm sterilized petri plate and allowed to solidify. After solidification, each plate was inoculated with five mm discs of 3 day old culture of *R. solani* were cut with sterile cork borer and transferred aseptically to the centre of the poisoned medium. Similarly, control plates were maintained by placing 5 mm disc of the pathogen in the centre of the non-poisoned PDA medium. Four replications were maintained in respect of each isolate and each concentration. All the inoculated petriplate were incubated at 27±2 °C in BOD incubator for seven days. The colony diameter was measured when the

**Comment [MT15]:** Who said this avoid plagiarism

**Comment [MT16]:** rephrase this sentence

**Comment [MT17]:** Replace under taken with conducted

**Comment [MT18]:** Replace this with under

**Comment [MT19]:** relative efficacy of what

**Comment [MT20]:** sentence not clear

**Comment [MT21]:** what is the meaning of non poisoned pda

control plates were filled with fungal growth. Per cent inhibition over control was calculated by the following formula suggested by Vincent (1947).

$$I = \frac{C - T}{C} \times 100$$

Where,

I = Per cent inhibition in growth of *R. solani*

C = Mycelial growth (mm) in control

T = Mycelial growth (mm) in treatments

## RESULTS AND DISCUSSION

The data on bio efficacy of fungicides against Sheath blight of rice is presented in Table-1. and Fig-1. The data indicate that among the seven tested fungicides at different concentrations, Azoxystrobin 11% + tebuconazole 18.3% w/w SC at 200ppm was found to be highly effective (0.0mm) followed by Azoxystrobin 18.2% w/w + difenoconazole 11.4% w/w SC (20.2mm) over untreated check (74.0mm) after 96 hrs of incubation. The study also reveals that Azoxystrobin 11% + tebuconazole 18.3% w/w SC at low concentration was even superior to other fungicide for inhibiting the mycelia growth. Thus, it may be summarized that Azoxystrobin 11% + tebuconazole 18.3% w/w SC and Azoxystrobin 18.2% w/w + difenoconazole 11.4% w/w SC at 50ppm, 100ppm and 200ppm drastically inhibited the growth of test fungi. However, it was observed that Trifloxystrobin 25% + tebuconazole 50% WG at 200ppm drastically inhibited the growth of test fungus (23mm) after 96 hrs of incubation whereas, Zineb 68% + hexaconazole 4% WP, Tricyclazole 18% + mancozeb 62% WP, Flusilazole 12.5% + carbendazim 25% SC and Mencozeb 50% + carbendazim 25% WS gave moderate inhibition of mycelial growth 25.6mm, 27.3 mm, 28.5mm and 30.0 mm respectively over untreated check (74.0mm) under study. The other tested concentration 50 ppm and 100 ppm also gave inhibitory effect to check the mycelial growth of test fungus.

Azoxystrobin 11% + tebuconazole 18.3% w/w SC to recorded maximum % inhibition over control at 200 ppm and caused complete inhibition of the *R. solani* followed by Azoxystrobin 18.2% w/w + difenoconazole 11.4% w/w SC (72.2%). Other tested fungicides caused growth inhibition in the range of 59.4 to 68.9% at 200 ppm concentration. At 100 ppm concentration, Azoxystrobin 11% + tebuconazole 18.3% w/w SC caused complete inhibition of growth of the *R. solani* followed by Azoxystrobin 18.2% w/w +

**Comment [MT22]:** I suggest you use most effective not highly

**Comment [MT23]:** using superior is not scientifically use other word and the sentence is not clear

difenoconazole 11.4% w/w SC caused 60.8% inhibition of growth of the *R. solani*. Rest of the fungicides caused growth inhibition in the range of 49.2 to 56.5 at 100 ppm concentration. At lower concentration of 50ppm, 52.0% growth inhibition was recorded with Azoxystrobin 11% + tebuconazole 18.3% w/w SC followed by Azoxystrobin 18.2% w/w + difenoconazole 11.4% w/w SC effected 49.3% growth inhibition. Other fungicides exhibited moderate inhibition. Among the fungicides Mancozeb 50% + carbendazim 25% WS showed least inhibition. The similar results was also reported by Agrawal *et al.*(2012), Srinivas *et al.*(2014) and Mushineni *et al.*(2017) and confirmed the present findings.

Comment [MT24]: discuss your results

## CONCLUSION

It can be concluded that among the tested fungicides, Azoxystrobin 11%+Tebuconazole 18.3% w/w SC at 200 ppm and 100 ppm was found significantly superior in inhibiting the mycelial growth of *R. solani* over untreated check at 96 hrs. after incubation.

## REFERENCES

- Anonymous. Agriculture Statistics at a glance, Ministry of Agriculture. 2022;28.
- Agrawal M and Sunder S. Effect of fungicides and non-conventional chemicals on *Rhizoctonia solani* AG-1 1A and sheath blight disease of rice. Plant Disease Research. 2012; 28 (1): 39-44.
- Chahal K K S, Sokhi SS and Rattan GS. Investigation on sheath blight of rice in Punjab. Indian Phytopathology. 2003; 56(1): 22-26.
- Mushineni AJ, Khalko S and Thapa S. *In vitro* Study of New Generation Chemicals against *Rhizoctonia solani* Kuhn Causing Sheath Blight of Rice. International Journal of Agriculture Sciences. 2017; 19(9):4201-4203.
- Nene Y L and Thapliyal PN. Fungicides in Plant Disease Control, Oxford and IBH Publishing House, New Delhi. 1979; 163.
- Ou SH. Rice Diseases. Commonwealth mycological Institute, Kew, England. 1985;368.
- Srinivas P, Ratan V, Narayan RP and Bindu MG. *In vitro* evaluation of fungicides, bio control agents and plant extracts against rice sheath blight pathogen *Rhizoctonia solani*. International Journal of Applied Biology and Pharmaceutical Technology. 2014; 5:121-126.
- Vincent J M. Distortion of fungal hyphae in presence of certain inhibitors. Nature. 1947; 150-850.

UNDER PEER REVIEW

**Table -1. Mycelial growth of *R. solani* as influenced by different fungicides *in Vitro***

Sl. No.	Treatments	Mycelial growth (mm)			% inhibition over control
		48 hrs	72 hrs	96 hrs	
T1	Flusilazole 12.5% + carbendazim 25% SC	24.25	34.5	47	

					35.6
<b>T2</b>	Azoxystrobin 18.2% w/w + difenoconazole 11.4% w/w SC	21	32	37	49.3
<b>T3</b>	Azoxystrobin 11% + tebuconazole 18.3% w/w SC	18	30	35	52.0
<b>T4</b>	Tricyclazole 18% + mancozeb 62% WP	24	34	45	38.3
<b>T5</b>	Zineb 68% + hexaconazole 4% WP	23.8	33.5	43	41.0
<b>T6</b>	Trifloxystrobin 25% + tebuconazole 50% WG	23	32.9	40	45.2
<b>T7</b>	Mencozeb 50% + carbendazim 25% WS	24.7	35.2	49	32.8
<b>T8</b>	Control	28	45	74	0.0
	<b>CD (5%)</b>	<b>2.226</b>	<b>2.551</b>	<b>2.349</b>	
	<b>Concentration 100 ppm</b>				
<b>T1</b>	Flusilazole 12.5% + carbendazim 25% SC	14.1	27.1	37.2	50.4
<b>T2</b>	Azoxystrobin 18.2% w/w + difenoconazole 11.4% w/w SC	11	22.2	29.4	60.8
<b>T3</b>	Azoxystrobin 11% + tebuconazole 18.3% w/w SC	8	16	0	100
<b>T4</b>	Tricyclazole 18% + mancozeb 62% WP	13.5	26.5	35	53.3
<b>T5</b>	Zineb 68% + hexaconazole 4% WP	13	26.3	34	54.6
<b>T6</b>	Trifloxystrobin 25% + tebuconazole 50% WG	12.4	24.7	32.6	56.5
<b>T7</b>	Mencozeb 50% + carbendazim 25% WS	15.1	29	38	49.2
<b>T8</b>	Control	28	45	74	0.0
	<b>CD (5%)</b>	<b>2.462</b>	<b>3.015</b>	<b>2.418</b>	
	<b>Concentration 200 ppm</b>				
<b>T1</b>	Flusilazole 12.5% + carbendazim 25% SC	9.1	18	28.5	61.5

<b>T2</b>	Azoxystrobin 18.2% w/w + difenoconazole 11.4% w/w SC	6	12.2	20.2	72.2
<b>T3</b>	Azoxystrobin 11% + tebuconazole 18.3% w/w SC	0	0	0	100
<b>T4</b>	Tricyclazole 18% + mancozeb 62% WP	8.8	16.3	27.3	63.1
<b>T5</b>	Zineb 68% + hexaconazole 4% WP	7.6	15.9	25.6	65.4
<b>T6</b>	Trifloxystrobin 25% + tebuconazole 50% WG	7	14	23	68.9
<b>T7</b>	Mencozeb 50% + carbendazim 25% WS	10.6	20	30	59.4
<b>T8</b>	Control	28	45	74	0.0
	<b>CD (5%)</b>	<b>1.764</b>	<b>3.000</b>	<b>2.514</b>	

UNDER PEER REVIEW




Fig-1. Mycelial growth of *R. solani* as influenced by different fungicides *in Vitro*

UNDER PEER REVIEW